

Article

Taphonomic and diagenetic pathways to protein preservation, part II: the case of *Brachylophosaurus canadensis* specimen MOR 2598

Paul V. Ullmann ^{1,*}, Richard D. Ash² and John B. Scannella^{3,4}

¹ Department of Geology, Rowan University, Glassboro NJ 08028, USA

² Department of Geology, University of Maryland, College Park MD 20742, USA; rdash@umd.edu

³ Museum of the Rockies, Montana State University, Bozeman MT 59717, USA; john.scannella@montana.edu

⁴ Department of Earth Sciences, Montana State University, Bozeman MT 59717, USA

* Correspondence: ullmann@rowan.edu

Citation: Ullmann, P.V.; Ash, R.D.; Scannella, J.B. Taphonomic and diagenetic pathways to protein preservation, part II: the case of *Brachylophosaurus canadensis* specimen MOR 2598. *Biology* **2022**, *10*, x. <https://doi.org/xxxxx>

Academic Editor: Firstname Lastname
Received: date
Accepted: date
Published: date

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

Simple Summary: Reports of recovery of proteins and other molecules from fossils have become so common over the last two decades that some paleontologists now focus almost entirely on studying how biologic molecules can persist in fossils. In this study, we explored the fossilization history of a specimen of the hadrosaurid dinosaur *Brachylophosaurus* which was previously shown to preserve original cells, tissues, and structural proteins. Trace element analyses of the tibia of this specimen revealed that after its bones were buried in a brackish estuarine channel, they fossilized under moist conditions which shifted in redox state multiple times. Successful recovery of proteins from this specimen, despite this complex history of chemical alterations, shows that the processes which bind and stabilize biologic molecules shortly after death provide them remarkable physical and chemical resiliency. By uniting our results with those of similar studies on other dinosaur fossils known to also preserve original proteins, we also conclude that exposure to oxidizing conditions in the initial ~48 hours postmortem likely promote molecular stabilization reactions and that retention of early-diagenetic trace element signatures may be a useful proxy for molecular recovery potential.

Abstract: Recent recoveries of peptide sequences from two Cretaceous dinosaur bones require paleontologists to rethink traditional notions about how fossilization occurs. As part of this shifting paradigm, several research groups have recently begun attempting to characterize biomolecular decay and stabilization pathways in diverse paleoenvironmental and diagenetic settings. To advance these efforts, we assessed the taphonomic and geochemical history of *Brachylophosaurus canadensis* specimen MOR 2598, the left femur of which was previously found to retain endogenous cells, tissues, and structural proteins. Combined stratigraphic and trace element data show that after brief fluvial transport, this articulated hind limb was buried in a sandy, likely-brackish, estuarine channel. During early diagenesis, percolating groundwaters stagnated within the bones, forming reducing internal microenvironments. Recent exposure and weathering also caused surficial leaching of trace elements from the specimen. Despite these shifting redox regimes, proteins within the bones were able to survive through diagenesis, attesting to their remarkable resiliency over geologic time. Synthesizing our findings with other recent studies reveals that oxidizing conditions in the initial ~48 hours postmortem likely promote molecular stabilization reactions and that retention of early-diagenetic trace element signatures may be a useful proxy for molecular recovery potential.

Keywords: REE; *Brachylophosaurus*; molecular paleontology; geochemical taphonomy; diagenesis; bone; protein; collagen; Judith River Formation

1. Introduction

1.1. Shifting Views on Molecular Preservation

Fossilization has historically been viewed as a “harsh” process involving forced mineralization coincident with wholesale loss of organic tissues and their component biomolecules (e.g., [1]). However an ever-expanding wealth of recent studies employing methods from histochemistry and immunoassays to genomics and proteomics have shattered this ‘traditional’ paradigm. As reviewed by ourselves and others [2–7], it is now clear that not only is long-term preservation of endogenous DNA, proteins, and other biomolecules possible in fossils, but select burial circumstances may actually promote molecular preservation in both invertebrates and vertebrates, interred in a diverse array of depositional environments. For example, drastic advances in analytical resolution over the last two decades have enabled recovery of nuclear and mitochondrial genomes and/or proteomes from fossils of a number of charismatic Pleistocene mammals, such as cave bears [8], mammoths [9–12], horses [13], saber-toothed cats [14], and ancient hominins [15,16].

Given their greater resilience to decay than DNA [17], structural proteins (e.g., collagen I, actin, β -keratin) are now known from vertebrate fossils dating all the way back to the Jurassic (e.g., [18–22]). Considerable attention has been paid especially to the protein collagen I due to its sheer abundance in bioapatitic tissues [23] and inferred high preservation potential [18,24–47]. Recent studies have employed numerous independent techniques to identify collagen within fossils, with arguably the most convincing evidence being the identification of original peptide sequences and diagenetiforms (protein remains demonstrably modified by diagenetic alterations [48]) in fossil bones via high-resolution tandem mass spectrometry [18,28,30,42,45]. Remarkably, cladistic analyses incorporating collagen I peptides recovered from two Cretaceous nonavian dinosaur bones confirmed their archosaurian identities and thus endogeneity [45,49]. Such findings unequivocally demonstrate both that biomolecules can ‘survive’ fossilization and that portions of them can persist over strikingly-long geologic timescales.

However, the idea of molecular preservation in Mesozoic (and possibly even older) fossils remains controversial to some due to our incomplete understanding of soft-tissue fossilization in general and the geochemical reactions which may stabilize biomolecules within cells and tissues over such immense time frames. While it is universally agreed that processes such as rapid burial can facilitate the preservation of soft tissues in fossils [50,51], it largely remains unclear how other taphonomic processes and the physical (e.g., sedimentology, hydrodynamics) and chemical attributes (e.g., aqueous geochemistry) of depositional environments influence decay at the molecular level (but see [52] for an informative initial foray into this subject). It is therefore vital for researchers to not only demonstrate the authenticity of biomolecular remnants in fossils, but to also identify the physicochemical factors acting within the diagenetic settings which permitted such cases of “exceptional” preservation. Pioneering actualistic studies by Schweitzer et al. [4] and Boatman et al. [53] demonstrated that iron free radicals in diagenetic pore fluids likely play a role by inducing intramolecular crosslinking, but are other aspects of groundwater chemistry (i.e., redox state; cf. [54]) and diffusion history (i.e., duration spent saturated; cf. [47]) equally important in determining whether or not biomolecules persist in fossils? And if they are, which depositional settings and diagenetic histories most favor long-term molecular preservation? In short, we are just beginning to explore these questions.

1.2. Insights from Trace Element Analyses

One of the most effective means of clarifying the geochemical history of a fossil is through studying its trace element composition. After being freed from surrounding sediments by percolating groundwaters, trace element ions, including those of the rare earth elements (REE: lanthanum-lutetium), uranium, and scandium, are ubiquitously

adsorbed by bone hydroxyapatite during diagenesis [55]. Since these elements are essentially absent in bone tissue *in vivo*, their presence in fossil bones derives almost entirely from postmortem interactions with surface and groundwaters [55,56]. As a result, the proportions and spatial distributions of trace elements within a fossil bone provide detailed insights into the chemistries of past pore fluids and the geochemical milieus to which a specimen was exposed throughout its term of burial (e.g., [57–64]). Trace element signatures have thus been successfully utilized to: 1) infer the relative degree of chemical alteration a specimen has endured [65,66]; 2) characterize the chemistry of pore fluids in past environments (e.g., [60,63,67]); 3) track spatiotemporal trends in redox conditions within specimens throughout diagenesis (e.g., [62–64]), and; 4) clarify the number and relative timing of exposures to pore fluids throughout diagenesis (e.g., [64,68]). REE signatures, in particular, have also been shown to potentially be viable proxies for molecular preservation in fossil bones (i.e., they can help identify the most ideal specimens for paleomolecular investigation [47]).

Several of our recent studies capitalized on these diverse utilities of trace elements to elucidate the paleoenvironmental, geochemical, and diagenetic history of two fossils from the Cretaceous Hell Creek Formation which were each documented to preserve endogenous collagen I [28,47], specifically an *Edmontosaurus* fibula [63] and a femur of *Tyrannosaurus rex* specimen Museum of the Rockies (MOR) 1125 [64] (also see [69] for alternative taxonomic assignment of MOR 1125). These studies provided intriguing insights into taphonomic pathways to protein preservation, but they still merely constitute two case studies in the same geologic Formation; the full suite of taphonomic and diagenetic variables at play in molecular preservation remains to be clarified.

In this study, we conducted trace element analyses on the only Mesozoic fossil other than *T. rex* MOR 1125 known to yield endogenous peptide sequences: *Brachylophosaurus canadensis* specimen MOR 2598. Schweitzer et al. [30] and Schroeter et al. [45] each recovered numerous peptides of collagen I from the left femur of this hadrosaur, the authenticity of which were independently corroborated by multiple forms of microscopy, infrared spectroscopy, mass spectrometry, and immunoassays replicated in multiple laboratories by separate researchers each using dedicated equipment and reagents (see each reference for further details). The cumulatively-comprehensive approach undertaken by these studies to demonstrate reproducibility and authenticate the endogeneity of collagen in MOR 2598 set a rigorous standard that has yet to be matched again, despite over a decade of ensuing research on other specimens. Given this great significance of MOR 2598 in providing concrete foundations for the field of molecular paleontology, it is only right to resolve the taphonomic and diagenetic history of this specimen in equally comprehensive detail.

2. Taphonomic and Geologic Context

MOR 2598 consists of an articulated left hind limb of a subadult *Brachylophosaurus canadensis* recovered from an outcrop of the Campanian Judith River Formation north of Malta, MT, on lands managed by the Montana Department of Natural Resources and Conservation (Figure 1). The specimen was found within a thick sequence (~7 m) of channel sandstones exposed along the southern side of Cottonwood Creek. The tibia, fibula, and pes were collected in the summer of 2006, and the femur was collected in a separate plaster jacket the following year. The incomplete nature of the tibia (see below) and slightly lighter color of this bone than the femur indicate that portions of the tibia were exposed by modern erosion/weathering upon discovery (whereas the femur was not [30]). As discussed by Schweitzer et al. [30], the femur was collected with 10–12 cm of sediment still encasing it to maintain geochemical equilibrium for as long as possible before examination by demineralization, scanning electron microscopy, multiple immunoassays, and liquid chromatography-tandem mass spectrometry. All of the skeletal elements appear well-preserved morphologically in that they lack any signs of weathering or abrasion, but the tibia (examined herein) is missing its distal end and a

section of the posterior and medial portions of the shaft near its proximal end. The tibia is also highly fractured, exhibiting numerous transverse and longitudinal fractures arising from compaction after fossilization. Though the medullary cavity of this bone is partly 'filled' due to compaction and partial crushing of cancellous trabeculae, there are no signs of permineralization or infilling by sedimentary matrix (pers. observations, and [70]).




Figure 1. (A) Map showing the locality from which MOR 2598 was recovered in Phillips County, Montana. (B) Left tibia of MOR 2598 examined in this study, shown in lateral view. Map redrawn and modified from ([71], fig. 1).

3. Materials and Methods

3.1. Materials

It was not possible to acquire a sample from the left femur of *Brachylophosaurus canadensis* MOR 2598 for this study as prior histologic and paleomolecular studies of this particular skeletal element of the specimen [30,45,70] and reconstructive efforts undertaken during preparation to maintain permanent stability of the bone left no portion of the cortex easily removable without compromising the integrity of the fossil. Therefore, we instead extracted a fragment of the cortex from the midshaft of the left tibia of MOR 2598. This bone was found in articulation with the left femur in the field, and was accordingly buried at the same stratigraphic position within the same stratum as the left femur, so it was almost certainly exposed to the same environmental conditions postmortem and early-diagenetic regime(s) after burial as the left femur. The cortex at the midshaft of tibiae also possesses similar thickness, density, and histologic microstructure to the midshaft cortex of femora in hadrosaurids (e.g., [72,73]). For these reasons, we are confident that trace element signatures within the left tibia should be very similar to those that would be identified in the left femur examined by prior studies, and that this tibia therefore represents a suitable choice for examining the geochemical history of the hindlimb of MOR 2598 as a whole.

The excised cortical fragment encompasses the majority of the cortical thickness of the bone, including the external margin, but fragile cancellous bone within the medullary cavity disintegrated away during preparation. Because of this, we infer that the innermost portion of the internal cortex is not included in our analyses.

3.2. Methods

3.2.1. Sample Preparation

The cortical sample was embedded in Silmar 41™ resin (US Composites) under vacuum, then sectioned using a Hillquist SF-8 trim saw. The resulting ~3 mm thick section was briefly rinsed with distilled water then briefly polished with 600 grit silicon

carbide to acquire an evenly smooth surface for ensuing laser ablation-inductively coupled plasma mass spectrometry (LA-ICPMS) analyses.

3.2.2. LA-ICPMS Analyses

We employed the same LA-ICPMS methods as Ullmann et al. [63,64] in this study, and refer the reader to the Supplementary Materials and those publications for thorough details. In brief, LA-ICPMS was chosen as a powerful means of examining this spatial distribution of REE and other minor and trace elements within the tibia of MOR 2598, which can provide unique insights into the diagenetic history of a fossil bone and any geochemical shifts it endured through its fossilization. Concentrations of elements are reported in parts per million (ppm) except for iron, which is reported in weight percent (wt. %). REE concentrations were normalized against the North American Shale Composite (NASC; [74,75]) to facilitate comparisons to other fossil bones from other localities. Use of a subscript _N denotes NASC-normalized values and ratios. Reproducibility of our results was taken as the percent relative standard deviation for all REE in the NIST 610 glass standard; it averaged 1.5% and was below 3% for all analyzed elements. NASC-normalized REE ratios were used to calculate $(\text{Ce}/\text{Ce}^*)_N$, $(\text{Ce}/\text{Ce}^{**})_N$, $(\text{Pr}/\text{Pr}^*)_N$, and $(\text{La}/\text{La}^*)_N$ anomalies following Herwartz et al. [60]: $(\text{Ce}/\text{Ce}^*)_N = \text{Ce}_N / (0.5 \cdot \text{La}_N + 0.5 \cdot \text{Pr}_N)$, $(\text{Ce}/\text{Ce}^{**})_N = \text{Ce}_N / (2 \cdot \text{Pr}_N - \text{Nd}_N)$, $(\text{Pr}/\text{Pr}^*)_N = \text{Pr}_N / (0.5 \cdot \text{Ce}_N + 0.5 \cdot \text{Nd}_N)$, and $(\text{La}/\text{La}^*)_N = \text{La}_N / (3 \cdot \text{Pr}_N - 2 \cdot \text{Nd}_N)$.

4. Results

4.1. Overall REE Composition

As a whole (i.e., by summing all transect data), the left tibia of MOR 2598 exhibits a ΣREE value of 256 ppm (Table 1). The three most abundant trace elements in the cortex of the bone are iron (Fe), strontium (Sr), and barium (Ba), which exhibit concentrations of 0.94 wt%, 2,499 ppm, and 1,448 ppm, respectively (Table 1). All of these elements, as well as manganese (Mn), are present in concentrations approximately one to two orders of magnitude higher than REE (Table 1). Whereas average scandium (Sc) enrichment (59 ppm) is around the same magnitude as those of most LREE (~10–90 ppm), the average yttrium (Y) concentration (190 ppm) is more than double that of the highest REE (89 ppm for cerium [Ce]; Table 1). Among REE, there is substantially-greater enrichment in light rare earth elements (LREE, La–Nd) than middle (MREE, Sm–Gd) and heavy rare earths (HREE, Tb–Lu), clearly indicative of fractionation during uptake (see Discussion). The average whole-bone concentration of uranium (U), 51 ppm, is distinctly higher than those of bones from the Hell Creek Formation known to also yield endogenous collagen I (2–38 ppm [63,64]).

4.2. Intra-Bone Concentration Depth Profiles

Each REE exhibits a steeply-declining concentration profile from the cortical margin. As an example, lanthanum (La) concentrations decrease from ~1,000 ppm at the outer edge of the cortex to ~10 ppm at 5 mm, thus constituting an order of magnitude decrease across this distance (Figure 2A). Concentrations of HREE and the latter half of the MREE series, as well as U, Sc, Y, and lutetium (Lu), all increase toward the internal end of the transect (Figure 2A–C). For example, Yb concentrations increase from ~1–2 ppm at a depth of 25 mm to ~15–20 ppm at the internal end of the transect (Figure 2A, Data S1). Such increases signify secondary diffusion from within the medullary cavity (see Discussion and Supplementary Materials). Among REE, Ce exhibits the highest concentration at the cortical margin (~3,000 ppm) whereas Lu exhibits the lowest (~10 ppm). LREE generally exhibit the steepest concentration profiles, reflective of spatially-heterogeneous uptake, whereas HREE generally exhibit flatter profiles reflective of comparatively more spatially-homogenous uptake. MREE profiles are generally intermediate in steepness, and MREE concentrations commonly fall below lower detection limit in the middle and internal cortices (Data S1).

Table 1. Average whole-bone trace element composition of the left tibia of *Brachylophosaurus canadensis* MOR 2598. Numbers presented are averages of all transect data acquired across the cortex. Iron (Fe) is presented in weight percent (wt. %), all other elements are in parts per million (ppm). Absence of $(Ce/Ce^*)_N$, $(Pr/Pr^*)_N$, $(Ce/Ce^{**})_N$, and $(La/La^*)_N$ anomalies occurs at 1.0. The Y/Ho value reflects this mass ratio.

Element	Concentration
Sc	59.23
Mn	834
Fe	0.94
Sr	2499
Y	190
Ba	1448
La	40.65
Ce	88.61
Pr	9.86
Nd	35.26
Sm	7.41
Eu	2.61
Gd	12.41
Tb	2.09
Dy	17.33
Ho	4.44
Er	15.26
Tm	2.21
Yb	15.42
Lu	2.62
Th	0.18
U	51.13
ΣREE	256
$(Ce/Ce^*)_N$	1.04
$(Pr/Pr^*)_N$	0.95
$(Ce/Ce^{**})_N$	1.10
$(La/La^*)_N$	1.12
Y/Ho	42.85

Brief spikes in concentrations typically encountered in osteonal tissue around Haversian canals are rare and generally of miniscule magnitude. Although this would seem to imply a lack of major uptake through vascular systems, most REE (e.g., La in Figure 2A) profiles exhibit a subtle deflection near 2.5 mm reflective of uptake via double medium diffusion (*sensu* [76]). Near 1 mm, numerous elements, especially Y, HREE, Sc, and U, alternatively exhibit a roughly 80% increase in concentrations over values at the cortical margin (e.g., Figure 2A–C), perhaps reflective of late diagenetic near-surface leaching (see Discussion).

Fe, Ba, Sr, and Mn each exhibit very flat profiles at higher concentrations than all other elements (Figure 2C,D), with Fe exhibiting both the highest values and greatest range in variation of concentrations across the transect among these four elements. Unlike all other elements we investigated, Sc and U each exhibit “W-shaped” profiles with increasing cortical depth: after slowly decreasing from the cortical margin, each profile includes a broad, moderate peak in concentrations in the central portion of the middle cortex followed by steadily increasing concentrations across the internal cortex (Figure 2B). Y exhibits the same profile shape as HREE in the tibia of MOR 2598 (Figure 2C), indicating similar uptake behavior for these elements in this fossil.




Figure 2. Intra-bone concentration gradients of various elements in the tibia of MOR 2598. (A) Lanthanum (La) and ytterbium (Yb). (B) Scandium (Sc) and uranium (U). (C) Iron (Fe) and yttrium (Y). (D) Barium (Ba), manganese (Mn), and strontium (Sr). Note the different concentration scales for each panel. The laser track is denoted by the yellow line in each bone cross section. Gray text labels in (A) span the approximate regions considered as the ‘external’, ‘middle’, and ‘internal’ cortices. Scale bars, in white over bone images, each equal 5 mm.

4.3. NASC-Normalized REE Patterns

Spider diagrams of NASC-normalized REE concentrations reveal overall HREE enrichment in the bone as a whole (Figure 3B,C), but significant relative enrichment of LREE within the external 250 μm of the cortex (Figure 4A). A ternary plot of Nd_N - Gd_N - Yb_N confirms this trend of relative HREE enrichment by revealing that a data point for the specimen as a whole plots more closely to the Yb corner (Figure 3C). Whereas there is no apparent Ce anomaly in the bone as a whole (Figure 3B), the external-most 250 μm of the cortex exhibits a modest positive Ce anomaly (seen as an upward deflection of the pattern at this element; Figure 4A). REE concentrations range from ~25–50 times NASC values in the external 250 μm of the cortex.

Substantial spatial heterogeneity in REE composition is evident in both a ternary plot of La_N - Gd_N - Yb_N (Figure 3D) and a spider diagram of individual laser runs compiled into the full transect (Figure 4B). Both of these figures reveal significantly greater LREE content in the external-most laser run compared to all other laser runs (i.e., variation exceeds two standard deviations), signifying variations in composition are largely controlled by cortical depth. In general, the bone becomes increasingly enriched in MREE and HREE relative to LREE with increasing cortical depth, with transects through the middle and internal cortices exhibiting both similar magnitudes of REE enrichment and drastic relative enrichment in HREE (Figure 4B). Proportionally, transects through these internal regions of the bone exhibit one to two orders of magnitude enrichment in HREE over LREE, compared to just half an order of magnitude HREE enrichment in the external-most transect.

All laser runs through the middle and internal cortices exhibit isolated peaks at gadolinium (Gd; Figure 4B), likely attributable to isobaric interference effects from LREE oxides [77,78]. Most spider diagrams, especially those which separately plot data from the external cortex (e.g., Figure 4), also exhibit subtle peaks at europium (Eu) and

holmium (Ho). These peaks impart a weak 'M' shape to the shale-normalized patterns, which most authors (e.g., [79] and references therein) attribute to influences of tetrad effects during uptake (also see Supplementary Materials for further discussion on potential tetrad effects in MOR 2598).




Figure 3. REE composition of the tibia of MOR 2598. (A) Three-point moving average profile of La concentrations in the outermost 5 mm of the bone. (B) Average NASC-normalized REE composition of the fossil specimen as a whole. (C,D) Ternary diagrams of NASC-normalized REE. (C) Average composition of the bone. (D) REE compositions divided into data from each individual laser transect (~5 mm of data each). Compositional data from the transect that included the outer bone edge is denoted by a dark diamond; all other internal transect data are indicated by gray circles. The 2σ circle represents two standard deviations based on ± 5% relative standard deviation.




Figure 4. Spider diagrams of intra-bone NASC-normalized REE distribution patterns within the tibia of MOR 2598. (A) Average composition of the outermost 250 μm of the cortex, demonstrating substantially-greater LREE and MREE content in the outermost cortex compared to in the bone as a whole (Figure 4B). (B) Variation in compositional patterns by laser transects. The pattern which includes the external margin of the bone is shown in black, those from deepest within the bone by dotted, light-gray lines, and all other analyses in between by solid, dark-gray lines.

4.4. $(La/Yb)_N$ vs. $(La/Sm)_N$ Ratio Patterns

The tibia of MOR 2598 exhibits a whole-bone average $(La/Sm)_N$ value of 0.99 and a $(La/Yb)_N$ of 0.26. These values signify modest HREE enrichment relative to many environmental water samples, dissolved loads, and sedimentary particulates. Specifically, these values place the bone within the compositional range of river waters, brackish estuary waters, and marine pore fluids (Figure 5A).




Figure 5. $(La/Yb)_N$ and $(La/Sm)_N$ ratios of the tibia of MOR 2598. **(A)** Comparison of the whole-bone average $(La/Yb)_N$ and $(La/Sm)_N$ ratios of the fossil to ratios from various environmental waters and sedimentary particulates. Literature sources for environmental samples are provided in the Supplementary Materials. **(B)** REE compositions of individual laser transects expressed as NASC-normalized $(La/Yb)_N$ and $(La/Sm)_N$ ratios. The transect including the external bone margin is denoted by the black symbol whereas all other (internal) transects are represented by gray symbols.

When plotted by individual laser run (Figure 5B), REE ratios from the middle and internal cortices plot with similar $(La/Sm)_N$ values to those seen in the external cortex but with consistently lower $(La/Yb)_N$ values. This difference encompasses roughly one order of magnitude, on average. The most internal laser run exhibits the lowest $(La/Yb)_N$ ratio (0.006), and all but two laser runs across the middle and internal cortices exhibit $(La/Yb)_N$ ratios < 0.1. $(La/Sm)_N$ ratios range between 0.8–1.5 and exhibit no apparent relationship with cortical depth.

4.5. REE Anomalies

Due to concentrations of many trace elements in the middle cortex being so low that they fall below lower detection limit (Data S1), every anomaly examined exhibits major gaps in coverage through this region of the bone (Figure S1). Occasional instances of significantly higher neodymium (Nd) than praseodymium (Pr) concentrations also create gaps in the anomaly profiles. Whereas $(Ce/Ce^*)_N$ and La-corrected $(Ce/Ce^{**})_N$ anomalies are absent at the outer cortex edge, $(La/La^*)_N$ anomalies are slightly negative in the external-most ~280 μ m (Figure S1). All three of these anomalies exhibit substantial positive and negative fluctuations across the transect.

Although $(Ce/Ce^*)_N$ anomalies fluctuate from ~0.2–20 across the transect, they are largely positive throughout most of the internal half of the transect (Figure S1). This trend is not reflected, however, in the whole-bone $(Ce/Ce^*)_N$ average for the tibia, which is essentially absent (1.04; Table 1). $(Ce/Ce^*)_N$ values were also plotted against $(Pr/Pr^*)_N$ values (following [80]) to aid us in differentiating true, redox-related cerium anomalies from apparent anomalies induced by variations in $(La/La^*)_N$ anomalies. The majority of $(Ce/Ce^*)_N$ values from the external 1 mm of the bone plot near the lower margins of fields 3a and 4a (Figure 6), reflective of slightly negative La anomalies in the external cortex (in agreement with Figure S1). In contrast, anomaly values from inner regions of the cortex plot over a broad range encompassing every field of the diagram (Figure 6), indicative of

substantial heterogeneity at the sub-millimeter scale in the middle and internal cortices. Within this broad spectrum, there are relatively few data points in fields 1 and 2b (Figure 6); regions of the internal cortex represented by these data points lack a Ce anomaly.




Figure 6. $(\text{Ce}/\text{Ce}^*)_{\text{N}}$ vs. $(\text{Pr}/\text{Pr}^*)_{\text{N}}$ plot (after [80]) of five-point averages along the transect across the cortex of MOR 2598 recorded by LA-ICPMS. Separate fields (labeled by blue text) are as follows: 1, neither Ce nor La anomaly; 2a, no Ce and positive La anomaly; 2b, no Ce and negative La anomaly; 3a, positive Ce and negative La anomaly; 3b, negative Ce and positive La anomaly; 4a, negative Ce and negative La anomaly; 4b, positive Ce and positive La anomaly. Measurements from the outer 1 mm of the external cortex are plotted as black triangles, and all measurements from deeper within the bone are plotted as grey diamonds. $(\text{Ce}/\text{Ce}^*)_{\text{N}}$ vs. $(\text{Pr}/\text{Pr}^*)_{\text{N}}$ anomalies calculated as in the Materials and Methods section of the text.

$(\text{La}/\text{La}^*)_{\text{N}}$ anomalies and La-corrected $(\text{Ce}/\text{Ce}^{**})_{\text{N}}$ anomalies were also directly calculated (see Methods) to quantitatively assess these qualitative inferences. Unfortunately, as mentioned above, frequent drops in concentrations of LREE below detection limit severely limit coverage in these profiles. However, based on Figure S1, it is clear that $(\text{La}/\text{La}^*)_{\text{N}}$ anomalies are exclusively negative in the internal 29 mm of the cortex. The average $(\text{La}/\text{La}^*)_{\text{N}}$ value across this region is 0.20. $(\text{Ce}/\text{Ce}^{**})_{\text{N}}$ anomalies are also almost exclusively negative in the middle and internal cortices, exhibiting a similarly-low average (0.65) across this same span. Values of both of these anomalies fluctuate by roughly two orders of magnitude across the transect. As a whole, the tibia of MOR 2598 exhibits slightly positive $(\text{Ce}/\text{Ce}^{**})_{\text{N}}$ and $(\text{La}/\text{La}^*)_{\text{N}}$ anomalies (1.10 and 1.12, respectively; Table 1), but these are each clearly biased by overweighting of data from the external cortex (caused by abundant missing data from internal regions of the bone, as discussed above). Plotting $(\text{Ce}/\text{Ce}^{**})_{\text{N}}$ anomalies against U concentrations for each laser run yielded a poor correlation between these two redox-sensitive signatures ($r^2 = 0.29$; Figure S2).

Yttrium/holmium (Y/Ho) ratios are slightly above chondritic (26; [81]) in the outer ~10 mm of the bone and the innermost ~7 mm of the transect, wherein they range ~20–300. Though data are sporadic through the middle cortex due to very low concentrations, ratios from this region form a broad swale below these peaks in the external and internal cortices (Figure S1). Specifically, the average of Y/Ho ratios through the central 15 mm of the cortex (10–25 mm along the transect) is 35, and values across this region mostly fall between ~10–80. These spatial contrasts partially negate one another when data are averaged for the entire transect, which yields a slightly positive whole-bone average anomaly of 43 (Table 1).

5. Discussion

5.1. MOR 2598's Paleoenvironmental and Taphonomic Context

Though limited taphonomic and stratigraphic data are available, the excellent preservation quality of MOR 2598 suggests it was protected by rapid burial postmortem. Full articulation of the hind limb, negligible signs of (ancient) weathering and abrasion, and excellent histological preservation [70] each support the interpretation that burial took place within a few years postmortem, perhaps even just weeks after death [82]. However, absence of the remainder of the skeleton implies this hind limb became disarticulated from the remainder of the carcass during brief subaerial decay, as well as probable short-distance transport of the limb (from an upstream site of death) prior to burial (cf., [83,84]). Recovery of MOR 2598 from a channel sandstone [30] strongly suggests that: 1) decay primarily occurred subaqueously under oxygenated conditions (cf., [85]); 2) fluvial currents likely caused the separation of the limb from the body, and; 3) the carcass likely reached the “bloat” or active decay phase of postmortem decomposition for this to occur (cf., [86,87]). Ultimately, a lull in flow competency induced deposition and burial of the limb within the channel. Based on the great thickness of the succession of channel sandstone horizons from which MOR 2598 was recovered (7 m), it appears burial occurred within a well-established lowland channel rather than a recently-formed avulsion channel. This conclusion is consistent with prior interpretations of the Judith River Formation as generally representing lowland fluvial environments close to the coastline of the Western Interior Cretaceous Seaway (WIKS; [88,89]).

To briefly summarize, the available data reveal that this *Brachylophosaurus canadensis* individual (MOR 2598) died within or near a fluvial channel on the coastal lowlands. Its carcass experienced fairly brief decay within the channel, where currents eventually led to disarticulation of the left hind limb which was carried shortly downstream. Either an obstruction in the channel, a temporal lull in flow competency, or slowing of currents due to gradual channel broadening caused deposition of the limb on the channel floor where it became quickly buried and fossilized within cross-stratified sands. Our trace element data provide illuminating insights into the ensuing diagenetic history of MOR 2598, which we now characterize in an effort to constrain geochemical pathways to cellular, soft tissue, and biomolecular preservation.

5.2. Reconstructing the Geochemical History of MOR 2598

The tibia of MOR 2598 exhibits low REE concentrations near the cortical margin (e.g., ~800 ppm for La; Data S1) and a low whole-bone Σ REE (256 ppm) compared to many other bones from the Cretaceous period (Table 2), which have been found to possess Σ REE ranging from 1,110–25,000 ppm [58,63,90–93]. Notably, however, these values each fall within the range of other dinosaur bones we have examined from the Cretaceous Hell Creek Formation [63,64] which have also been found to yield endogenous proteins [28,29,47]. Compared to those specimens from the Hell Creek Formation, MOR 2598 exhibits lower average concentrations of Fe (0.94 wt.%), Mn (834 ppm), and Y (190 ppm), a higher concentration of U (51 ppm), and similar concentrations of Sr (2,499 ppm), Ba (1,448 ppm) and Lu (3 ppm). Although these comparisons do not take into account differences in taxon, cortical width, histology, or diagenetic regimes, they still reveal that the tibia of MOR 2598 is (for most elements examined) less chemically altered than the majority of fossil bones of similar age. We have previously attributed such cases of minimal alteration to various sequestration processes limiting the availability of trace element ions in early-diagenetic pore fluids (e.g., complexation with humic acids and/or dissolved carbonates [94–97], coprecipitation with phosphates in entombing sediments [98–101]), and those processes may also account for the modest alteration of MOR 2598 (see Supplementary Materials for further discussion).

Table 2. Summary of the REE composition of the left tibia of *Brachylophosaurus canadensis* MOR 2598. Qualitative Σ REE content is based on the value shown in Table 1 (256 ppm) in comparison to values from other Mesozoic bones (as listed in the main text). Abbreviations: DMD, double medium diffusion *sensu* [76]; LREE, light rare earth elements.

Clear DMD Kink for LREE?	Relative Noise in Outer Cortex for La	REE Suggest Flow in Marrow Cavity?	Relative Σ REE Content (Whole Bone)	Relative U Content (Whole Bone)	Relative Porosity of the Cortex
Yes	Moderate	Yes	Low	Moderate	Low

Ba, Fe, and Mn each exhibit flat concentration profiles (Figure 2C,D) probably indicative of incorporation into homogeneously-distributed, minute, secondary mineral phases, presumably barite, goethite, and Mn oxides [102]. Sr likely exhibits a similarly-flat profile shape due to spatially-homogenous substitution for Ca in bone hydroxyapatite [103,104]. In contrast to these more abundant elements, all REE exhibit steep declines in concentrations from the cortical margin, with LREE exhibiting the steepest declines and HREE the shallowest (due to crystal-chemical controls based on ionic radius [105]). Meanwhile, MREE concentrations commonly drop below detection limit in the middle and internal cortices (Data S1). These trends are typical of fossil bones which have experienced relatively brief uptake largely by simple ‘external-to-internal’ diffusion and did not equilibrate with external pore fluids during diagenesis (cf., [60,63,64,68,106]). However, clear kinks in concentration profiles for many REE and substantial, locally-restricted variations in their concentrations in the external cortex (e.g., La in Figure 2A) also indicate at least partial uptake via double medium diffusion (*sensu* [76]) through Haversian canals.

Spider diagrams reveal even proportions of REE within the external-most cortex (Figure 4A) yet significant relative HREE enrichment throughout the bone as a whole (Figures 3B,C and 4B). Relative HREE enrichment is especially evident in the internal cortex, where, for example, concentrations of Yb rise to more than double those of La (Figure 2A). These signatures are very similar to, but less pronounced than, those observed in a *Tyrannosaurus rex* femur recovered from an estuarine channel sandstone in the Hell Creek Formation [64]. As with that specimen, it is likely that such HREE enrichment reflects protracted trace element uptake from relatively HREE-enriched brackish waters and/or diagenetic pore fluids under oxidizing conditions. This interpretation is supported by the whole bone composition of this specimen being similar to those of lowland river waters, estuarine waters, and marine pore fluids (Figure 5A) which typically exhibit such relative HREE enrichment [55,90,107]. Interestingly, these findings strongly suggest that the channel in which MOR 2598 was interred was likely tidally-influenced, which in turn suggests that it was recovered from an estuarine channel, not a (strictly-speaking) fluvial channel – an insight not apparent from the sedimentology/stratigraphy of the quarry.

Regarding redox regimes through diagenesis, at the whole-bone level the tibia exhibits a slightly-positive $(\text{Ce}/\text{Ce}^{**})_{\text{N}}$ anomaly (1.10) reflective of a weakly-oxidizing overall diagenetic history; this is consistent with the inferred burial setting having been an estuarine channel (see above). Generally high U concentrations throughout much of the cortex (Figure 2B and Table 1) and a relatively high average Sc concentration (59 ppm) corroborate this signal, as U and Sc enrichment have each been linked with uptake under oxidizing conditions [108–110]. Plotting of numerous data points from the middle and internal cortices in fields 3b and 4a of the $(\text{Ce}/\text{Ce}^{*})_{\text{N}}$ vs. $(\text{Pr}/\text{Pr}^{*})_{\text{N}}$ plot (Figure 6) also support the presence of oxidizing conditions within the bone. However, both $(\text{Ce}/\text{Ce}^{*})_{\text{N}}$ and $(\text{Ce}/\text{Ce}^{**})_{\text{N}}$ anomalies are essentially absent at the cortical margin, whereas their values fluctuate considerably (both positively and negatively) throughout the middle and internal cortices (Figure S1). These contrasts, as well as the broad distribution of data points in Figure 6, demonstrate the presence of considerable spatial heterogeneity in

redox conditions throughout the bone through diagenesis, especially the middle and internal cortices.

Although the trace element anomaly profiles in Figure S1 may seem somewhat stochastic, $(\text{Ce}/\text{Ce}^*)_{\text{N}}$ and $(\text{Ce}/\text{Ce}^{**})_{\text{N}}$ values are generally positive and negative, respectively, across the inner half of the transect. If these trends are taken as reliable records of redox regimes during early diagenesis/fossilization, as all indications appear to support (see below), then these anomaly trends would signify that uptake occurred under prevailingly reducing conditions within the interior of the bone. High Sc concentrations in the internal cortex (Figure 2B) appear incompatible with this interpretation (as this should be a product of uptake under oxidizing conditions), but we attribute this apparent “contrast” to temporal changes in redox conditions in this region of the bone through early diagenesis. Specifically, these conflicting signals could arise via significant uptake of Sc in the internal cortex under initially-oxidizing conditions followed by a shift to reducing conditions, during which latter time the internal cortex secondarily acquired positive $(\text{Ce}/\text{Ce}^*)_{\text{N}}$ anomalies (and Sc ions remained sequestered by, as in adsorbed to, bone crystallites; also see the Supplementary Materials for further discussion of the peculiar shapes of Sc and U concentration profiles in MOR 2598).

The redox scenario just described would necessitate supply of significant amounts of Sc to the interior of the bone. This would have to be supplied by a pore fluid percolating through the medullary cavity, which would presumably also supply numerous other trace elements to the internal cortex. Concentration profiles of HREE (e.g., Yb in Figure 2A), U (Figure 2B), Y, Lu, and the latter half of the MREE series (Data S1) each exhibit increases toward the internal end of the transect (in the internal cortex), providing concrete evidence of uptake from a second diffusion front in the interior of the bone. That LREE exhibit negligible rises in concentrations toward the internal end of the transect (e.g., La in Figure 2A) indicates that the majority of elements supplied by the pore fluid passing through the medullary cavity were mostly those with comparatively-modest to low diffusivities (based on [106]). This bias signifies that the pore fluid must have been a chemically-‘evolved’, highly-fractionated fluid which, based on the magnitude of select elemental enrichments in the internal cortex (e.g., Figure 2B), either flowed through the medullary cavity for an extended period of time or, more likely due to burial and compaction, became pooled there, allowing protracted uptake. It also apparent that this pore fluid was likely not simply an HREE-enriched solution passing through during some later phase of late diagenesis because there are no clear signs of similar HREE enrichment in the external-most cortex (e.g., Figure 2A,B). Instead, the majority of elements exhibiting enrichment toward the internal end of the transect (e.g., U, Y, HREE) exhibit a subtle ‘plateau’ of stable concentrations in the outermost ~1 mm of the cortex followed by a ~80% increase near ~1.5–2 mm (Figure 2A–C). We interpret this pattern to reflect modest leaching of trace elements from the outermost ~1 mm of the external cortex, most likely during late diagenesis and under slightly oxidizing conditions (based on the weakly positive $(\text{Ce}/\text{Ce}^*)_{\text{N}}$ anomalies in this region; Figure S1). This conclusion may also be supported by: 1) the common presence of negative $(\text{La}/\text{La}^*)_{\text{N}}$ anomalies in the outermost ~500 μm of the external cortex (potentially reflective of near-surface loss of La; Figure S1), and; 2) lack of a correlation between U concentrations and $(\text{Ce}/\text{Ce}^{**})_{\text{N}}$ anomalies for each laser run ($r^2 = 0.29$; Figure S2), implying uptake of U and REE over differing timescales [110].

Absence of more major signs of leaching or late-diagenetic trace element uptake at the cortical edge (Figure 2), as well as retention of clear evidence of spatiotemporal changes in pore fluid compositions (described in the last few paragraphs), imply that late-diagenetic overprinting of trace element signatures in MOR 2598 was not substantial, and, therefore, that the tibia at least partially retains early-diagenetic signatures. Indeed, there are numerous signs of relatively brief interaction with pore fluids. For example, a spider diagram (Figure 4B) and ternary plot (Figure 3D) of REE proportions by individual laser runs each reveal clear signs of significant fractionation during uptake

from circum-neutral pH surface/groundwaters (cf., [111]) in the form of increasing relative LREE depletion/HREE enrichment with increasing cortical depth (as in, e.g., [59,105,112]). These fractionation effects are also evident from significant spatial variations in $(\text{La}/\text{Yb})_N$ ratios (Figure 5B), a positive $(\text{La}/\text{La}^*)_N$ anomaly and Y/Ho ratio for the bone as a whole (1.12 and 43, respectively; cf., [60]), and differing concentration profile shapes (contrast Figure 2A,B) for U and REE (which should otherwise have similar shapes due to their similar diffusivities [106]). Similarly-high Y/Ho ratios in the external and internal cortices (Figure S1) further imply fractionation also occurred during uptake from the chemically ‘evolved’ pore fluid pooled in the medullary cavity.

To review, our trace element data thus identify that after brief subaqueous decay and transport down a fluvial stream, the hind limb of MOR 2598 became fossilized after burial in a sandy, oxic, estuarine channel (Figure 7). Its bones experienced relatively brief primary uptake of REE and other trace elements from circum-neutral pH, HREE-enriched, and potentially-brackish channel waters and groundwaters under oxidizing conditions during early diagenesis. Comparatively-slower percolation of pore fluids through the medullary cavity of the tibia than around its exterior led to the development of reducing conditions inside the bone. Recent erosion, typical of that in desert/badlands environments from which most fossil bones are recovered, re-exposed MOR 2598 to oxidizing conditions and caused minor leaching of trace elements from the outermost ~1 mm of the cortex, but had no significant effects on the overall chemistry of the bone. As a result, the specimen remains modestly altered compared to others of similar age.

5.3. Insights into Molecular Taphonomy from Comparative Geochemistry

MOR 2598 is only the third vertebrate fossil of pre-Cenozoic age to have both yielded endogenous protein and have its geochemical history characterized through trace element analyses. Both other specimens in this short list, namely *Tyrannosaurus rex* MOR 1125 [28,29,64] and *Edmontosaurus annectens* SRHS-DU-231 [47,63], are also large nonavian dinosaurs recovered from Late Cretaceous strata in Montana, USA. Adding MOR 2598 into this comparative framework reveals both geochemical similarities to, and differences from, these two other specimens which: 1) bolster prior hypotheses about protein preservation pathways, and; 2) add new insights into the complexity of post-burial diagenetic alterations which biomolecules can withstand.

Taken as a whole, the biostratinomic history of *Brachylophosaurus* MOR 2598 is quite similar to that which we recently explicated for *T. rex* MOR 1125 [64]. After death and a short period of (likely subaqueous) decay, brief fluvial transport brought each specimen into coastal estuaries along the western coast of the WIKS where they were rapidly buried in sandy estuarine channels, and throughout this history, the bones of each specimen were acquiring trace elements from brackish, HREE-enriched surface and groundwaters (this study and [64]). As at the MOR 1125 quarry [64], early cementation of the sediments entombing MOR 2598 appears to have limited trace element uptake by the fossil bones, allowing them to exhibit minimal alteration at the elemental level. This is evident in the low ΣREE of the tibia compared to many other bones of Cretaceous age (as discussed above), as well as its steep declines in REE concentrations from the cortical margin (Figure 2A) and very low concentrations of elements with ionic radii similar to that of Ca^{2+} (i.e., MREE) in the middle cortex (Data S1). Thus, MOR 2598 adds further support to the assertions of Schweitzer [113], Herwartz et al. [59], and Ullmann et al. [64] that: 1) early-diagenetic cementation of sediments can effectively thwart protracted decay and chemical alteration of bones after burial (presumably by minimizing exposure to percolating groundwaters and the exogenous microbes they carry with them), and; 2) this diagenetic pathway also facilitates rapid molecular stabilization. Fossils from the Standing Rock Hadrosaur Site (SRHS; [47,63,114]) demonstrate that rapid burial in fine-grained sediments with low-permeability and/or encasement in early-diagenetic

concretion can similarly hinder decay of endogenous cells, tissues, and their component biomolecules.




Figure 7. Reconstruction of the diagenetic history of *Brachylophosaurus canadensis* MOR 2598, summarizing macroscale and microscale event effecting its skeletal elements after death. Events are classified into approximate taphonomic stages, with a visual portrayal of the diagenetic processes affecting the bones during each stage shown at left.

As for SRHS-DU-231 [63] and MOR 1125 [64], MOR 2598 exhibits high Sc enrichment and a slightly positive whole-bone $(\text{Ce}/\text{Ce}^{**})_{\text{N}}$ anomaly (Table 1) reflective of a generally-oxidizing diagenetic history. Although this pattern superficially supports the proposition by Wiemann et al. [54] that oxidizing depositional environments may be more favorable settings for molecular preservation than reducing environments, it is clear that both MOR 2598 and bones from SRHS also experienced reducing conditions during diagenesis. This is evident from consistently-positive $(\text{Ce}/\text{Ce}^{*})_{\text{N}}$ anomalies in the internal cortex of the tibia of MOR 2598 (Figure S1) and external cortices of many SRHS bones (figure 6 of [63]). Microbial decay of organics (including those within bones) is known to decrease local pH and create dysoxic to anoxic conditions, thereby eliciting the

production of reducing conditions [115], especially within confined microenvironments, such as those within and around a carcass in compacted or low-permeability sediments (e.g., [50,116]) or within the internal pore spaces of bones (e.g., [102,117,118]). It is also well known that reducing early-diagenetic conditions do not preclude exquisite morphologic and biochemical preservation of structural soft tissues, perhaps due to the induced dysoxia/anoxia (e.g., [119–122]). Thus, there are reasons that fossil bones preserved under largely-reducing conditions may still yield original molecules.

Though the acidic pH that would have temporarily accompanied reducing conditions within the medullary cavity of this specimen may seem (at face value) preclusive to molecular preservation, weak acidity has been implicated in the rapid nucleation of inert, protective, microcrystalline goethite crystals within ‘osteocytes’ and ‘blood vessels’ recovered from fossil bones [4,53]. For this reason, initial redox conditions in the immediate ~48 hrs after death may ultimately be the most critical, as it has been demonstrated that inter- and intramolecular crosslinking (i.e., stabilization) reactions can operate in this brief timeframe, promoting equilibration with the early-diagenetic environment which may then persist through fossilization and late diagenesis [53]. However, we must note that actualistic studies examining molecular stability through temporal shifts in redox regimes would be necessary to further evaluate this hypothesis. Regardless, our recognition of varied pH and redox conditions over time within the tibia of MOR 2598 is thus not incompatible with these prior studies, but rather augments them by revealing that biomolecular remains may survive multiple changes in redox regimes through diagenesis (Figure 7).

In particular, as discussed above, bones of MOR 2598 experienced two comparatively ‘extra’ diagenetic events which MOR 1125 and bones from SRHS did not [63,64]: protracted trace element uptake from stagnant pore fluids ‘pooled’ in the central medullary cavity (evidenced by elevated concentrations of HREE, U, Sc, Y, and Lu in the internal cortex; Figure 2, Data S1), and; 2) modest late-diagenetic leaching of the cortical surface (evidenced by negative $(La/La^*)_N$ anomalies and reduced concentrations of many trace elements in the outermost ~0.5–1 mm; Figures 2 and S1). Despite this complex diagenetic history, cortical bone from the femur of MOR 2598 has yielded numerous microstructures retaining endogenous peptide sequences of multiple proteins (namely collagen, actin, tubulin, histones, myosin, and tropomyosin [18,30,45]). This fact indicates that processes which stabilized diagenetiforms within this specimen in the initial hours to days postmortem imparted remarkable long-term resiliency. Novel experiments by Schwietzer et al. [4] and Boatman et al. [53] have begun to shed light on how this may occur, but testing of more fossils and further actualistic studies are needed to fully resolve the endurance of diagenetiforms under the wide array of physicochemical/thermodynamic regimes of natural diagenetic environments.

Finally, all three specimens examined in this discussion (MOR 2598, SRHS-DU-231, MOR 1125) also each exhibit abundant signals of fractionation of REE during uptake, which when combined with their low Σ REE content, indicate at least partial retention of early-diagenetic trace element signatures. These signs include positive whole-bone Y/Ho anomalies (Table 1, and table 1 in [63] and [64]), negative $(La/La^*)_N$ anomalies in the middle and internal cortices (Figure S1, and figure S2 of [64]), steeper concentration profiles for LREE than HREE (Figure 2A, figure 2 of [63], and figure 4A of [64]), and increasing relative-HREE enrichment with cortical depth (Figure 4B, figure 4 of [63], and figure 6B of [64]). As suggested by Ullmann et al. [64], retention of such early-diagenetic signatures may constitute a useful proxy for molecular recovery potential because it indicates a specimen has avoided protracted interactions with any late-diagenetic pore fluids (e.g., phreatic groundwaters) which could plausibly cause hydrolysis and other decay processes.

6. Conclusions

Synthesizing our results with those of other recent experimental and actualistic studies in molecular taphonomy leads us to conclude the following:

1. By allowing quick characterization of spatial patterns of diagenetic alteration within a fossil, trace element analyses constitute a useful and effective means of screening fossil tissues prior to paleomolecular analyses;
2. Retention of early-diagenetic trace element signatures may constitute a useful proxy for molecular recovery potential because it indicates a specimen has avoided protracted interactions with any late-diagenetic pore fluids.
3. Although burial in coarse, permeable sediments in oxic environments appears conducive to molecular preservation, it is also possible for fossils preserved in reducing paleoenvironments to yield endogenous molecules. This dichotomy suggests that presence of oxidizing conditions in the initial ~48 hours postmortem may be more key to molecular preservation than the redox state of the final setting of burial;
4. Rapid burial in fine-grained sediments with low permeability, encasement in early-diagenetic concretion, and/or early-diagenetic cementation of entombing sediments can effectively thwart protracted decay and chemical alteration of bones (and their component cells, tissues, and biomolecules) by minimizing exposure to percolating groundwaters and the exogenous microbes they carry with them, and;
5. Biomolecular remains may survive multiple changes in redox regimes through diagenesis, and this indicates that processes which stabilize biomolecules in the initial hours to days postmortem can impart remarkable long-term resiliency.

While these insights are obviously enlightening, it must be reiterated that they primarily derive from fossils from just three Cretaceous localities. In agreement with Schroeter et al. [123], we propose that paleomolecular and trace element analyses on Paleogene and Miocene fossils are direly needed to close the long window of the Cenozoic for which protein preservation has yet to be explored. Based on all data currently available, it seems very likely that future studies will continue to broaden the suite of depositional and diagenetic circumstances known to be conducive to molecular preservation.

Supplementary Materials: The following supplementary data to this article can be found online at [_](#): Supplementary Materials narrative: A DOCX Word file including further details on our LA-ICPMS methods, discussion of potential tetrad effects in the left tibia of MOR 2598, discussion of potential sequestration processes that may have limited REE availability to the bones of MOR 2598, discussion of potential causes for the peculiar concentration profile shapes of U and Sc in the left tibia of MOR 2598, sources for environmental data in Figure 5, and additional data on the trace element composition of the left tibia of MOR 2598 (Figures S1, S2). Data S1: Raw transect data acquired and analyzed in this study, provided as an XLSX Excel spreadsheet.

Author Contributions: Conceptualization, P.V.U.; methodology, P.V.U.; formal analysis, P.V.U. and R.D.A.; investigation, P.V.U.; resources, J.B.S.; data curation, P.V.U. and J.B.S.; writing—original draft preparation, P.V.U.; writing—review and editing, P.V.U. and J.B.S.; visualization, P.V.U.; project administration, P.V.U.; funding acquisition, P.V.U. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Rowan University.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: All data generated by this study are available in this manuscript and the accompanying Supplementary Materials.

Acknowledgments: We thank Bob Harmon, Nels Peterson, and the rest of their Museum of the Rockies field crews for their efforts to excavate and collect this specimen over the course of two

field seasons in 2006 to 2007. Gratitude is also extended to the Montana Department of Natural Resources and Conservation for access to the land where MOR 2598 was recovered. We also thank Ellen-Thérèse Lamm for specimen sampling and Elena Schroeter, Jack Horner, and Mary Schweitzer for helpful discussions.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Curry, G.B. Molecular palaeontology: new life for old molecules. *Trends Ecol. Evol.* **1987**, *2*, 161–165.
- Schweitzer, M.H. Soft tissue preservation in terrestrial Mesozoic vertebrates. *Annu. Rev. Earth Pl. Sc.* **2011**, *39*, 187–216.
- Hofreiter, M.; Collins, M.; Stewart, J.R. Ancient biomolecules in Quaternary palaeoecology. *Quaternary Sci. Rev.* **2012**, *33*, 1–13.
- Schweitzer, M.H.; Zheng, W.; Cleland, T.P.; Goodwin, M.B.; Boatman, E.; Theil, E.; Marcus, M.A.; Fakra, S.C. A role for iron and oxygen chemistry in preserving soft tissues, cells and molecules from deep time. *Proc. R. Soc. B* **2014**, *281*, 1–10.
- Schweitzer, M.H.; Schroeter, E.R.; Cleland, T.P.; Zheng, W. Paleoproteomics of Mesozoic dinosaurs and other Mesozoic fossils. *Proteomics* **2019**, 1800251.
- Thomas, B.; Taylor, S. Proteomes of the past: the pursuit of proteins in paleontology. *Expert Rev. Proteomic.* **2019**, *16*, 881–895.
- Bailleul, A.M.; Li, Z. DNA staining in fossil cells beyond the Quaternary: reassessment of the evidence and prospects for an improved understanding of DNA preservation in deep time. *Earth-Sci. Rev.* **2021**, *216*, 103600.
- Orlando, L.; Bonjean, D.; Bocherens, H.; Thenot, A.; Argant, A.; Otte, M.; Hanni, C. Ancient DNA and the population genetics of cave bears (*Ursus spelaeus*) through space and time. *Mol. Biol. Evol.* **2002**, *19*, 1920–1933.
- Gilbert, M.T.P.; Drautz, D.I.; Lesk, A.M.; Simon, Y.W.H.; Qi, J.; Ratan, A.; Hsu, C.-H.; Sher, A.; Dalén, L.; Götherström, A.; Tomsho, L.P.; Rendulic, S.; Packard, M.; Campos, P.F.; Kuznetsova, T.V.; Shidlovskiy, F.; Tikhonov, A.; Willerslev, E.; Iacumin, P.; Buigues, B.; Ericson, P.G.P.; Germonpre, M.; Kosintsev, P.; Nikolaev, V.; Nowak-Kemp, M.; Knight, J.R.; Irzyk, G.P.; Perbost, C.S.; Fredrikson, K.M.; Harkins, T.T.; Sheridan, S.; Miller, W.; Schuster, S.C. Intraspecific phylogenetic analysis of Siberian woolly mammoths using complete mitochondrial genomes. *Proc. Natl Acad. Sci. USA* **2008**, *105*, 8327–8332.
- Miller, W.; Drautz, D.I.; Ratan, A.; Pusey, B.; Qi, J.; Lesk, A.M.; Tomsho, L.P.; Packard, M.D.; Zhao, F.; Sher, A.; Tikhonov, A.; Raney, B.; Patterson, N.; Lindblad-Toh, K.; Lander, E.S.; Knight, J.R.; Irzyk, G.P.; Fredrikson, K.M.; Harkins, T.T.; Sheridan, S.; Pringle, T.; Schuster, S.C. Sequencing the nuclear genome of the extinct woolly mammoth. *Nature* **2008**, *456*, 387–390.
- Cappellini, E.; Jensen, L.J.; Szklarczyk, D.; Ginolhac, A.; da Fonseca, R.A.R.; Stafford, Jr., T.W.; Holen, S.R.; Collins, M.J.; Orlando, L.; Willerslev, E.; Gilbert, M.T.P.; Olsen, J.V. Proteomic analysis of a Pleistocene mammoth femur reveals more than one hundred ancient bone proteins. *J. Proteome Res.* **2012**, *11*, 917–926.
- van der Valk, T.; Pečnerová, P.; Díez-del-Molino, D.; Bergström, A.; Oppenheimer, J.; Hartmann, S.; Xenikoudakis, G.; Thomas, J.A.; Dehasque, M.; Sağlıcan, E.; Fidan, F.R.; Barnes, I.; Liu, S.; Somel, M.; Heintzman, P.D.; Nikolskiy, P.; Shapiro, B.; Skoglund, P.; Hofreiter, M.; Lister, A.M.; Götherström, A.; Dalén, L. Million-year-old DNA sheds light on the genomic history of mammoths. *Nature* **2021**, *591*, 265–269.
- Orlando, L.; Ginolhac, A.; Zhang, G.; Froese, D.; Albrechtsen, A.; Stiller, M.; Schubert, M.; Cappellini, E.; Petersen, B.; Moltke, I.; Johnson, P.L.F.; Fumagalli, M.; Vilstrup, J.T.; Raghavan, M.; Korneliussen, T.; Malaspina, A.-S.; Vogt, J.; Szklarczyk, D.; Kelstrup, C.D.; Vinther, J.; Dolocan, A.; Stenderup, J.; Velazquez, A.M.V.; Cahill, J.; Rasmussen, M.; Wang, X.; Min, J.; Zazula, G.D.; Sequin-Orlando, A.; Mortensen, C.; Magnussen, K.; Thompson, J.F.; Weinstock, J.; Gregersen, K.; Røed, K.H.; Eisenmann, V.; Rubin, C.J.; Miller, D.C.; Antczak, D.F.; Bertelsen, M.F.; Brunak, S.; Al-Rasheid, A.S.; Ryder, O.; Andersson, L.; Mundy, J.; Krogh, A.; Gilbert, M.T.P.; Kjær, K.; Sicheritz-Ponten, T.; Jensen, L.J.; Olsen, J.V.; Hofreiter, M.; Nielsen, R.; Shapiro, B.; Wang, J.; Willerslev, E. Recalibrating *Equus* evolution using the genome sequence of an early Middle Pleistocene horse. *Nature* **2013**, *499*, 74–78.
- Barnett, R.; Westbury, M.V.; Sandoval-Velasco, M.; Vieira, F.G.; Jeon, S.; Zazula, G.; Martin, M.D.; Ho, S.Y.W.; Mather, N.; Gopalakrishnan, S.; Ramos-Madrugal, J.; de Manuel, M.; Zepeda-Mendoza, M.L.; Antunes, A.; Baez, A.C.; De Cahsan, B.; Larson, G.; O'Brien, S.J.; Eizirik, E.; Johnson, W.E.; Koepfli, K.-P.; Wilting, A.; Fickel, J.; Dalén, L.; Lorenzen, E.D.; Marques-Bonet, T.; Hansen, A.J.; Zhang, G.; Bhak, J.; Yamaguchi, N.; Gilbert, M.T.P. Genomic adaptations and evolutionary history of the extinct scimitar-toothed cat, *Homotherium latidens*. *Curr. Biol.* **2020**, *30*, 1–8.
- Noonan, J.P.; Coop, G.; Kudaravalli, S.; Smith, D.; Krause, J.; Alessi, J.; Chen, F.; Platt, D.; Pääbo, S.; Pritchard, J.K.; Rubin, E.M. Sequencing and analysis of Neanderthal genomic DNA. *Science* **2006**, *314*, 1113–1118.
- Welker, F.; Ramos-Madrugal, J.; Gutenbrunner, P.; Mackie, M.; Tiwary, S.; Jersie-Christensen, R.R.; Chiva, C.; Dickinson, M.R.; Kuhlwilm, M.; de Manuel, M.; Gelabert, P.; Martín-Torres, M.; Margvelashvili, A.; Arsuaga, J.L.; Carbonell, E.; Marques-Bonet, T.; Racimo, F.; Lalueza-Fox, C.; Bermúdez de Castro, J.M.; Willerslev, E.; Cappellini, E.; The dental proteome of *Homo antecessor*. *Nature* **2020**, *580*, 235–238.
- Briggs, D.E.G.; Evershed, R.P.; Lockheart, M.J. The biomolecular paleontology of continental fossils. *Paleobiology* **2000**, *26*, 169–193.
- Cleland, T.P.; Schroeter, E.R.; Zamborg, L.; Zheng, W.; Lee, J.E.; Tran, J.C.; Bern, M.; Duncan, M.B.; Lebleu, V.S.; Ahlf, D.R.; Thomas, P.M.; Kalluri, R.; Kelleher, N.L.; Schweitzer, M.H. Mass spectrometry and antibody-based characterization of blood vessels from *Brachyophosaurus canadensis*. *J. Proteome Res.* **2015**, *14*, 5252–5260.

19. Pan, Y.; Zheng, W.; Moyer, A.E.; O'Connor, J.K.; Wang, M.; Zheng, X.; Wang, X.; Schroeter, E.R.; Zhou, Z.; Schweitzer, M.H. Molecular evidence of keratin and melanosomes in feathers of the Early Cretaceous bird *Eoconfuciusornis*. *Proc. Natl Acad. Sci. USA* **2016**, *113*, 7900–7907. 767
20. Pan, Y.; Zheng, W.; Sawyer, R.H.; Pennington, M.W.; Zheng, X.; Wang, X.; Wang, M.; Hu, L.; O'Connor, J.; Zhao, T.; Li, Z.; Schroeter, E.R.; Wu, F.; Xu, X.; Zhou, Z.; Schweitzer, M.H. The molecular evolution of feathers with direct evidence from fossils. *Proc. Natl Acad. Sci. USA* **2019**, *116*, 3018–3023. 768
21. Lindgren, J.; Kuriyama, T.; Madsen, H.; Sjövall, P.; Zheng, W.; Uvdal, P.; Engdahl, A.; Moyer, A.E.; Gren, J.A.; Kamezaki, N.; Ueno, S.; Schweitzer, M.H. Biochemistry and adaptive colouration of an exceptionally preserved juvenile fossil sea turtle. *Sci. Rep.* **2017**, *7*, 13324. 769
22. Lindgren, J.; Sjövall, P.; Thiel, V.; Zheng, W.; Ito, S.; Wakamatsu, K.; Hauff, R.; Kear, B.P.; Engdahl, A.; Alwmark, C.; Eriksson, M.E.; Jarenmark, M.; Sachs, S.; Ahlberg, P.E.; Marone, F.; Kuriyama, T.; Gustafsson, O.; Malmberg, P.; Thomen, A.; Rodríguez-Meizoso, I.; Uvdal, P.; Ojika, M.; Schweitzer, M.H. Soft-tissue evidence for homeothermy and crypsis in a Jurassic ichthyosaur. *Nature* **2018**, *564*, 359–365. 770
23. Ambler, R.P.; Daniel, M. Proteins and molecular paleontology. *Philos. T. Roy. Soc. Lon. B* **1991**, *333*, 381–389. 771
24. Baird, R.F.; Rowley, M.J. Preservation of avian collagen in Australian Quaternary cave deposits. *Palaeontology* **1990**, *33*, 447–451. 772
25. Wotjowicz, A.; Yamauchi, M.; Montella, A.; Bandiera, P.; Sotowski, R.; Ostrowski, K. Persistence of bone collagen cross-links in skeletons of the Nuraghi population living in Sardinia 1500–1200 B.C. *Calcif. Tissue Int.* **1999**, *64*, 370–373. 773
26. Tuross, N. Alterations in fossil collagen. *Archaeometry* **2002**, *44*, 427–434. 774
27. Avci, R.; Schweitzer, M.H.; Boyd, R.D.; Wittmeyer, J.L.; Arce, F.T.; Calvo, J.O. Preservation of bone collagen from the Late Cretaceous period studied by immunological techniques and atomic force microscopy. *Langmuir* **2005**, *21*, 3584–3590. 775
28. Asara, J.M.; Schweitzer, M.H.; Freimark, L.M.; Phillips, M.; Cantley, L.C. Protein sequences from mastodon and *Tyrannosaurus rex* revealed by mass spectrometry. *Science* **2007**, *316*, 280–285. 776
29. Schweitzer, M.H.; Suo, Z.; Avci, R.; Asara, J.M.; Allen, M.A.; Arce, F.T.; Horner, J.R. Analyses of soft tissue from *Tyrannosaurus rex* suggest the presence of protein. *Science* **2007**, *316*, 277–280. 777
30. Schweitzer, M.H.; Zheng, W.; Organ, C.L.; Avci, R.; Suo, Z.; Freimark, L.M.; Lebleu, V.S.; Duncan, M.B.; Vander Heiden, M.G.; Neveu, J.M.; Lane, W.S.; Cottrell, J.S.; Horner, J.R.; Cantley, L.C.; Kalluri, R.; Asara, J.M. Biomolecular characterization and protein sequences of the Campanian hadrosaur *B. canadensis*. *Science* **2009**, *324*, 626–631. 778
31. Lingham-Soliar, T.; Wesley-Smith, J. First investigation of the collagen D-band ultrastructure in fossilized vertebrate integument. *Proc. R. Soc. B* **2008**, *275*, 2207–2212. 779
32. Buckley, M.; Collins, M.; Thomas-Oates, J.; Wilson, J.C. Species identification by analysis of bone collagen using matrix-assisted laser desorption/ionisation time-of-flight mass spectrometry. *Rapid Commun. Mass Spectrom.* **2009**, *23*, 3843–3854. 780
33. Buckley, M.; Kansa, S.W.; Howard, S.; Campbell, S.; Thomas-Oates, J.; Collins, M. Distinguishing between archaeological sheep and goat bones using a single collagen peptide. *J. Archaeol. Sci.* **2010**, *37*, 13–20. 781
34. Buckley, M.; Larkin, N.; Collins, M. Mammoth and *Mastodon* collagen sequences; survival and utility. *Geochim. Cosmochim. Ac.* **2011**, *75*, 2007–2016. 782
35. Buckley, M.; Harvey, V.L.; Orihuela, J.; Mychajliw, A.M.; Keating, J.; Almonte Milan, J.N.; Lawless, C.; Chamberlain, A.T.; Egerton, V.M.; Manning, P.L. Collagen sequence analysis reveals evolutionary history of extinct West Indies *Nesophontes* ('island-shrews'). *Mol. Biol. Evol.* **2020**, *37*, 2931–2943. 783
36. Dobberstein, R.C.; Collins, M.J.; Craig, O.E.; Taylor, G.; Penkman, K.E.H.; Ritz-Timme, S. Archaeological collagen: why worry about collagen diagenesis? *Archaeol. Anthropol. Sci.* **2009**, *1*, 31–42. 784
37. Lindgren, J.; Uvdal, P.; Engdahl, A.; Lee, A.H.; Alwmark, C.; Bergquist, K.; Nilsson, E.; Ekstrom, P.; Rasmussen, M.; Douglas, D.A.; Polyn, M.J.; Jacobs, L.L. Microspectroscopic evidence of Cretaceous bone proteins. *PLoS ONE* **2011**, *6*, e19445. 785
38. San Antonio, J.D.; Schweitzer, M.H.; Jensen, S.T.; Kalluri, R.; Buckley, M.; Orgel, J. Dinosaur peptides suggest mechanisms of protein survival. *PLoS ONE* **2011**, *6*, e20381. 786
39. Zylberberg, L.; Laurin, M. Analysis of fossil bone organic matrix by transmission electron microscopy. *C. R. Palevol* **2011**, *10*, 357–366. 787
40. Wang, S.-Y.; Cappellini, E.; Zhang, H.-Y. Why collagens best survived in fossils? Clues from amino acid thermal stability. *Biochem. Biophys. Res. Co.* **2012**, *422*, 5–7. 788
41. Buckley, M. Ancient collagen reveals evolutionary history of the endemic South American 'ungulates'. *Proc. R. Soc. B* **2015**, *282*, 20142671. 789
42. Cleland, T.P.; Schroeter, E.R.; Feranec, R.S.; Vashishth, D. Peptide sequences from the first *Castoroides ohioensis* skull and the utility of old museum collections for palaeoproteomics. *Proc. R. Soc. B* **2016**, *283*, 20160593. 790
43. Hill, R.C.; Wither, M.J.; Nemkov, T.; Barrett, A.; D'Alessandro, A.; Dzieciatkowska, M.; Hansen, K.C. Preserved proteins from extinct *Bison latifrons* identified by tandem mass spectrometry; hydroxyllysine glycosides are a common feature of ancient collagen. *Mol. Cell. Proteomics* **2015**, *14*, 1946–1958. 791
44. Lee, Y.-C.; Chiang, C.-C.; Huang, P.-Y.; Chung, C.-Y.; Huang, T.D.; Wang, C.-C.; Chen, C.-I.; Chang, R.-S.; Liao, C.-H.; Reisz, R.R. Evidence of preserved collagen in an Early Jurassic sauropodomorph dinosaur revealed by synchrotron FTIR microspectroscopy. *Nat. Commun.* **2017**, *8*, 14220. 792

45. Schroeter, E.R.; DeHart, C.J.; Cleland, T.P.; Zheng, W.; Thomas, P.M.; Kelleher, N.L.; Bern, M.; Schweitzer, M.H. Expansion for the *Brachylophosaurus canadensis* collagen I sequence and additional evidence of the preservation of Cretaceous protein. *J. Proteome Res.* **2017**, *16*, 920–932.
46. Anné, J.; Edwards, N.P.; Brigidi, F.; Gueriau, P.; Harvey, V.L.; Geraki, K.; Slimak, L.; Buckley, M.; Wogelius, R.A. Advances in bone preservation: Identifying possible collagen preservation using sulfur speciation mapping. *Palaeogeog. Palaeoclimatol.* **2019**, *520*, 181–187.
47. Ullmann, P.V.; Voegelé, K.K.; Grandstaff, D.E.; Ash, R.D.; Zheng, W.; Schroeter, E.R.; Schweitzer, M.H.; Lacovara, K.J. Molecular tests support the viability of rare earth elements as proxies for fossil biomolecule preservation. *Sci. Rep.* **2020**, *10*, 15566.
48. Cleland, T.P.; Schroeter, E.R.; Colleary, C. Diagenetiforms: A new term to explain protein changes as a result of diagenesis in paleoproteomics. *J. Proteomics* **2021**, *230*, 103992.
49. Organ, C.L.; Schweitzer, M.H.; Zheng, W.; Freemark, L.M.; Cantley, L.C.; Asara, J.M. Molecular phylogenetics of *Mastodon* and *Tyrannosaurus rex*. *Science* **2008**, *320*, 499.
50. Allison, P.A. The role of anoxia in the decay and mineralization of proteinaceous macro-fossils. *Paleobiology* **1988**, *14*, 139–154.
51. Eglinton, G.; Logan, G.A. Molecular preservation. *Philos. T. R. Soc. Lon. B* **1991**, *333*, 315–328.
52. Colleary, C.; Lamadrid, H.M.; O'Reilly, S.S.; Dolocan, A.; Nesbitt, S.J. Molecular preservation in mammoth bone and variation based on burial environment. *Sci. Rep.* **2021**, *11*, 2662.
53. Boatman, E.M.; Goodwin, M.B.; Holman, H.-Y.N.; Fakra, S.; Zheng, W.; Gronsky, R.; Schweitzer, M.H. Mechanisms of soft tissue and protein preservation in *Tyrannosaurus rex*. *Sci. Rep.* **2019**, *9*, 15678.
54. Wiemann, J.; Fabbri, M.; Yang, T.-R.; Stein, K.; Sander, P.M.; Norell, M.A.; Briggs, D.E.G. Fossilization transforms vertebrate hard tissue proteins into N-heterocyclic polymers. *Nat. Commun.* **2018**, *9*, 4741.
55. Trueman, C.N. Trace element geochemistry of bonebeds. In *Bonebeds: genesis, analysis, and paleobiological significance*; Rogers, R.R.; Eberth, D.A.; Fiorillo, A.R., Eds.; University of Chicago Press: Chicago, IL, USA, **2007**; pp. 397–435.
56. Trueman, C.N.; Behrensmeyer, A.K.; Tuross, N.; Weiner, S. Mineralogical and compositional changes in bones exposed on soil surfaces in Amboseli National Park, Kenya: diagenetic mechanisms and the role of sediment pore fluids. *J. Archaeol. Sci.* **2004**, *31*, 721–739.
57. Suarez, C.A.; Macpherson, G.L.; Gonzalez, L.A.; Grandstaff, D.E. Heterogeneous rare earth element (REE) patterns and concentrations in a fossil bone: implications for the use of REE in vertebrate taphonomy and fossilization history. *Geochim. Cosmochim. Acta* **2010**, *74*, 2970–2988.
58. Suarez, C.A.; Morschhauser, E.M.; Suarez, M.B.; You, H.; Li, D.; Dodson, P. Rare earth element geochemistry of bone beds from the Lower Cretaceous Zhonggou Formation of Gansu Province, China. *J. Vert. Paleo.* **2019**, *38* (Suppl.), 22–35.
59. Herwartz, D.; Tütken, T.; Munker, C.; Jochum, K.P.; Stoll, B.; Sander, P.M. Timescales and mechanisms of REE and Hf uptake in fossil bones. *Geochim. Cosmochim. Acta* **2011**, *75*, 82–105.
60. Herwartz, D.; Munker, C.; Tütken, T.; Hoffmann, J.E.; Wittke, A.; Barbier, B. Lu-Hf isotope systematics of fossil biogenic apatite and their effects on geochronology. *Geochim. Cosmochim. Acta* **2013**, *101*, 328–343.
61. Kowal-Linka, M.; Jochum, K.P.; Surmik, D. LA-ICP-MS analysis of rare earth elements in marine reptile bones from the Middle Triassic bonebed (Upper Silesia, S Poland): impact of long-lasting diagenesis, and factors controlling the uptake. *Chem. Geol.* **2014**, *363*, 213–228.
62. Decrée, S.; Herwartz, D.; Mercadier, J.; Miján, I.; de Buffrénil, V.; Leduc, T.; Lambert, O. The post-mortem history of a bone revealed by its trace element signature: the case of a fossil whale rostrum. *Chem. Geol.* **2018**, *477*, 137–150.
63. Ullmann, P.V.; Grandstaff, D.E.; Ash, R.D.; Lacovara, K.J. Geochemical taphonomy of the Standing Rock Hadrosaur Site: exploring links between rare earth elements and cellular and soft tissue preservation. *Geochim. Cosmochim. Acta* **2020**, *269*, 223–237.
64. Ullmann, P.V.; Macauley, K.; Ash, R.D.; Shoup, B.; Scannella, J.B. Taphonomic and diagenetic pathways to protein preservation, part I: the case of *Tyrannosaurus rex* specimen MOR 1125. *Biology* **2021**, *10*, 1193.
65. McCormack, J.M.; Bahr, A.; Gerdes, A.; Tütken, T.; Prinz-Grimm, P. Preservation of successive diagenetic stages in Middle Triassic bonebeds: evidence from *in situ* trace element and strontium isotope analysis of vertebrate fossils. *Chem. Geol.* **2015**, *410*, 108–123.
66. Grimstead, D.N.; Clark, A.E.; Rezac, A. Uranium and vanadium concentrations as a trace element method for identifying diagenetically altered bone in the inorganic phase. *J. Archaeol. Method. Theory* **2018**, *25*, 689–704.
67. Kocsis, L.; Gheerbrant, E.; Mouflih, M.; Cappetta, H.; Ulianov, A.; Chiaradia, M.; Bardet, N. Gradual changes in upwelled seawater conditions (redox, pH) from the late Cretaceous through early Paleogene at the northwest coast of Africa: negative Ce anomaly trend recorded in fossil bio-apatite. *Chem. Geol.* **2016**, *421*, 44–54.
68. Trueman, C.N.; Palmer, M.R.; Field, J.; Privat, K.; Ludgate, N.; Chavagnac, V.; Eberth, D.A.; Cifelli, R.; Rogers, R.R. Comparing rates of recrystallisation and the potential for preservation of biomolecules from the distribution of trace elements in fossil bones. *C. R. Palevol* **2008**, *7*, 145–158.
69. Paul, G.S.; Persons, W.S.; Van Raalte, J. The tyrant lizard king, queen and emperor: multiple lines of morphological and stratigraphic evidence support subtle evolution and probable speciation within the North American genus *Tyrannosaurus*. *Evol. Biol.* **2022**, *49*, 156–179.

70. Schweitzer, M.H.; Moyer, A.E.; Zheng, W. Testing the hypothesis of biofilm as a source for soft tissue and cell-like structures preserved in dinosaur bone. *PLoS ONE* **2016**, *11*, e0150238. 884
71. LaRock, J.W. Sedimentology and taphonomy of a dinosaur bonebed from the Upper Cretaceous (Campanian) Judith River Formation of north central Montana. M.S. thesis, Montana State University, Bozeman, 2000. 885
72. Horner, J.R.; de Ricqlès, A.; Padian, K. Variation in dinosaur skeletochronology indicators: implications for age assessment and physiology. *Paleobiology* **1999**, *25*, 295–304. 886
73. Horner, J.R.; De Ricqlès, A.; Padian, K. Long bone histology of the hadrosaurid dinosaur *Maiasaura peeblesorum*: growth dynamics and physiology based on an ontogenetic series of skeletal elements. *J. Vertebr. Paleontol.* **2000**, *20*, 115–129. 887
74. Haskin, L.A.; Haskin, M.A.; Frey, F.A.; Wildeman, T.R. Relative and absolute terrestrial abundances of the rare earths. In *Origin and Distribution of the Elements*; Ahrens, L.H., Ed.; Pergamon: Oxford, UK, 1968; pp. 889–912. 888
75. Gromet, L.P.; Dymek, R.F.; Haskin, L.A.; Korotev, R.L. The "North American shale composite": its compilation, major and trace element characteristics. *Geochim. Cosmochim. Ac.* **1984**, *48*, 2469–2482. 889
76. Kohn, M.J. Models of diffusion-limited uptake of trace elements in fossils and rates of fossilization. *Geochim. Cosmochim. Ac.* **2008**, *72*, 3758–3770. 890
77. Kemp, R.A.; Trueman, C.N. Rare earth elements in Solnhofen biogenic apatite: geochemical clues to the paleoenvironment. *Sediment. Geol.* **2003**, *155*, 109–127. 891
78. Smirnova, E.V.; Mysovskaya, I.N.; Lozhkin, V.I.; Sandimirova, G.P.; Pakhomova, N.N.; Smagunova, A.A. Spectral interferences from polyatomic barium ions in inductively coupled plasma mass spectrometry. *J. Appl. Spectrosc.* **2006**, *73*, 813–818. 892
79. Grandstaff, D.E.; Terry, D.O. Rare earth element composition of Paleogene vertebrate fossils from Toadstool Geologic Park, Nebraska, USA. *Appl. Geochem.* **2009**, *24*, 733–745. 893
80. Bau, M.; Dulski, P. Anthropogenic origin of positive gadolinium anomalies in river waters. *Earth Planet. Sc. Lett.* **1996**, *143*, 245–255. 894
81. Pack, A.; Russell, S.S.; Shelley, J.M.G.; van Zuilen, M. Geo- and cosmochemistry of the twin elements yttrium and holmium. *Geochim. Cosmochim. Ac.* **2007**, *71*, 4592–4608. 895
82. Behrensmeyer, A.K. Taphonomic and ecologic information from bone weathering. *Palaeogeogr. Palaeocl.* **1978**, *4*, 150–162. 896
83. Hill, A.; Behrensmeyer, A.K. Disarticulation patterns of some modern East African mammals. *Paleobiology* **1984**, *10*, 366–376. 897
84. Brand, L.R.; Hussey, M.; Taylor, J. Decay and disarticulation of small vertebrates in controlled experiments. *J. Taphonomy* **2003**, *1*, 69–95. 898
85. Rogers, R.R. Taphonomy of three dinosaur bone beds in the Upper Cretaceous Two Medicine Formation of northwestern Montana: evidence for drought-related mortality. *PALAIOS* **1990**, *5*, 394–413. 899
86. Payne, J.A. A summer carrion study of the baby pig *Sus scrofa* Linnaeus. *Ecology* **1965**, *46*, 592–602. 900
87. Syme, C.E.; Salisbury, S.W. Patterns of aquatic decay and disarticulation in juvenile Indo-Pacific crocodiles (*Crocodylus porosus*), and implications for the taphonomic interpretation of fossil crocodyliform material. *Palaeogeogr. Palaeocl.* **2014**, *412*, 108–123. 901
88. Rogers, R.R.; Kidwell, S.M. Associations of vertebrate skeletal concentrations and discontinuity surfaces in terrestrial and shallow marine records: a test in the Cretaceous of Montana. *J. Geol.* **2000**, *108*, 131–154. 902
89. Rogers, R.R.; Kidwell, S.M.; Deino, A.L.; Mitchell, J.P.; Nelson, K.; Thole, J.T. Age, correlation, and lithostratigraphic revision of the Upper Cretaceous (Campanian) Judith River Formation in its type area (north-central Montana), with a comparison of low- and high-accommodation alluvial records. *J. Geol.* **2016**, *124*, 99–135. 903
90. Trueman, C.N. Rare earth element geochemistry and taphonomy of terrestrial vertebrate assemblages. *PALAIOS* **1999**, *14*, 555–568. 904
91. Suarez, C.A.; Suarez, M.B.; Terry, D.O.; Grandstaff, D.E. Rare earth element geochemistry and taphonomy of the Early Cretaceous Crystal Geyser Dinosaur Quarry, east-central Utah. *PALAIOS* **2007**, *22*, 500–512. 905
92. Rogers, R.R.; Fricke, H.C.; Addona, V.; Canavan, R.R.; Dwyer, C.N.; Harwood, C.L.; Koenig, A.E.; Murray, R.; Thole, J.T.; Williams, J. Using laser ablation-inductively coupled plasma-mass spectrometry (LA-ICP-MS) to explore geochemical taphonomy of vertebrate fossils in the Upper Cretaceous Two Medicine and Judith River Formations of Montana. *PALAIOS* **2010**, *25*, 183–195. 906
93. Suarez, C.A.; Kohn, M.J. Caught in the act: a case study on microscopic scale physicochemical effects of fossilization on stable isotopic composition of bone. *Geochim. Cosmochim. Ac.* **2020**, *268*, 277–295. 907
94. Cantrell, K.J.; Byrne, R.H. Rare earth element complexation by carbonate and oxalate ions. *Geochim. Cosmochim. Ac.* **1987**, *51*, 597–605. 908
95. Zhang, J.; Létolle, R.; Jusserand, C. Major element chemistry of the Huanghe (Yellow River), China – weathering processes and chemical fluxes. *J. Hydrol.* **1995**, *168*, 173–203. 909
96. Luo, Y.-R.; Byrne, R.H. Carbonate complexation of yttrium and the rare earth elements in natural waters. *Geochim. Cosmochim. Ac.* **2004**, *68*, 691–699. 910
97. Pourret, O.; Davranche, M.; Gruau, G.; Dia, A. Rare earth elements complexation with humic acid. *Chem. Geol.* **2007**, *243*, 128–141. 911
98. Byrne, R.H.; Liu, X.; Schijf, J. The influence of phosphate coprecipitation on rare earth distributions in natural waters. *Geochim. Cosmochim. Ac.* **1996**, *60*, 3341–3346. 912
99. Föllmi, K.B. The phosphorous cycle, phosphogenesis and marine phosphate-rich deposits. *Earth-Sci. Rev.* **1996**, *40*, 55–124. 913

100. Liu, X.; Byrne, R.H.; Schijf, J. Comparative coprecipitation of phosphate and arsenate with yttrium and the rare earths: the influence of solution complexation. *J. Solution Chem.* **1997**, *26*, 1187–1198. 944
101. Brunet, R.-C.; Astin, K.B. Variation in phosphorus flux during a hydrological season: the River Ardour. *Wat. Res.* **1998**, *32*, 547–558. 945
102. Wings, O. Authigenic minerals in fossil bones from the Mesozoic of England: poor correlation with depositional environments. *Palaeogeogr. Palaeoclimatol.* **2004**, *204*, 15–32. 946
103. Janssens, K.; Vincze, L.; Vekemans, B.; Williams, C.T.; Radtke, M.; Haller, M.; Knochel, A. The non-destructive determination of REE in fossilized bone using synchrotron radiation induced K-line X-ray microfluorescence analysis. *J. Anal. Chem.* **1999**, *363*, 413–420. 947
104. Hinz, E.A.; Kohn, M.J. The effect of tissue structure and soil chemistry on trace element uptake in fossils. *Geochim. Cosmochim. Ac.* **2010**, *74*, 3213–3231. 948
105. Trueman, C.N.; Kocsis, L.; Palmer, M.R.; Dewdney, C. Fractionation of rare earth elements within bone mineral: a natural cation exchange system. *Palaeogeogr. Palaeoclimatol.* **2011**, *310*, 124–132. 949
106. Kohn, M.J.; Moses, R.J. Trace element diffusivities in bone rule out simple diffusive uptake during fossilization but explain in vivo uptake and release. *P. Natl. Acad. Sci. USA* **2013**, *110*, 419–424. 950
107. Elderfield, H.; Upstill-Goddard, R.; Sholkovitz, E.R. The rare earth elements in rivers, estuaries, and coastal seas and their significance to the composition of ocean waters. *Geochim. Cosmochim. Ac.* **1990**, *54*, 971–991. 951
108. Williams, C.T.; Potts, P.J. Element distribution maps in fossil bones. *Archaeometry* **1988**, *30*, 237–247. 952
109. Williams, C.T.; Henderson, P.; Marlow, C.A.; Molleson, T.I. The environment of deposition indicated by the distribution of rare earth elements in fossil bones from Olduvai Gorge, Tanzania. *Appl. Geochem.* **1997**, *12*, 537–547. 953
110. Metzger, C.A.; Terry, D.O.; Grandstaff, D.E. Effect of paleosol formation on rare earth element signatures in fossil bone. *Geology* **2004**, *32*, 497–500. 954
111. Martin, J.E.; Patrick, D.; Kihm, A.J.; Foit, F.F.; Grandstaff, D.E. Lithostratigraphy, tephrochronology, and rare earth element geochemistry of fossils at the classical Pleistocene Fossil Lake area, south central Oregon. *J. Geol.* **2005**, *113*, 139–155. 955
112. Ferrante, C.; Cavin, L.; Vennemann, T.; Martini, R. Histology and geochemistry of *Allosaurus* (Dinosauria: Theropoda) from the Cleveland-Lloyd Dinosaur Quarry (Late Jurassic, Utah): paleobiological implications. *Front. Earth Sci.* **2021**, *9*, 641060. 956
113. Schweitzer, M.H. Molecular paleontology: some current advances and problems. *Ann. Paleontol.* **2004**, *90*, 81–102. 957
114. Ullmann, P.V.; Pandya, S.H.; Nelleremoe, R. Patterns of soft tissue and cellular preservation in relation to fossil bone microstructure and overburden depth at the Standing Rock Hadrosaur Site, Maastrichtian Hell Creek Formation, South Dakota, USA. *Cretaceous Res.* **2019**, *99*, 1–13. 958
115. Allison, P.A. Decay processes. In *Paleobiology: A Synthesis*; Briggs, D.E.G.; Crowther, P.R., Eds.; Blackwell Scientific Publications, Oxford, **1990**; pp. 213–216. 959
116. Martill, D.M. Macromolecular resolution of fossilized muscle tissue from an elopomorph fish. *Nature* **1990**, *346*, 171–172. 960
117. Hubert, J.F.; Panish, P.T.; Chure, D.J.; Probst, K.S. Chemistry, microstructure, petrology, and diagenetic model of Jurassic dinosaur bones, Dinosaur National Monument, Utah. *J. Sediment. Res.* **1996**, *66*, 531–547. 961
118. Pfretzschner, H.-U. Pyrite in fossil bone. *N. Jb. Geol. Paläont. Abh.* **2001**, *220*, 1–23. 962
119. Franzen, J.L. Exceptional preservation of Eocene vertebrates in the Lake Deposit of Grube Messel (West Germany). *Philos. T. Roy. Soc. B Lon.* **1985**, *311*, 181–186. 963
120. McNamara, M.E.; Orr, P.J.; Kearns, S.L. High-fidelity organic preservation of bone marrow in ca. 10 Ma amphibians. *Geology* **2006**, *34*, 641–644. 964
121. McNamara, M.E.; Orr, P.J.; Kearns, S.L.; Alcalá, L.; Anadón, P.; Penalver Mollar, E. Soft-tissue preservation in Miocene frogs from Libros, Spain: insights into the genesis of decay microenvironments. *PALAIOS* **2009**, *24*, 104–117. 965
122. Manning, P.; Morris, P.M.; McMahon, A.; Jones, E.; Gize, A.; Macquaker, J.H.S.; Wolff, G.; Thompson, A.; Marshall, J.; Taylor, K.G.; Lyson, T.; Gaskell, S.; Reamtong, O.; Sellers, W.I.; Van Dongen, B.E.; Buckley, M.; Wogelius, R.A. Mineralized soft-tissue structure and chemistry in a mummified hadrosaur from the Hell Creek Formation, North Dakota (USA). *Proc. R. Soc. B* **2009**, *276*, 3429–3437. 966
123. Schroeter, E.R.; Cleland, T.P.; Schweitzer, M.H. Deep time paleoproteomics: looking forward. *J. Proteome Res.* **2022**, *21*, 9–19. 967