

Comprehensive assessment of plant and water productivity responses in negative pressure irrigation technology: A meta-analysis

Yuwen Jin¹, Xiang Gao¹, Renlian Zhang¹, Xueping Wu¹, Huaiyu Long¹, Zhimei Sun² and Shuxiang Zhang^{1,*}

¹ Institute of Agricultural Resources and Regional Planning, Chinese Academy of Agricultural Sciences, Beijing 100081, China; 1525980610@qq.com (Y.J.); gxkochung@gmail.com (X.G.); zhangrenlian@caas.com (R.Z.); wuxueping@caas.com (X.W.); longhuaiyu@caas.com (H.L.)

² College of Resources and Environmental Sciences, Hebei Agricultural University, Baoding 071000, China; sunzhm2002@163.com (Z.S.)

* Correspondence: zhangshuxiang@caas.cn.

Yuwen Jin and Xiang Gao contributed equally to this work.

Abstract: Negative pressure irrigation (NPI) is an important water management strategy that can improve crop yields and water use efficiency (WUE). However, because NPI is affected by vital factors such as the negative pressure values, soil properties and fertilization dosages, there is a lack of systematic analysis on the application effects of NPI on various crops. Hence, this study collected the results of 44 published studies and established valid 142 crop yields, 121 WUEs, 138 crop qualities, and 138 crop nutrient status in a database for NPI systems. The meta-analysis method was used to analyze NPI in comparison to conventional irrigation (CI) conditions. The results showed that NPI yields and WUEs significantly improved by 17% and 63% compared to CI, respectively. Meanwhile, the negative pressure values were -2 ~ -5 kPa, the improvement effects on yields were the best; and the WUEs exhibited the highest performance with negative pressure values of -6 ~ -10 kPa. NPI promoted crop quality and plant nutrient uptakes under appropriate NPI conditions. The synergistic impacts for sandy loam, alkaline soils and leafy vegetables were greater than for clay loam, neutral soils, and fruit vegetables under NPI conditions. Simultaneously, it was shown that soil available phosphorus content and application of P fertilizer have a greater impact on NPI and CI crop yields. Therefore, the meta-analysis demonstrated the impacts of NPI on crop yields, WUEs, quality and nutrient absorption, and quantified the effects of NPI on crop growth under various conditions, which provides an important water-saving technology for greenhouse production.

Citation: *Agronomy* **2022**, *12*, x.
<https://doi.org/10.3390/xxxxx>

Academic Editor:

Received: date

Accepted: date

Published: date

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Keywords: Negative pressure irrigation; Yield; Quality of plant; Water use efficiency; Fertilization

1. Introduction

The growth and development of crops depends on the absorption of large amounts of water, and artificial irrigation is the main source of water to satisfy crop water requirements [1–3]. Conventional irrigation (CI) systems, such as furrows, depressions, and flood irrigation, have relatively low water use efficiencies (WUE), while advanced water-saving technologies, including sprinklers, drip, microspray, and alternate root zone irrigation, provide significantly improved WUEs [4–7]. The introduction of irrigation water into farmland requires human labor and equipment, and the process for water to enter soils depends entirely on human irrigation [3,8]. Crop and soil systems are consistently in a state of passively receiving irrigation water [5,9]. In the past 20 years, negative pressure irrigation (NPI) technology has been developed, which is based on the active absorption of soil water by crops, and water enters the soil under the driving forces of the physiological activities of crop transpiration and water consumption characteristics, which can

continuously and smoothly supply crops with water for its absorption and utilization from soil [5,10–12].

NPI is a new type of subsurface irrigation technology, which is also known as active water absorption by crops. The processes of soil supplying water to crop roots, water supply time, intensity and water consumption mainly depend on the physiological needs of dryland crops [13–15]. Due to transpiration and the absorption of soil water by root systems, which cause the soil matrix potential to have a gradient between the soil and emitter, the water absorption force spreads out through the soil and attracts water away from the rhizosphere to move into root systems [3,10,16]. When the water absorption force of the crops is less than or equal to the external water suction force, crops suspend their absorption of water from the soil, and the outside biotope stops supplying water to the soil [5,15,17]. This ensures that crops can be supplied in a stable manner with suitable water amounts for growth in NPI systems. Since the water supply process is regulated by the controllable negative pressure that is generated in the system, the technology is named “negative pressure” [3,5,18].



At present, NPI has been used in practical research in greenhouse production, which is mainly manifested by the fact that dryland crops have suitable pressure values (e.g., within the range of -2 to -10 kPa) [3,8,19]. For example, the yields of rape, pepper, cucumber, lettuce, tomato and other crops under suitable negative pressure conditions have shown significant increases of 29% to 77%, and the WUEs have increased by 35% to 166% compared with CI [3,20,21]. NPI also improves the absorption of nutrients and utilization efficiency of nutrients by crops [15–17]. In addition, NPI affects the soil nutrient availabilities and spatial distributions, soil microbial diversities, and soil and plant enzyme activities [9,11,22,23]. The study also found that NPI increased the microbial diversities in the rhizospheres of rape and tomato (OUT, Chao1, Shannon) and the activities of catalase, urease, and phosphatase in pepper and tomato soils [9,24,25], but decreased the levels of the superoxide anion, malondialdehyde, proline, catalase and polyphenol oxidase in maize under NPI conditions [26]. This is proof enough that NPI has emerged as a key potential solution for reducing water extraction in the context of increasing agricultural intensification and climate change. Given its importance, Yang et al (2022) systematically reviewed the development of NPI in recent years through literature analysis [8]. However, the study lacks quantitative analysis of crop yield and WUE by NPI. In addition, although NPI has provided many research achievements, the results occasionally appear to be discrepancies or contradictions, but there is a lack of independent systematic combined analyses of results with common research destinations, analyses of the diversity among studies, and quantitative comprehensive evaluations of research results.

Previous studies have focused on the effects of different negative pressure values on crop growth and WUEs. The results indicated that different crops favor different negative pressure values of NPI. In fact, the efficiency of NPI is also affected by soil texture, soil property, water and fertilizer applications, which cannot be ignored. To comprehensively understand the comprehensive effect of NPI on crop growth and its relationship with soil physicochemical properties, published experimental data were collected for a synthetic analysis between NPI and crop growth and were systematically quantified. The data collection step was carried out by using the published articles related to NPI, and the meta-analysis method was used to systematically analyze the effects of NPI on crop yield and quality, water use efficiency, and nutrient absorption and to clarify the negative pressure values. According to the existing data that were mined, the appropriate fertilization dosages under NPI were quantitatively recommended and would provide a theoretical basis for the development of NPI technology.

2. Materials and Methods

2.1. Data collection

We collected data from published papers by using the Web of Science (<https://www.webofscience.com>) and China National Knowledge Infrastructure (<https://www.cnki.net>) databases between 2000 and 2021 and used the following 4 search terms: “Negative pressure irrigation”, “Negative emitter irrigation”, “Negative pressure water supply technology”, and “Negative emitter water supply”. The literature screening was conducted based on the following conditions: (1) the experiment was conducted by using potted and greenhouse experiments; (2) the same experiment must include both an NPI treatment and conventional irrigation treatment and provide yield or water use efficiency or quality or crop nutrient status index data; and (3) the basic physical and chemical indices for the 0–20 cm soil layer are included in the literature. A total of 44 peer-reviewed papers spanning valid 142 sets of crop yield data, 121 sets of water use efficiency data, 138 sets of crop quality data, and 138 sets of crop nutrient status data were included in this meta-analysis.



Schematic diagram of the negative pressure irrigation system designed by the Chinese Academy of Agricultural Sciences [5,9] (Figure 1a). Figure 1b and 1c show the cultivation of cucumber and rapeseed with NPI in greenhouse production, respectively [3,9,25].




Figure 1. Schematic of plant growth when using a negative pressure irrigation system (a). Negative pressure irrigation applied to cucumber (b) and rapeseed (c) in greenhouse production.

2.2. Data analysis

In the meta-analysis method, the standard deviations and number of repetitions of the experimental group and control group are very important parameters. If the original literature clearly mentions the sample sizes, mean and standard deviations (or standard errors), the data were directly extracted; if the standard deviations were not provided directly in the literature, these studies were processed in the following two ways: (1) The data for each treatment are listed. When there were multiple replicates of experimental

data, the experimental data were sorted with Excel to obtain the means and standard deviations. (2) The corresponding standard deviations are estimated based on the ratios of the existing standard deviations to the means [27]. If there only graphs of the above data parameters are presented in the literature, GetData Graph Digitizer software was used for data extraction.

2.3. Data calculation

Only the standard error (SE) was provided in the literature, which was converted to the standard deviation (SD) using Formula (1):

$$SD = SE \times \sqrt{n} \quad (1)$$

The statistical indicators were expressed as the response ratio (RR) and calculated by Formula (2) and the logarithm of the response ratio (lnRR) to reflect the effect of negative pressure irrigation on conventional irrigation:

$$RR = \frac{A_t}{A_c} \quad (2)$$

$$\ln RR = \frac{\ln A_t}{\ln A_c} = \ln A_t - \ln A_c \quad (3)$$

The sampling variance for each lnRR was calculated as Formula (4)

$$V = \frac{SD_t^2}{A_t^2 n_t} + \frac{SD_c^2}{A_c^2 n_c} \quad (4)$$

In the formula, SD_t, A_t, and n_t are the standard deviation, mean, and number of samples of the experimental group, respectively; SD_c, A_c, and n_c are the standard deviation, mean, and number of samples of the control group, respectively.

To facilitate the interpretation, the change percentage of crop yield, WUE, quality and nutrient uptake under NPI treatment compared with CI treatment was calculated by the following formula :

$$\text{Effect size (\%)} = (\exp(\ln RR) - 1) \times 100 \quad (5)$$

The meta analysis was performed including the studies as random factors using the rma function in metafor (setting method "REML"), which was necessary to reduce data dispersion due to the experimental design or the measuring method.

If the 95% confidence interval does not include zero points, this means that NPI has a significant effect when compared with conventional irrigation; otherwise, this means that there was no significant effect; if all values fall on the negative semiaxis, this means that negative pressure irrigation provided a negative effect; otherwise, this means that there was a positive effect. To evaluate the quality of our meta analysis, we determined Egger's tests. The results show that there is no publication bias in all data ($p > 0.05$, Table 1). We also used the Leave1out function in metafor for sensitivity analysis of meta-analysis results, and the results showed that the conclusions of this study were relatively reliable.

Table 1. Publication bias test of relevant data - Egger's test.

	Sample size (n)	Z value	P value
Crop yields	135	0.5472	0.5842
Water use efficiencies	113	-0.1597	0.8731
Crop quality	132	0.7764	0.4375
Nutrient content	54	1.7533	0.0796
Nutrient concentration	63	-0.9302	0.3523

Note: $p > 0.05$ indicates that there is no publication bias.

The above data processing was performed in R 4.1.0, and Origin 2019b was used for graphing.

3. Results

3.1. Effect size on crop yield under NPI

As shown in Figure 2, NPI significantly improved by 17.4% crop yields compared to conventional irrigation (CI) (all effect sizes and their 95% confidence). However, the crop yield responses showed discrepancies for various levels of negative pressure values, and crop yields were noticeably improved for the ranges of -2 ~ -5 kPa and -6 ~ -10 kPa. Among them, -2 ~ -5 kPa provided the best yield increase (41.6%), while negative pressure values of -12 ~ -15 kPa remarkably reduced crop yields (-26.2%). The yield-promoting effects of NPI on a variety of crops were different; leafy vegetables had the best yield increase (30.2%), which were followed by fruit vegetables (8.0%).




Figure 2. Effect size analysis of various factors on crop yields under negative pressure irrigation compared with conventional irrigation conditions.

The soil impacts on crop yields vary according to texture (Figure 2). NPI produces a noteworthy yield increase for sandy loam (22.7%), but not for clay loam. Simultaneously, NPI boosts crop yields in alkaline soils (16.8%), but the effect size was not seen for neutral soils. The effect of NPI on crop yields increased with decreasing soil organic matter contents; the soil organic matter contents were less than 20 g kg⁻¹, and NPI prominently increased crop yields, especially the organic matter contents, which ranged from 0~10 g kg⁻¹ (22.9%). The soil available phosphorus contents ranged from 0~20 mg kg⁻¹, and NPI significantly increased crop yields, while this effect was not significant in other areas. When soil total nitrogen content was 0 ~ 1 g kg⁻¹, NPI significantly increased crop yield (17.0%).

3.2. Effect size on water use efficiency under NPI

NPI significantly increased by 62.6% the WUEs of crops under various factors (Figure 3). Different from the yields, the WUEs are associated with minor differences under various negative pressure values, with the highest WUEs occurring for the range of -6 ~ -10 kPa (75.1%), which are followed by -12 ~ -15 kPa (67.3%) and then by -2 ~ -5 kPa (53.5%). The WUE results for various crop species were not consistent with the yields, and leafy vegetables (72.8%) were slightly better than fruit vegetables. The WUEs were higher in sandy loam soils than clay loam soils, which were consistent with the yield results. NPI had the greatest impact on the WUEs in alkaline soils (pH > 7.5). The effects of the soil organic matter, available phosphorus and total N changed significantly with the difference in content under NPI conditions. NPI could significantly increase WUE under different organic matter content, and the increase was the best under low organic matter content (0~10 g kg⁻¹). The subgroups of available P and total N showed the same trend of results as organic matter.




Figure 3. Effect size analysis of various factors on water use efficiencies under negative pressure irrigation compared with conventional irrigation conditions.

3.3. Crop quality and nutrient uptake on NPI

The quality indicators used in this study were the soluble sugar content, Vc, nitrate, and soluble protein, as shown in Figure 4. NPI markedly increased the crop soluble sugar and Vc contents, and the effect size was 15.0% and 16.5%, respectively. The nitrate contents were significantly reduced 11.7%. However, the effects were not significant for soluble proteins. NPI also significantly promoted the uptakes of the nitrogen (N), phosphorus (P), and potassium (K) contents and concentrations in plants.




Figure 4. Effect size analysis of various factors on crop quality and nutrient uptake under negative pressure irrigation compared with **conventional irrigation conditions**.

3.4. Influencing factors analysis of yield under NPI and CI

In order to evaluate the effect of NPI application, the random forest model in the R Programming Language was used to analyze the influence of various soil factors on crop yield (Figure 5). The variance explanation degrees of the nine factors included in the analysis on NPI and CI crop yield were 76.98% and 93.68%, respectively, indicating that these nine factors could well explain the impact on crop yield.




Figure 5. Influencing factors analysis under NPI (a) and CI (b). AP represents available soil phosphorus, PF represents application amount of phosphorus fertilizer, TN represents total soil nitrogen, NF represents application amount of nitrogen fertilizer, KF represents application amount of potassium fertilizer, AK represents available soil potassium, SOM represents soil organic matter, and SBD represents soil bulk density.

The major factors affecting yield under NPI were soil available phosphorus, phosphorus fertilizer application and soil pH, and soil organic matter (SBD) had the least effect (Figure 5a). Potassium and phosphorus fertilizer application were the main factors affecting CI yield, followed by soil available phosphorus (Figure 5b). It suggestion that soil available phosphorus content and application of P fertilizer have a greater impact on NPI and CI crop yields, while SBD has the least impact.

4. Discussion

4.1. NPI improves crop yields and WUEs

The advantage of NPI is that it can generate a stable water supply, assist the soil water contents to remain within a stable range, and ensure that crops take up water according to their physiological needs [5,21,26]. Determining a suitable water supply pressure value is a basic key approach when applying NPI. Through the meta-analysis, it was found that the crop yields generally showed trends of first increasing and then decreasing with supply pressure reductions (the absolute value of the pressure value increased) on a negative pressure generator (Figure 2). The meta-analysis indicated that negative pressure values of -2 ~ -5 kPa had the best performance with respect to yields, and negative pressure values of -12 ~ -15 kPa were too low. The crops exhibited symptoms of water shortage, which resulted in reduced yields. This is primarily because the negative pressure water supply devices are buried in the soil without water saturation, and the pressure of the external water is lower than the atmospheric pressure, which limits the supply of water [3]. For example, when the negative pressure value was -5 kPa, cabbage yields increased by 51% compared with a value of -10 kPa [28], and cucumber yields increased by 45% at -5 kPa compared to 0 kPa [29]. The water and fertilizer supply modes of NPI are conducive to the uptake of water and fertilizer by root systems through the soil, which thereby improve the operation and accumulation of substances assimilated in plants [14,30]. An appropriate water supply increases the biomass of crops, improves photosynthesis, and improves the ability of roots to absorb soil nutrients [9,25]. Therefore, it is meaningful to discuss crop yields and WUEs under suitable pressure values.

In the 121 sets of data analyzed, the WUEs significantly improved under NPI (Figure 3). For example, tomato and rapeseed decreased their water consumption levels by 20% and 23% compared to CI, respectively, while their WUEs also improved [9,15]. Furthermore, negative pressure values in the range of -2 to -15 kPa could improve the WUE, while the range of -6 ~ -10 kPa resulted in the best performance, but the WUEs had a biased response on crop yields (Figure 3). However, it was difficult to achieve the effect of the lowest irrigation amount and highest yield and quality. For example, when the negative pressure is -5 kPa, the WUE was 11% higher than that at 0 kPa [29]; simultaneously, for pepper, the WUE was 32% higher at -5 kPa than at -10 kPa [31]. In an experiment conducted on Chinese cabbage, the yields increased by 32% at -5 kPa compared to -10 kPa, but the WUEs decreased by 52% [28]. The fundamental reason is that crops take up water according to their physiological needs by maintaining unsaturated water in the rhizosphere and by reducing surface evaporation and deep leakage loss of water from the soil [9,12,21]. Simultaneously, NPI is able to transport water directly to the rhizosphere to accurately and constantly supply water to crops [5,19], and different negative pressure values result in different WUEs. In addition, the inconsistent effect of NPI on crop yields and WUEs may be because leafy vegetables have shorter growth periods, consume less water and have lower biomass than fruit vegetables, which may explain why the WUEs of leaf vegetables are slightly lower than those of fruit vegetables (Figure 3).

Soil texture is closely related to the distribution and transport of water and nutrients in the soil layer under NPI conditions [32,33]. There are significant positive impacts on sandy loam, alkaline soils ($\text{pH} > 7.5$), soils with low organic matter contents (e.g., $0\sim 10 \text{ g kg}^{-1}$), and available P levels of $10\sim 20 \text{ mg kg}^{-1}$ under NPI conditions (Figs. 2 and 3). Mainly, the cumulative infiltration, horizontal and vertical distances, and wet volumes of clay loam soils under NPI were significantly higher than those for sandy loam soils [34]. For the same pressures and water supply times, the same results were also observed [32]. However, soil layers with higher clay contents are prone to salt accumulation at their interfaces [35]. For the same soil matrix potential level, clay loam soils have better water and fertilizer retention capacities than sandy loam soils, and the soil volumetric water contents and water uptake capacities are greater [3,22]. The characteristics of NPI supplying water from the rhizosphere can significantly improve the low infiltration of sandy loam soils and increase the WUEs of sandy loam.

The analysis of the 72 groups of collected data concluded that sandy loam soils exhibit more obvious impacts on increasing yields under NPI conditions (Figure 2). The higher air permeabilities of sandy loam soils are more conducive to microbial activity. NPI significantly increases the numbers and activities of soil microbial populations and promotes the decomposition of organic matter such as humus [3,9,24]. NPI simultaneously stimulates the growth of rhizosphere soil microorganisms and decomposition of soil humus, which is beneficial to the transformation, absorption and utilization of nutrients [36]. The pH values of the applied water-soluble fertilizers are approximately 5.5, which makes it easier to neutralize the pH values in alkaline soils, and the root systems can adjust the soil pH after nutrient uptake, which avoids reductions in pH values in acidic soils. However, soils with low organic matter contents and available P levels can significantly increase crop yields under NPI, which is mainly because NPI directly supplies water and nutrients to the rhizospheres of crops, which can be actively absorbed and utilized by the root systems and avoids using traditional irrigation methods that drench the soil surface [9,21], which results in the loss and volatilization of nutrients, so low-nutrient soil environments are suggested to be more conducive to applications of NPI in greenhouse production.

The root systems of leafy vegetables are not strongly developed, and they can only take up shallow soil water, which is easily lost by evaporation, but NPI mainly supplies water at depths of 15 to 20 cm from the soil surface [9,17,21]. This is just enough to meet the water uptake by the root systems in this area, so the effect of increasing the yields of leaf vegetables is better than for fruit vegetables, such as cabbage and tomato [21,28]. Another consideration is that leafy vegetables consume less water and biomass than fruit vegetables, which results in higher yield increase ratios than fruit vegetables.

4.2. NPI benefits crop quality and nutrient uptake

Crop quality can be improved by reasonably controlling the soil water contents to provide a suitable water environment for crops under NPI conditions [9,17,19,25]. For example, the soil water contents of rapeseed under NPI were relatively stable, with a range of 9.7% to 11.7%, compared with the soil water contents that appear to vary greatly with CI, with a range of 8.6% to 13.3% and result in improved quality and N, P and K contents in rapeseed of 21% ~ 57% [9]. Meanwhile, the vitamin C and soluble sugar contents of pakchoi, lettuce and tomato increased by 23% to 35% with NPI compared to CI [3,15,37]. This is because appropriate water and nutrient contents are beneficial for increasing the stomatal conductances and chlorophyll contents of plants, reducing osmoregulation in the pericarp, promoting vitamin C, and increasing the concentrations of sugar entering the phloem [21,38,39]. In addition, decreasing the application of N fertilizer by 30% with NPI also increased the N uptake of plants by 11% and the nitrogen use efficiency compared with CI [25]. Meanwhile, increasing the K absorption by crops can promote nitrate metabolism, reduce nitrate contents, and increase the soluble sugar, soluble sugar and vitamin C contents in plants [15,29]. Furthermore, the K uptake levels increased by 48% and the K

utilization rates increased by 12% with NPI (-10 kPa value) compared with CI [40]. However, nutrient applications on the soil surface, especially N fertilizer, easily cause volatilization and leaching, which not only cause the loss of N but also decreases the quality of crops due to the uptake of excessive nitrate.

4.3. Fertilizer savings under NPI conditions

Planting crops decreases the fertilization dosages that are needed with NPI systems compared with CI, but excessive fertilizer applications did not significantly improve crop yields [3,9,25]. It was found that the NPI application rates of N, P, and K decreased by 30% compared with CI, but the yields were not significantly different in greenhouse production [23]. Meanwhile, the N dosage for lettuce under NPI was 20% lower than that under CI, and its yields still significantly increased by 14% [25]. The primary reason is that the fertilizer dosages that are applied to vegetables are generally high in China's greenhouses [41,42], which mainly rely on furrow and drip irrigation technologies, which are applied to the soil surface and easily cause the volatilization and leaching of N fertilizer. However, NPI can solve this shortcoming; the utilization efficiency of N fertilizer in cucumber increased by 52%, and the K utilization efficiency increased by 20%; simultaneously, the uptakes of NPK by tomato, lettuce and cucumber increased by 12% to 60% compared with CI [3,20,23].

In addition, using an appropriate fertilizer dosage improves crop yields and fertilizer use efficiencies under NPI conditions [9,23,25]. For example, for rapeseed irrigation with water and fertilization under NPI, the production was highest with a dose of 225 N kg ha⁻¹ compared with other N application treatments, and dosages of excessive amounts of fertilizer resulted in significant production declines [9]. Compared with the same N fertilizer dosage as CI, the N amounts can be reduced by 30% under NPI, but the yields did not decrease, the utilization of ammonium nitrogen can be improved, and the nitrate nitrogen levels can be reduced, which result in the mitigation of N₂O emissions [25]. Excessive applications of N fertilizer pollute rivers and groundwater and lead to high levels of greenhouse gas emissions [43,44].

This study found that NPI yields increased by 17% compared to a uniform dosage with CI, it provides guidance on fertilizer dosages for future research of NPI by using the meta-analysis, which would be helpful for fertilization management. The analysis of yield influencing factors also showed that phosphorus fertilizer and potassium fertilizer had greater influence on crop yield under negative pressure and conventional irrigation (Figure 5). Therefore, while considering the reductions in fertilizer applications under NPI, their impact on the environment can also be studied, especially the large-scale N fertilizer applications and mechanisms of soil and greenhouse gas emissions, which deserve further study.

At present, the research on NPI is mainly applied to vegetables, especially those with shallow root systems, such as rapeseed, spinach, and lettuce [9,24,25], but is less frequently applied to cereals and fruits, and there are also few experimental studies in the field [3]. In addition, it is necessary to use NPI technology in water-deficient areas to solve water shortage problems. Moreover, most of the water emitters used for NPI consist of ceramic heads, which are expensive and have a service life of only approximately 1 to 2 years, and the phenomenon of micropore blockage will occur, which is also one of the technologies that needs improvement [3,8]. In short, the combination of the water and fertilizer integration technologies of NPI can save water and nutrients and improve the yields and quality of crops. Negative pressure irrigation is an important strategy to solve the water resource shortage in China.

5. Conclusions

Overall, NPI significantly promoted crop yields, water use efficiencies, quality and nutrient uptakes when compared to conventional irrigation. Negative pressure values of



-2 ~ -5 kPa were the best indicator of crop yields, which were followed by values of -6 ~ -10 kPa and -12 ~ -15 kPa, which decreased yields, while the WUEs exhibited the highest performance at -6 ~ -10 kPa. The yield increases of leafy vegetables were significantly higher than those of fruit vegetables, while the WUEs were slightly higher for fruit vegetables. There was a dominance of sandy loam over clay soils and of alkaline soils over neutral soils under NPI conditions. The soil available phosphorus (P) content and application of P fertilizer have a greater impact on NPI and CI crop yields.

Author Contributions: Conceptualization, H.L. and S.Z.; methodology, Y.J., X.G., and S.Z.; software, Y.J.; validation, Y.J., X.G. and S.Z.; investigation, Y.J. and X.G.; resources, H.L., R. Z., X.W. and S.Z.; data curation, Y.J. and X.G.; writing—original draft preparation, Y.J., X.G. and S.Z.; writing—review and editing, Y.J., X.G. and S.Z.; supervision, H.L., X.W., Z.S. and S.Z.; project administration, R.Z. and H.L.; funding acquisition, H.L. and S.Z. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the National Key Research and Development Program of China (2018YFE0112300).

Acknowledgments: This research was funded by the National Key Research and Development Program of China (2018YFE0112300).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data sets generated and/or analyzed during the current study are available from the corresponding author on reasonable request.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Kadiresan, K., Khanal P. Rethinking irrigation for global food security. *Irrig. Drain.* **2018**, *67*, 8–11. <https://doi.org/10.1002/ird.2219>
- Peng, Y., Xiao, Y., Fu, Z., Dong, Y., Zheng, Y., Yan, H., Li, X. Precision irrigation perspectives on the sustainable water-saving of field crop production in China: Water demand prediction and irrigation scheme optimization. *J. Clean. Prod.* **2019**, *230*, 365–377. <https://doi.org/10.1016/j.jclepro.2019.04.347>
- Long, H., Wu, X., Zhang, S., Wang, J., Drohan, P. J., Zhang, R. Connotation and research progress of crop initiate water drawing technology *Trans. Chin. Soc. Agric. Eng.* **2020**, *36*, 139–152. (Chinese)
- Fan, Z., Lin, S., Zhang, X., Jiang, Z., Yang, K., Jian, D., Chen, Y., Li, J., Chen, Q., Wang, J. Conventional flooding irrigation causes an overuse of nitrogen fertilizer and low nitrogen use efficiency in intensively used solar greenhouse vegetable production. *Agric. Water Manage.* **2014**, *144*, 11–19. <https://doi.org/10.1016/j.agwat.2014.05.010>
- Long, H., Zhang, H., Yue, X., Zhang, R. Design and experiment of heavy-type negative pressure value used for negative pressure irrigation. *Trans. Chin. Soc. Agric. Eng.* **2018**, *34*, 85–92. (Chinese)
- Michael, O., David, O., Frederick, A., Paul, A., Kwame, A., Meta-analysis of crop yields of full, deficit, and partial root-zone drying irrigation. *Agric. Water Manage.* **2018**, *197*, 79–90. <https://doi.org/10.1016/j.agwat.2017.11.019>
- Lv, H., Lin, S., Wang, Y., Lian, X., Zhao, Y., Li, Y., Du, J., Wang, Z., Wang, J., Butterbach-Bahl, K. Drip fertigation significantly reduces nitrogen leaching in solar greenhouse vegetable productions system. *Environ. Pollut.* **2019**, *245*, 694–701. <https://doi.org/10.1016/j.envpol.2018.11.042>
- Yang, P., Bai, J., Yang, M., Ma, E., Yan, M., Long, H., Liu, J., L. L. Negative pressure irrigation for greenhouse crops in China: A review. *Agric. Water Manage.* **2022**, *164*, 107497. <https://doi.org/10.1016/j.agwat.2022.107497>
- Zhao, X., Gao, X., Zhang, S., Long, H. Improving the growth of rapeseed (*Brassica chinensis* L.) and the composition of rhizosphere bacterial communities through negative pressure irrigation. *Water Air Soil Poll.* **2019**, *239*, 9. <https://doi.org/10.1007/s11270-018-4061-1>
- Moniruzzaman, S., Fukuhara, T., Terasaki, H. Experimental study on water balance in a negative pressure difference irrigation system. *J. Japan Soc. Civ. Engin.* **2011**, *67*, 103–108. https://doi.org/10.2208/jscejhe.67.I_103
- Zhang, J., Ji, J., Wang, P., Long, H., Wu, X. Molecular mechanism of negative pressure irrigation inhibiting root growth and improving water use efficiency in maize. *Plant Soil*, **2021**, <https://doi.org/10.1007/s11104-021-05190-7>
- Zhang, J., Wang, P., Ji, J., Long, H., Wu, X., Transcriptome analysis reveals molecular mechanism of yield increases in maize under stable soil water supply. *PLoS One*, **2021**, <https://doi.org/10.1371/journal.pone.0257756>

13. Iwama, H., Kubota, T., Ushiroda, T., Osazawa, S., Katou, H. Control of soil water potential using negative pressure water circulation technique. *Soil Sci. Plant Nutr.* **1991**, *37*, 1, 7-14. <https://doi.org/10.1080/00380768.1991.10415004>
14. Bian, Y., Yang, P., Long, H., Ding, Y.H., Li, D. Water use efficiency and nutrient absorption of spinach (*Spinacia oleracea* L.) under two material emitters and negative water supply pressure. *J. Plant Nutr. Fert.* **2018**, *24*: 507-518. (In Chinese)
15. Gao, X., Zhang, S., Zhao, X., Long, H. Stable water and fertilizer supply by negative pressure irrigation improve tomato production and soil bacterial communities. *SN App. Sci.* **2019**, *1*, 178. <https://doi.org/10.1007/s42452-019-0719-6>
16. Li, Y., Xue, X., Guo, W., Wang, L., Duan, M., Chen, H., Chen, F. Soil moisture and nitrate-nitrogen dynamics and economic yield in the greenhouse cultivation of tomato and cucumber under negative pressure irrigation in the North China Plain. *Sci. Rep.* **2019**, *9*, 4439. <https://doi.org/10.1038/s41598-019-38695-4>
17. Song, Y., Zhao, X., Zhang, S. Reducing nitrogen input and improving yield and quality of rape through combination of fertigation and nitrification/urease inhibitor addition. *J. Plant Nut. Fertil.* **2017**, *23*: 632-640. (in Chinese)
18. Lipiec, J., Kubota, T., Iwama, H., Hirose, J. Measurement of plant water use under controlled soil moisture conditions by the negative pressure water circulation technique. *Soil Sci. Plant Nutr.* **1998**, *34*: 417-428. <https://doi.org/10.1080/00380768.1998.10415697>
19. Zhang, Z., Liu, S., J. S., Du, F., Qi, H., Li, J., Song X., Zhao, N., Nie, L., Fan, F. Precise soil water control using a negative pressure irrigation system to improve the water productivity of greenhouse watermelon. *Agric. Water Manage.* **2021**, *263*, 107144. <https://doi.org/10.1016/j.agwat.2021.107144>
20. Ding, Y., Xiao, H., Wang, D. Effect of potassium fertigation on potassium absorption and growth of flue-cured tobacco. *J. Plant Nutr. Fert.* **2017**, *23*: 1238-1248. (In Chinese)
21. Gao, X., Zhang, S., Zhao, X., Long, H. Evaluation of potassium application on tomato performance and rhizosphere bacterial communities under negative pressure irrigation of greenhouse-grown. *J. Plant Nutr.* **2020**, *43*, 317-326. <https://doi.org/10.1080/01904167.2019.1683862>
22. Wang, J., Huang, Y., Long, Y., Hou, S., Xiang, A., Sun, Z. Simulations of water movement and solute transport through different soil texture configurations under negative-pressure irrigation. *Hydrol Process.* **2017**, *31*: 2599-2612. <https://doi.org/10.1002/hyp.11209>
23. Li, S., Tan, D., Wu, X., Degre, A., Long, H., Zhang, S., Lu, J., Gao, L., Zheng, F., Liu, X., Liang, G. Negative pressure irrigation increases vegetable water productivity and nitrogen use efficiency by improving soil water and NO₃-N distributions. *Agric. Water Manage.* **2021**, *251*, 106853. <https://doi.org/10.1016/j.agwat.2021.106853>
24. Li, D., Long, H. Y., Zhang, S. X., Wu, X. P., Shao, H. Y., Wang, P. Effect of continuous negative pressure water supply on the growth, development and physiological mechanism of *Capsicum annuum* L. *J. Integr. Agric.* **2017**, *16*, 1978-1989. [https://doi.org/10.1016/S2095-3119\(16\)61572-1](https://doi.org/10.1016/S2095-3119(16)61572-1)
25. Gao, X., Zhang, S., Song, Y., Long, H. Negative pressure irrigation system reduces soil nitrogen loss for lettuce during greenhouse production. *Agronomy.* **2021**, *11*, 2380. <https://doi.org/10.3390/agronomy11122380>
26. Zhang, J., Wang, P., Long, H., Su, S., Wu, Y., Wang H. Metabolomics analysis reveals the physiological mechanism underlying growth restriction in maize roots under continuous negative pressure and stable water supply. *Agric. Water Manage.* **2022**, *263*, 107452. <https://doi.org/10.1016/j.agwat.2021.107452>
27. Schutz, L., Gattinger, A., Meier, M., Muller A., Boller, T., Mader P., Mathimaran, N. Improving crop yield and nutrient use efficiency via biofertilization – A global meta – analysis. *Front. Plant Sci.* **2018**, *12*. <https://doi.org/10.3389/fpls.2017.02204>
28. Xiang, Y., Huang, Y., Huang, C., Long, H. Influence of different water supply negative pressures on the growth and main physiological indexes of Chinese cabbage. *Journal of Agricultural Science and Technology*, **2019**, *20*: 16-22. (in Chinese)
29. Li, S., Wu, X., Dang, J., Pei, X., Gao, L., Li, J., Wang, B., Liang, G., Long H. Effects of negative pressure irrigation on yield, quality and water and nitrogen use efficiency of cucumber. *Soils Fertil. Sci China*, **2017**, *2* : 55-62. (in Chinese)
30. Wang, C., Gu, F., Chen, J., Yang, H., Jiang, J., Du, T., Zhang, J. Assessing the response of yield and comprehensive fruit quality of tomato grown in greenhouse to deficit irrigation and nitrogen application strategies. *Agric. Water Manage.* **2015**, *161*, 9-19. <https://doi.org/10.1016/j.agwat.2015.07.010>
31. Xiang, Y., Huang, Y., Long, H., Long, Q., Zhang, Y. Effects of different negative pressures on the growth and water use efficiency of pepper plants. *Res. Agric. Mod.* **2019**, *40*: 161-168. (in Chinese)
32. Wang, J., Huang, Y., Long, H. Water and salt movement in different soil textures under various negative irrigation pressures. *J. Integr. Agric.* **2016**, *15*, 1874-1882. [https://doi.org/10.1016/S2095-3119\(15\)61209-6](https://doi.org/10.1016/S2095-3119(15)61209-6)
33. Ma Y., Ren X., Yang S., Zong, J., Sun, H., Lv, M., Wu, D. Effect of fertigation by keeping irrigating water under negative pressure on quality and yield of pepper grown in greenhouse. *J. Irrig. Drain.* **2017**, *36*, 17-20.
34. Jiang, P., Lei, T., Bralts, V. Effects of soil textures and emitter material on the soil water movement and efficiency of negatively pressurized irrigation system. *Trans. Chin. Soc. Agric. Eng.* **2006**, *22*: 19-22. (in Chinese)
35. Wang, J., Long, H., Huang, Y., Wang, X., Cai, B., Liu, W. Effects of different irrigation management parameters on cumulative water supply under negative pressure irrigation. *Agric. Water Manage.* **2019**, *224*, 105743. <https://doi.org/10.1016/j.agwat.2019.105743>
36. Singandupe, B., Rao, G., Patil, G., Brahmanand, P. Fertigation studies and irrigation scheduling in drip irrigation system in tomato crop (*Lycopersicon esculentum* L.). *Eur. J. Agron.* **2003**, *19*, 327-340. [https://doi.org/10.1016/S1161-0301\(02\)00077-1](https://doi.org/10.1016/S1161-0301(02)00077-1)

37. Zhao, X., Song, Y., Yue, X. Effect of different potassium levels on the growth of bok choy under negative pressure. *Scient Agr. Sinica*. **2017**, *50*, 689–697. (in Chinese) 485
486
38. Adiku, K., Renger, M., Wessolek, G., Facklam, M., Hecht-Bucholtz, C. Simulation of the dry matter production and seed yield of common beans under varying soil water and salinity conditions. *Agric. Water Manage.* **2001**, *47*, 55–68. 487
[https://doi.org/10.1016/S0378-3774\(00\)00094-9](https://doi.org/10.1016/S0378-3774(00)00094-9) 488
489
39. Patane, C., Tringali, S., Sortino, O. Effects of soil water on yield and quality of processing tomato under a Mediterranean climate conditions. *Agric. Water Manage.* **2010**, *97*, 131–138. <https://doi.org/10.1016/j.agwat.2009.08.021> 490
491
40. Xiao, H., Ding, Y., Huang, C., Yang, H., Guan, E., Gao, K., Long, H. Effect of negative pressure irrigation on water fertilizer utilization and flue-cured tobacco growth. *Acta Tab. Sin.* **2016**, *22*, 52–60. (in Chinese) 492
493
41. Guo, Y., Chen, Y., Searchinger, T., Zhou, M., Pan, D., Yang, J., Wu, L., Cui, Z., Zhang, W., Zhang, F., Ma, L., Sun, Y., Zondlo, M., Zhang, L., Mauzerall, D. Air quality, nitrogen use efficiency and food security in China are improved by cost-effective agricultural nitrogen management. *Nat. Food*. **2020**, *1*, 648–658. <https://doi.org/10.1038/s43016-020-00162-z> 494
495
496
42. Wang, X., Dou, Z., Shi, X., Zou, C., Liu, D., Wang, Z. Innovative management programme reduces environmental impacts in Chinese vegetable production. *Nat Food*, **2021**, *2*, 47–53. <https://doi.org/10.1038/s43016-020-00199-0> 497
498
43. Subbarao, G. V., Searchinger, T. D. A “more ammonium solution” to mitigate nitrogen pollution and boost crop yields. *Proc. Natl. Acad. Sci. (USA)*. **2021**, *118*, e2107576118. <https://doi.org/10.1073/pnas.2107576118> 499
500
44. Subbarao, G. V., Kishii, M., Bozal-Leorri, A., Ortiz-Monasterio, I., Gao, X., Ibba, M. I., Karwat, H., Gonzalez-Moro, M. B., Gonzalez-Murua, C., Yoshihashi, T., Tobita, S., Kommerell, V., Braun, H., Iwanaga, M. Enlisting wild grass genes to combat nitrification in wheat farming: A nature-based solution. *Proc. Natl. Acad. Sci. (USA)*. **2021**, *118*, e2106595118. 501
<https://doi.org/10.1073/pnas.2106595118> 502
503
504