



Flavonoid baicalein suppresses oral biofilms and protects enamel hardness to combat dental caries

Hong Chen^{1,2,3,4#}, Sihong Xie^{1,2,3#}, Jin Gao^{1,2,3,4}, Liwen He^{1,2,3,4}, Wenping Luo^{1,2,3,4}, Michael D. Weir⁵, Thomas W. Oates⁵, Hockin H. K. Xu^{5,6,7}, Deqin Yang^{1,2,3,4*}

- ¹ Department of Endodontics, Stomatological Hospital of Chongqing Medical University, Chongqing, 404100, P.R. China
- ² Stomatological Hospital of Chongqing Medical University, Chongqing Key Laboratory of Oral Diseases and Biomedical Sciences, Chongqing 404100, P. R. China
- ³ Chongqing Municipal Key Laboratory of Oral Biomedical Engineering of Higher Education, Chongqing 404100, P. R. China
- ⁴ Chongqing Key Laboratory of Oral Diseases and Biomedical Sciences, 426 Songshi North Road, Yubei District, Chongqing 401147, P. R. China
- ⁵ Department of Advanced Oral Sciences and Therapeutics, University of Maryland Dental School, Baltimore, MD 21201, USA;
- ⁶ Center for Stem Cell Biology & Regenerative Medicine, University of Maryland School of Medicine, Baltimore, MD 21201, USA;
- ⁷ Marlene and Stewart Greenebaum Cancer Center, University of Maryland School of Medicine, Baltimore, MD 21201, USA

Co-first authors.

* Correspondence: yangdeqin@hospital.cqmu.edu.cn

Abstract: The objectives of this study were to investigate the effects of a novel method using flavonoids to inhibit *Streptococcus mutans* (*S. mutans*), *Candida albicans* (*C. albicans*) and dual-species biofilms, and to protect enamel hardness in a biofilm-based caries model for the first time. Several flavonoids, including baicalein, naringenin and catechin, were tested. Gold-standard chlorhexidine (CHX) and untreated (UC) groups served as controls. Optimal concentrations were determined by cytotoxicity assay. Biofilm MTT, colony-forming-units (CFUs), biofilm biomass, lactic acid and polysaccharide production were evaluated. Real-time-polymerase-chain reaction (qRT-PCR) was used to determine gene expressions in biofilms. Demineralization of human enamel was induced via *S. mutans*-*C. albicans* biofilms, and enamel hardness was measured. Compared to CHX and UC groups, the baicalein group achieved the greatest reduction in *S. mutans*, *C. albicans* and *S. mutans*-*C. albicans* biofilms, yielding the least metabolic activity, polysaccharide synthesis and lactic acid production ($p < 0.05$). The biofilm CFU was decreased in baicalein group by 5 logs, 4 logs, 5 logs, for *S. mutans*, *C. albicans* and *S. mutans*-*C. albicans* biofilms, respectively, compared to UC group. When tested in a *S. mutans*-*C. albicans* *in vitro* caries model, the baicalein group substantially reduced enamel demineralization under biofilms, yielding an enamel hardness that was 2.75 times greater than that of UC group. Hence, the novel baicalein method is promising to inhibit dental caries by reducing biofilm formation and protecting enamel hardness.

Citation: Lastname, F.; Lastname, F.; Lastname, F. Title. *Int. J. Mol. Sci.* **2022**, *23*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor: Firstname Lastname

Received: date

Accepted: date

Published: date



Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Short title: Flavonoid baicalein suppresses biofilms and protects enamel hardness

Keywords: Baicalein, *Streptococcus mutans*, *Candida albicans*, Biofilm inhibition, Dental caries.

1. Introduction

Dental caries is one of the most widespread and costly biofilm-mediated oral infectious diseases, affecting all ages of people worldwide [1]. Among the several hundred bacterial species in the dental plaque, *Streptococcus mutans* (*S. mutans*) is a principal causative agent of caries [2]. *Candida albicans* (*C. albicans*) is one of the most common fungi in

mouth [3, 4]. And oral colonization of *C. albicans* are increased in the root caries (RC) and early childhood caries (ECC) lesions [5, 6]. *C. albicans* has an extraordinary acidogenic capacity and acid tolerance, and its association with *S. mutans* results in increased exopolysaccharides (EPS) formation and acid production, yield enhancing biofilm cariogenicity [7, 8]. Indeed, the positive correlation between *C. albicans* and the risk of dental caries has been demonstrated [9], and individuals with higher yeast counts in saliva and dental biofilms have a higher incidence of RC and ECC [6, 7, 10].

Preventing biofilm formation is a key to avoid the occurrence of dental caries [11]. Chlorhexidine (CHX) is a potent dental agent against oral infections, due to its broad-spectrum antimicrobial efficacy [12, 13]. CHX can increase the cellular membrane permeability of bacteria or fungi by interacting with the anionic receptors on the cell surface [14, 15]. CHX is widely used in clinical dentistry, and its therapeutic benefit has been demonstrated in reducing dental plaque [16]. However, bacterial spores and mycobacteria are highly resistant to CHX [17]. Thus, novel and effective methods should be explored to fight and prevent cariogenic biofilms.

Flavonoids are a family of polyphenolic compounds possessing great antibacterial and antifungal action [18]. Studies have shown that flavonoids have potentially beneficial effects as antimicrobial agents in the therapy for human disease [19, 20]. In view of the fact that flavonoids represent an emerging threat of dental microbe, the present study was designed to evaluate the effect of selected single flavonoids (baicalein, naringenin, and catechin) for their activities against dental biofilm bacteria and fungi. In addition, to date, there has been no report on the role of flavonoids in affecting *S. mutans*, *C. albicans* and the dual-species biofilm formation. [This is not true. Here is at least one paper: https://doi.org/10.1016/j.micpath.2017.03.033](https://doi.org/10.1016/j.micpath.2017.03.033)

Therefore, the objectives of this study were to investigate, for the first time: (1) flavonoids could inhibit single (*S. mutans* or *C. albicans*) and dual-species (*S. mutans*-*C. albicans*) cariogenic biofilm formation, with CHX as control; (2) flavonoids would substantially reduce enamel demineralization and increase the enamel hardness under *S. mutans*-*C. albicans* biofilm. **The null hypothesis was that flavonoids would have no influence on biofilm formation of *S. mutans* and *C. albicans*.**

2. Results

The concentration of antimicrobial agent in this experiment was screened by cytotoxicity experiment. As shown in Fig. 1, the results of CCK-8 assay, the absorbance rates decreased with increasing concentrations based on negative values observed in negative control group. As the gold standard, the CHX were chosen as our positive control to evaluate the cell toxicity. Based on the cell cytotoxicity test, the antibacterial concentrations were selected for our study as follows: 0.25 mg/mL of baicalein, 1.00 mg/mL of naringenin, 0.25 mg/mL of catechin, and 0.000031% of CHX, respectively.

Why would you even start an experiment if you suspected it was the right thing to do in the first place?




Figure 1. Cell viability tests of HOK cells after being exposed to flavonoids (A) or CHX (B) (mean±sd; n=3).

Representative SEM micrographs of typical biofilms on adhesive disks in different groups are shown in Fig. 2. The five groups (CHX, baicalein, naringenin, catechin, and untreated control (UC)) were labeled in the images. The two types (*S. mutans* and dual-species) were labeled in these groups. Compared to the UC groups, biofilms of CHX, baicalein and naringenin groups on the adhesive disks did not fully develop even after 24 h. While the 0.250 mg/mL baicalein groups had the greatest antibacterial activity, only a few bacteria or fungi were observed on the surface of the disk.




Figure 2. SEM observation of (A) *S. mutans*, and (B) *S. mutans-C. albicans* biofilms after flavonoids or CHX treatment.

The CV staining image is presented in Fig. 3A, and further biofilm biomass results are shown in Fig. 3B (mean±sd; n = 6). The drugs greatly reduced the biomass significantly containing flavonoid or CHX, compared to the UC group ($p < 0.01$). And the inhibition effect of baicalein is the greatest, and much better than CHX ($p < 0.01$).

Six standard glucose concentrations of 0, 10, 20, 30, 40 and 50 µg/mL were used to plot the standard curve of OD_{620nm} versus polysaccharide concentrations. A linear curve $y = 0.0486x + 0.0614$ was obtained, and the coefficient of determination (R^2) was 0.9829. Fig. 3C presents the polysaccharide synthesis with or without drugs (mean±sd; n = 6). Two main results were obtained: (1). flavonoids and CHX remarkably decreased the polysaccharide production, compared to UC group (mean±sd; n = 6; $p < 0.01$). (2). Using baicalein, the polysaccharide synthesis decreased more than 90%, which is much better than CHX control ($p < 0.01$).

Fig. 3D plotted that *S. mutans* and dual-species biofilms had relatively higher levels of lactic acid production compared to *C. albicans* biofilms. The lactic acid production of single- and dual-species biofilms were greatly inhibited on the flavonoids and CHX groups ($p < 0.01$). And lactic acid production of baicalein on biofilms was relatively low.




Figure 3. Biofilm formation with or without drugs: (A) A CV staining photograph of biofilms grown in a 24-well polystyrene microtiter plate; (B) Quantification of biofilm biomass; (C) Lactic acid production; and (D) of the single-species and dual-species biofilms after flavonoids or CHX treatment. The values represent the mean±sd for six independent experiments (mean±sd, n= 6; *p < 0.05, **p < 0.01).

whether it is necessary to specify
p < 0.05 if you have it nowhere?

The antibacterial effects of flavonoids and CHX against the *S. mutans*, *C. albicans* and *S. mutans-C. albicans* biofilms are shown in Fig. 4 (mean±sd; n = 6). MTT metabolic activity and colony forming unit (CFU) were markedly decreased in baicalein ($p < 0.01$). Compared to UC group, the biofilm *S. mutans* CFU was decreased by 5 logs and 4 logs in *S. mutans* and *S. mutans-C. albicans* biofilms, at 0.250mg/mL baicalein, respectively (Fig. 4B). While 4 logs and 1 log in *C. albicans* and *S. mutans-C. albicans* biofilms was reduced for *C. albicans* CFU at 0.250mg/mL baicalein (Fig. 4C). And the usage of baicalein achieved the greatest reduction in biofilm activity, while the CHX group seemed to have no effect.

The sentence is not clear enough. Do you mean....

Compared with the UC group, the CFU of the *S. mutans* biofilm decreased by 5 logs and 4 logs in the *S. mutans* and *S. mutans-C. albicans* biofilms, respectively, at 0.250 mg/ml baicalein (Fig. 4B). While 4 logs and 1 log in *C. albicans* and *S. mutans-C. albicans* biofilms were reduced for *C. albicans* CFU at 0.250mg/mL baicalein (Fig. 4C). The use of baicalein caused the greatest reduction in biofilm activity, while the CHX group appeared to have no effect.

How come you get different results for the same sample group (*S. mutans*-*C. albicans*) (diagram B and C)?




Figure 4. Antimicrobial effects of flavonoids or CHX (mean±sd; n = 6; **p* < 0.05, ***p* < 0.01): (A) MTT metabolic activity, (B) colony-forming units (CFU) for *S. mutans*, and (C) colony-forming units (CFU) for *C. albicans*.

Please, add the name for second sample group

CLSM showed that only small clusters of bacterial cells were observed in the baicalein and naringenin groups, while the CHX and catechin groups were similar with UC group (Fig. 5A). It was concluded that baicalein and naringenin impaired EPS production and biofilm formation. Furthermore, we calculated the EPS/bacterial volume ratio, and baicalein exhibited the least value, which indicated the best anti-bacterial effect of baicalein in the EPS architecture development (Fig. 5B and Fig. 5C).




Figure 5. Representative CLSM staining images (A). And the biofilm thickness (B) and EPS/bacterial or fungal ratio was calculated (C) (mean±sd; n = 6; * $p < 0.05$, ** $p < 0.01$).

The expression profiles of *gtfB/C/D*, and *comC/D/E* genes are shown in Fig. 6. After the biofilm cultured with flavonoids or CHX, the transcription levels of *gtfB/C/D*, and *comC/D/E* genes statistically declined ($p < 0.05$). Compared with the UC group in the *S. mutans* biofilm, the expression of *gtfB/C/D* in the baicalein group was down-regulated to 0.58, 0.56 and 0.35 folds, respectively (Fig. 6A); while in the naringenin group, it was down-regulated to about 0.20 folds significantly. In dual-species biofilm, catechin down-regulate the expression of *gtfB/C/D* gene to 0.69-0.73 folds. In addition, the data revealed that the transcriptional levels of *comC/D/E* were significantly decreased in dual-species biofilm (Fig. 6B). The expression of *comC/D/E* in the baicalein group was down-regulated to 0.34-0.70 folds, while in the naringenin group, it was down-regulated to 0.25-0.40 folds (Fig. 6B).




Figure 6. The *S. mutans* *gtfB/C/D*, *comC/D/E* gene expression in single-species and dual-species biofilms via qRT-PCR (mean±sd, n = 3; **p* < 0.05, ***p* < 0.01).

Enamel hardness at the enamel surface is plotted in Fig. 7 (mean±sd; n = 6). The sound enamel hardness was 3.08 ± 0.21 GPa. After 14 days under the *S. mutans-C. albicans* biofilm acid attack, enamel hardness in every group all decreased, but the baicalein group had the highest enamel hardness (*p* < 0.01). These results showed that: (1) Flavonoids substantially increased the enamel hardness, **by three folds**, compared to the UC group; (2) The efficacy of baicalein and naringenin was much better than CHX in protecting the enamel hardness.

Is this correct?
Enamel hardness has
tripled?
Can you back this up
with exact numbers,
since it is not
apparent from the
chart?




Figure 7. Enamel hardness at the enamel surface after *S. mutans*-*C. albicans* biofilm acid attack for 14 days (mean±sd, n = 6; * $p < 0.05$, ** $p < 0.01$).

3. Discussion

Flavonoids are promising therapy against dental caries due to their antimicrobial activity and low toxicity. In the present study we investigated flavonoids with regard to their cytotoxicity on oral HOK cells and antimicrobial effects on cariogenic biofilm, using CHX as control group. There existed a certain usage concentration with regard to their cytotoxic activities *in vitro*. The antimicrobial ability observation and biofilm-based caries model proved that baicalein exerted instant plaque-inhibiting effects, helping control biofilm formation and protect tooth structure, which suggested that baicalein exerted great clinical application potentials.

Biocompatibility is a necessary indicators for evaluating all kinds of biomedical agents [21]. The toxicity to HOK cells was tested to explore the potential of flavonoids for dental clinical application [22]. An ideal antimicrobial agent should had the effectiveness of killing microbes meanwhile and the safety to host cell [21, 23]. So *in vitro* and *in vivo* potential toxicity of antimicrobial agents should be evaluated before clinical use. But sometimes *in vitro* studies used to evaluate drug toxicity tended to display cell toxicity, but they are still widely used in clinical treatment under certain conditions [21, 24]. For example, 0.2% CHX is widely used in mouthwash, but 0.2% CHX in the present study revealed obvious toxic effects on human HOK cells viability. In the present study, we also evaluated the cytotoxic activities of flavonoids. The results indicated that flavonoids concentration of 0.25mg/mL in baicalein, 1.00mg/mL in naringenin, and 0.25mg/mL in

The cited literature (25) does not correspond to this content.

catechin showed significant no-cytotoxic effects on HOK cells, according to previous study [25]. Therefore, the concentrations above were used in the following experiments.

C. albicans is a gram-positive fungal microorganism, which plays an important role in the development of dental caries because its positive interactions with *S. mutans* [8]. The interaction between *C. albicans* and *S. mutans* can promote the occurrence and development of root caries and ECC [5, 6]. *C. albicans* can enhance the sugar metabolism pathways of *S. mutans* [8]. *S. mutans* possesses an exceptional ability to produce EPS, and the EPS matrix is a key virulence factor for cariogenic biofilm [26, 27]. As nutrients and the core of the matrix scaffold, EPS could enhance biofilm accumulation and stability [26, 27]. In the present study, flavonoids disrupted the polysaccharide synthesis and biofilm formation in the *S. mutans*, *C. albicans*, and the dual-species biofilms. Therefore, the use of flavonoids might be a potential strategy for caries prevention, because previous reports have indicated that the EPS initiates the adhesion for first colonizers and is critical for dental biofilms development.

The biofilm CFU reduction in CHX had no significance compared to the UC group. In sharp contrast, the baicalein reduced the biofilm *S. mutans* CFU by nearly 5 logs and 4 logs in *S. mutans* and dual-species biofilms. While 4 logs and 1 log in *C. albicans* and dual-species biofilms were reduced for *C. albicans* CFU at 0.25mg/mL baicalein. Therefore, the anti-microbe effects of flavonoids are much better than CHX.

Acidogenicity in *S. mutans* is a virulence factor associated with cariogenicity [1, 28]. Biofilms can produce organic acids (mainly lactic acid), and induce to teeth demineralization [26, 28]. In the present study, the lactic acid secretion by biofilms decreased significantly in drugs, especially baicalein. The usage of baicalein achieved the greatest reduction in lactic acid production by more than 90% in *S. mutans* and dual-species biofilms. Such a major drop in biofilm acid production is expected to effectively inhibit dental caries, which warrants further study.

Sucrose-dependent adhesion is one of the major virulence factors [29, 30]. The adhesion process is mainly mediated by glucans, which are synthesized from sucrose by glucosyltransferase (GTF) enzymes [31]. GTFB, GTFC, and GTFD are the three important GTF enzymes [32]. The corresponding genes involved are *gtfB*, *gtfC*, and *gtfD*. The mRNA expression of *gtfB*, *gtfC* and *gtfD* of *S. mutans* and dual-species biofilm was examined using qRT-PCR in this study. The results showed that naringenin and baicalein could inhibit the mRNA expression for *gtfB/C/D* in *S. mutans* biofilm. The relative expression of *gtfB/D* for *S. mutans*-*C. albicans* biofilms culturing in baicalein, and *gtfB/C* in naringenin and *gtfB/C/D* in catechin was also decreased. It demonstrated that flavonoids could reduce the sucrose-dependent adhesion of *S. mutans* and further decrease single- and dual-biofilm formation, which may neutralize the increase of sucrose independent adhesion, finally showing an inhibition effect.

The QS system controls biofilm formation and release of virulence factors in cariogenic biofilms [33]. This system plays a key role in the competition and coexistence between microbe in biofilms and other significant process[34]. In *S. mutans*, the most common intraspecific QS system is the competence-stimulating peptide (CSP)-QS system

Almost the same sentence as in the results; repetition of saideog.

I think this is a bit "too strong" a conclusion based on a one type of analysis.

[35]. Hence, *comC/D/E* genes were tested and the results proved that these genes were downregulated after incubation with naringenin and baicalein in the present study. In other words, naringenin and baicalein could inhibit the synthesis of sensor and regulator of QS system, which would reduce the internal signal transduction in *S. mutans* or *S. mutans* and *C. albicans*.

In the *S. mutans*-*C. albicans* *in vitro* caries model, the protection effects of flavonoids were verified. The hardness of the enamel in drug groups was all higher than that of UC groups. Compared to the CHX and UC groups, the enamel hardness in baicalein groups showed a significant increase at the enamel surface under 14 days of biofilm acid attack. These results demonstrate that baicalein could inhibit the biofilm formation and hindered the cariogenic activities, effectively inhibiting tooth demineralization and protect tooth structure.

When the single- and dual-species biofilms were cultured in the flavonoids, the biofilm formation, metabolic activities and cariogenic activities were all significantly reduced. In addition, we proved the effect of flavonoids baicalein in inhibiting tooth demineralization under *S. mutans*-*C. albicans* biofilms for longer than 14 days. This study demonstrates the excellent antibacterial and antifungal effects of baicalein. Based on these results, the usage of baicalein could successfully inhibit dental caries pathogen biofilm formation. Further studies are needed to investigate the effect of baicalein *in vivo*. And more species biofilms are also need to study as to the complexity of oral microbe.

Regarding potential clinical applications, there is a major need in dentistry to decrease biofilm formation, as well as to inhibit caries and protect tooth structures. The present study demonstrated for the first time that the use of topical baicalein-containing solutions effectively suppressed caries associated *S. mutans* or *C. albicans*, yielding much greater enamel hardness under biofilm attacks. Dental applications of this novel method could include baicalein-based drugs for anti-caries clinical applications, such as dentifrices, mouthrinses, moisturizing gels, varnishes, and chewing gums. Further studies are needed to investigate and realize these potentials.

4. Materials and Methods

4.1. Chemicals, bacterial and fungal strains and growth conditions

Baicalein, naringenin, catechin and CHX were purchased commercially (solarbio, China). The use of all the bacterial species was approved by *** Institutional Review Board. *S. mutans* strain UA159 (ATCC 700610), and *C. albicans* SC5314 (ATCC MYA-2876) were provided by *** [3, 15]. Precultures of *S. mutans* were grown in brain-heart infusion (BHI) medium at 37°C under 5% CO₂ [11]. Precultures of *C. albicans* were grown in YPD medium containing 1% yeast extract, 2% peptone, and 2% D-glucose at 37°C under 5% CO₂ as well [3]. The *S. mutans* biofilm were cultured in BHI supplemented with 1% sucrose (wt/vol) (BHIS). As to *C. albicans* and dual-species biofilm, YNBB (0.67% YNB, 75 mM Na₂HPO₄-NaH₂PO₄, 2.5 mM N-acetylglucosamine, 0.2% casamino acids, and 0.5% sucrose) was used [3].

4.2. Biofilm formation

Precultures of *S. mutans* and *C. albicans* from single colonies were incubated overnight [3, 11]. Then *S. mutans* were diluted to 2×10^6 CFU/mL into fresh BHIS with or without drugs, and 2 mL bacteria dilution were cultured to form biofilms in 24-well plates. For dual-species biofilm, inoculum for the experiment was adjusted to 2×10^6 CFU/mL of *S. mutans*, and 2×10^4 cell/mL of *C. albicans* [3]. Equal volumes of each strain (200 μ L) and 1.6 mL YNBB medium with and without drugs were also incubated in 24-well plates for dual-species biofilm formation [36]. The plates were incubated at 37°C under 5% CO₂ for 24 h. The 24 h biofilms were then used for subsequent experiments.

4.3. Cytotoxicity assays

Human oral keratinocyte cells (HOK cells) were used for cytotoxicity assays [37]. The HOK cells were cultured in a low-glucose Dulbecco's modified eagle's medium (DMEM, Gibco, Grand Island, NY, USA) supplemented with 10% fetal bovine serum (FBS) and 1% penicillin-streptomycin (Invitrogen, Carlsbad, CA) (control media) [25, 37]. The cells were incubated at 37 °C with 5% CO₂ and the culture medium was changed at every 2-3 days [25]. Each group with or without drugs, and 8000 HOK cells were placed into a well of 96-well plates. Control medium was added accordingly. A cell counting kit (CCK-8, Dojindo, Tokyo, Japan) was used to evaluate cell proliferation [25]. Three replicates in each group were used for this assay (n = 3). After 24 h, the 96-well plate was washed twice with PBS. 100 μ L of growth medium with 10% CCK-8 for 2 hours added into each well, and the cell proliferative rate was determined via measuring the absorbance at OD_{450nm} using a microplate reader (SpectraMax M5, Molecular Devices) [25]. Optimal tested concentrations were the drug concentrations had cell viability of above 85% [21]. Then the drugs (flavonoids and CHX) and UC groups were used in the following experiments.

4.4. Scanning electron microscopy (SEM)

The biofilm inhibition effects were observed using a scanning electron microscopy (SEM, FEI, Hillsboro, OR, USA). Each well of the 24-well plate contained a round glass slide (diameter = 14 mm) with 2 mL bacteria or fungal dilution for biofilm formation [38]. After 24 h anaerobic incubation, the biofilms were fixed with glutaraldehyde at room temperature for 12 hours, then serially dehydrated in ethanol, tertiary butyl alcohol and sputter-coated with gold [38]. The specimens were examined at 500 $\times$ and 10,000 $\times$ magnification, and representative pictures are shown.

4.5. Crystal violet (CV) staining

The biomass of the biofilms with or without drugs was evaluated via CV staining, following previous studies [11, 39]. After incubation for 24 h, the plates were washed twice with PBS [11]. The biofilms were air-dried and then stained with 0.1% (w/v) CV for 5 min at room temperature, washed three times with PBS to remove the unbound stain, dried, and dissolved in 1 mL of 33% acetic acid [11]. The biofilm biomass was quantified by the optical density measured at a wavelength of 600nm (OD_{600nm}) [38]. The experiments were performed for six times in each group.

4.6. Polysaccharide synthesis

The phenol-sulfuric acid method was used to measure the water-insoluble polysaccharides of biofilms [15]. Biofilms with or without drugs were collected by scraping, and then they were centrifuged (12,000 rpm) for 5 min at 4°C and washed twice with PBS [15]. The precipitate was then resuspended in 1M NaOH solution, placed in a constant temperature water bath at 37°C, and incubated for 3 hours. 100 µL of 6% phenol solution and 0.5 mL of 95-97% sulfuric acid were added, followed by incubation for 30 min [11]. Then, 200 µL of the solution was transferred into a 96-well plate and OD_{620nm} was determined with the microplate reader (SpectraMax M5, Molecular Devices) [11, 39]. Six glucose concentrations of 0, 10, 20, 30, 40 and 50 µg/mL were used to plot the standard curve of OD_{620nm} readings to polysaccharide concentrations.

4.7. Lactic acid secretion

After incubation, the biofilms in plates were rinsed twice with PBS, and then immersed in 1.5 mL buffered peptone water (BPW, Sigma-Aldrich) with 0.2% sucrose and incubated at 37°C for 3 h (n = 6) [15, 39]. After removing planktonic cells by centrifugation, the supernatants were decanted to measure lactate concentrations according to the manuscript of the Lactate Assay Kit (MAK064) [40]. The absorbance at 530nm was recorded using a microplate spectrophotometer and lactate concentrations were calculated by the standard curves [15].

4.8. Biofilm viability using the MTT assay

The 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) (VWR Chemicals, Ohio, USA) assay was used to estimate the viability of bacteria and fungi in biofilms [15]. Biofilms with or without drugs were washed twice with PBS (n=6). 1mL MTT dye (0.5 mg/mL MTT in PBS) was added into each well and incubated at 37 °C under 5% CO₂ for 1 h. 1 mL dimethyl sulfoxide (DMSO) was added in each well at room temperature for 20 min to dissolve the formazan crystals. After mixing via pipetting, 200 µL of the DMSO solution was collected and transferred into 96-well plate. OD_{570nm} was determined using the microplate reader (SpectraMax M5, Molecular Devices, Sunnyvale, CA, USA).

4.9. Biofilm CFU counts

Six wells of each group in 24-well plates were used for CFU counting [15]. Biofilms with or without drugs were transferred into tubes with 2mL of PBS, and the biofilms were harvested by scraping and sonication/vortexing (Fisher, Pittsburg, PA, USA) [15]. The suspensions were serially diluted and spread onto BHI or YPD agar plates. For dual-species, we added 8 µg/mL amphotericin B in BHI agar plates and 8 µg/mL gentamicin in YPD agar plates to inhibit the growth of *C. albicans* and *S. mutans*, respectively [41]. After 2-day incubation at 37 °C under 5% CO₂, the colony number was counted and then CFU counts were determined.

4.10. Confocal laser scanning microscopy (CLSM)

As in previous...

For analyzing EPS production and distribution within the biofilms with or without drugs, we used CLSM for EPS/bacterial staining. As previous studies, 1 μ M Alexa Fluor 647 (Invitrogen, Eugene, OR, USA) and 2.5 μ M SYTO 9 (Invitrogen, Carlsbad, CA, USA) were used to label EPS and bacterial cells, respectively [38, 42]. Biofilms were grown on coverslips, and observed by CLSM (Olympus FV1000, Japan) at a range of 495–515 nm for SYTO 9 and 655–690 nm for Alexa Fluor 647 [38, 42]. Images of six random fields of each group were captured. A three-dimensional reconstruction of the biofilms was analyzed, and the EPS/bacteria ratio was calculated.

4.11. Quantitative real-time-polymerase-chain reaction (qRT-PCR)

Where is Table 1.?

qRT-PCR was used to quantify expression of selected genes in biofilms, with *gyrA* of *S. mutans* as an internal control [38]. The cells were harvested from the biofilms and snap frozen in liquid nitrogen until they were needed [15]. Total bacterial RNA isolation, purification and reverse transcription of complementary DNA (cDNA) were performed as previously described [15]. All primers for qRT-PCR were obtained commercially (Sangon Biotech, Shanghai, China) and are listed in Table 1. Threshold cycle values (CT) were determined, and the data were analyzed by BIO-RAD CFX MANAGER software (version 2.0) using the $2^{-\Delta\Delta CT}$ method [38].

4.12. Enamel hardness measurement

Extracted caries-free human teeth were obtained from the *** following a protocol approved by the *** Institutional Review Board. Teeth were cleaned and stored in 0.1% thymol solution at 4 °C before use. As previous study, enamel slabs were prepared to have a diameter of 6 mm and a thickness of 2 mm [39]. Except the enamel surfaces, the rest area was coated with two layers of acid-resistant nail varnish. The specimens were polished using sandpapers with grit of # 600, 1200, 2400 and 4000, consecutively, with copious water [39]. The 30 enamel slabs were randomly divided into five groups of 6 slabs each and incubated with or without drugs.

A hardness tester (HVS-10Z, Jingbo Company, Zhejiang, China) was used with a Vickers indenter, under a 50 g load with a dwell time of 20 s [39]. The area selected for indentation was the center of enamel surface. Every enamel specimen had six indentations. Measurements were conducted before and after biofilm acid attacks. The sterile specimens were placed in 24-well plates containing 2 mL of *S. mutans* and *C. albicans* suspension, as previously described [39]. Each day, the slabs were placed into a well of 24-well plates, and immersed in 2 mL of YNBB at pH 7.4 at 37 °C for 4 hours. Then it was removed and placed into a new plate with the YNBB-R medium which reduced the sucrose content at pH 7 for 20 hours at 37 °C in 5% CO₂. The daily medium change was done under aseptic conditions. Since this biofilm model took 14 days to finish, to avoid the biofilm becoming old and too thick which would deprive the interior bacteria from nutrients, a sterile paper was used to remove the biofilm from the slab every 24 h, which was then inoculated again

to grow a new biofilm on the slab. This biofilm model and cyclic immersion treatment was repeated for 14 days.

4.13. Statistical analysis

Data analyses were performed using Statistical Package for the Social Sciences (SPSS 22.0, Chicago, IL, USA). All data were expressed as the mean value $\pm$ standard deviation (mean $\pm$ sd). One-way analysis of variance (ANOVA) was performed and Tukey's multiple comparison test was used to compare the means of each of the groups. The different level of p was set at 0.05.

? Please, explain this.
This is not correct neither used in this experiment.

5. Conclusions

This study demonstrated for the first time that flavonoid baicalein had strong biofilm-suppression, caries-inhibition and enamel-protection capabilities. Baicalein showed the most potent and the greatest reduction in biofilm biomass, polysaccharide and lactic acid production among the test groups of baicalein, naringenin and catechin, and CHX. Baicalein reduced biofilm CFU by more than 4 orders of magnitude, and reduced biofilm acids by 99.99%, for *S. mutans*-*C. albicans* biofilm, compared to UC group. In addition, the enamel hardness in baicalein group showed only a minimal decrease under 14-day biofilm acid attacks, resulting in an enamel hardness that was 2.75 times greater than that of the untreated control group. Baicalein-based drugs are promising for a wide range of applications to suppress biofilms, prevent dental caries and protect tooth structures.

Author Contributions: Author Contributions: Conceptualization, J.G., W.L., H.C., L.H. and H.H.K.X.; Methodology, H.C., D.Y., S.X., L.H., H.H.K.X., and M.D.W.; Investigation, S.X., H.C., W.L., and D.Y.; Analysis and Graphs— H.C., S.X., and T.W.O.; Writing—H.C., S.X., and D.Y.; Writing—Review & Editing, H.H.K.X., H.C., D.Y. and J.G.; Funding Acquisition, D.Y. and H.C.; Resources, S.X., J.G., H.C.; M.D.W. and T.W.O.; Supervision, M.D.W., W.L., and L.H.; Project Administration, H.H.K.X., D.Y. and H.C.. All authors have read and agreed to the published version of the manuscript.

Funding: This study was supported by National Natural Science Foundation of China (No.31970783 to D. Y., No. 82100991 to H.C.); Program for Top talent Distinguished Professor from Chongqing Medical University to D.Y., Program for Youth Innovation in Future Medicine from Chongqing Medical University (No. W0060 to D.Y.); and Chongqing Special Postdoctoral Science Foundation (No. 2010010005994583 to H.C.).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Pitts NB, Zero DT, Marsh PD et al. Dental caries. *Nat Rev Dis Primers* 2017; **3**: 1-16.
2. Ajdić D, McShan WM, McLaughlin RE et al. Genome sequence of *Streptococcus mutans* UA159, a cariogenic dental pathogen. *Proc Natl Acad Sci USA* 2002; **99**: 14434-9.
3. Du Q, Ren B, He J et al. *Candida albicans* promotes tooth decay by inducing oral microbial dysbiosis. *ISME J* 2021; **15**: 894-908.
4. Daniluk T, Tokajuk G, Stokowska W et al. Occurrence rate of oral *Candida albicans* in denture wearer patients. *Adv Med Sci* 2006; **51**: 77-80.
5. Takahashi N, Nyvad B. Ecological hypothesis of dentin and root caries. *Caries Res* 2016; **50**: 422-31.

6. Fakhruddin KS, Perera Samaranayake L, Egusa H et al. Candida biome of severe early childhood caries (S-ECC) and its cariogenic virulence traits. *J Oral Microbiol* 2020; **12**: 1724484. 437 438
7. Shen S, Samaranayake L, Yip H, Dyson J. Bacterial and yeast flora of root surface caries in elderly, ethnic Chinese. *Oral Dis* 2002; **8**: 207-17. 439 440
8. He J, Kim D, Zhou X et al. RNA-seq reveals enhanced sugar metabolism in *Streptococcus mutans* co-cultured with *Candida albicans* within mixed-species biofilms. *Front Microbiol* 2017; **8**: 1036. 441 442
9. Pereira D, Seneviratne C, Koga-Ito C, Samaranayake L. Is the oral fungal pathogen *Candida albicans* a cariogen? *Oral Dis* 2018; **24**: 518-26. 443 444
10. Zaremba M, Stokowska W, Klimiuk A et al. Microorganisms in root carious lesions in adults. *Adv Med Sci* 2006; **51**: 237-40. 445
11. Chen H, Tang Y, Weir MD et al. Effects of *S. mutans* gene-modification and antibacterial monomer dimethylaminohexadecyl methacrylate on biofilm growth and acid production. *Dent Mater* 2020; **36**: 296-309. 446 447
12. Varoni E, Tarce M, Lodi G, Carrassi A. Chlorhexidine (CHX) in dentistry: state of the art. *Minerva Stomatol* 2012; **61**: 399-419. 448
13. Seneviratne CJ, Leung KC-F, Wong C-H et al. Nanoparticle-encapsulated chlorhexidine against oral bacterial biofilms. *Plos One* 2014; **9**: e103234. 449 450
14. Kumar SB. Chlorhexidine mouthwash-a review. *J Pharm Pharm Sci* 2017; **9**: 1450. 451
15. Chen H, Zhang B, Weir MD et al. *S. mutans* gene-modification and antibacterial resin composite as dual strategy to suppress biofilm acid production and inhibit caries. *J Dent* 2020; **93**: 103278. 452 453
16. Sreenivasan P, Gaffar A. Antiplaque biocides and bacterial resistance: a review. *J Clin Periodontol* 2002; **29**: 965-74. 454
17. Mohammadi Z, Abbott PV. Antimicrobial substantivity of root canal irrigants and medicaments: a review. *Aust Endod J* 2009; **35**: 131-9. 455 456
18. Gutiérrez-Venegas G, González-Rosas Z. Apigenin reduce lipoteichoic acid-induced inflammatory response in rat cardiomyoblast cells. *Arch Pharm Res* 2017; **40**: 240-9. 457 458
19. Serpa R, França EJ, Furlaneto-Maia L et al. In vitro antifungal activity of the flavonoid baicalein against *Candida* species. *J Med Microbiol* 2012; **61**: 1704-8. 459 460
20. Gutiérrez-Venegas G, Gómez-Mora JA, Meraz-Rodríguez MA et al. Effect of flavonoids on antimicrobial activity of microorganisms present in dental plaque. *Heliyon* 2019; **5**: e03013. 461 462
21. Zhou Y, Wang S, Zhou X et al. Short-time antibacterial effects of dimethylaminododecyl methacrylate on oral multispecies biofilm in vitro. *Biomed Res Int* 2019; **2019**. 463 464
22. Weng L, Wu L, Guo R et al. Lactobacillus cell envelope-coated nanoparticles for antibiotic delivery against cariogenic biofilm and dental caries. *J Nanobiotechnol* 2022; **20**: 1-20. 465 466
23. Müller H-D, Eick S, Moritz A et al. Cytotoxicity and antimicrobial activity of oral rinses in vitro. *Biomed Res Int* 2017; **2017**. 467
24. Wyganowska-Swiatkowska M, Kotwicka M, Urbaniak P et al. Clinical implications of the growth-suppressive effects of chlorhexidine at low and high concentrations on human gingival fibroblasts and changes in morphology. *Int J Mol Med* 2016; **37**: 1594-600. 468 469 470
25. Chen H, Yang H, Weir MD et al. An antibacterial and injectable calcium phosphate scaffold delivering human periodontal ligament stem cells for bone tissue engineering. *RSC Adv* 2020; **10**: 40157-70. 471 472
26. Koo H, Falsetta M, Klein M. The exopolysaccharide matrix: a virulence determinant of cariogenic biofilm. *J Dent Res* 2013; **92**: 1065-73. 473 474
27. Koo H, Xiao J, Klein MI. Extracellular polysaccharides matrix—an often forgotten virulence factor in oral biofilm research. *Int J Oral Sci* 2009; **1**: 229. 475 476
28. Law V, Seow W, Townsend G. Factors influencing oral colonization of mutans streptococci in young children. *Aust Dent J* 2007; **52**: 93-100. 477 478

-
29. Rosenberg M. Microbial adhesion to hydrocarbons: twenty-five years of doing MATH. *FEMS Microbiol Lett* 2006; **262**: 129-34. 479-480
30. Strużycka I. The oral microbiome in dental caries. *Pol J Microbiol* 2014; **63**: 127. 481
31. Yue J, Yang H, Liu S et al. Influence of naringenin on the biofilm formation of *Streptococcus mutans*. *J Dent* 2018; **76**: 24-31. 482
32. Krzyściak W, Jurczak A, Kościelniak D et al. The virulence of *Streptococcus mutans* and the ability to form biofilms. *Eur J Clin Microbiol* 2014; **33**: 499-515. 483-484
33. Worthington RJ, Richards JJ, Melander C. Small molecule control of bacterial biofilms. *Org Biomol Chem* 2012; **10**: 7457-74. 485
34. Ding Y, Wang W, Fan M et al. Antimicrobial and anti-biofilm effect of Bac8c on major bacteria associated with dental caries and *Streptococcus mutans* biofilms. *Peptides* 2014; **52**: 61-7. 486-487
35. Suntharalingam P, Cvitkovitch DG. Quorum sensing in streptococcal biofilm formation. *Trends Microbiol* 2005; **13**: 3-6. 488
36. Liu S, Qiu W, Zhang K et al. Nicotine enhances interspecies relationship between *Streptococcus mutans* and *Candida albicans*. *Biomed Res Int* 2017; **2017**. 489-490
37. Nagai J, Shi H, Kubota Y et al. Quantitative structure–cytotoxicity relationship of pyrano [4, 3-b] chromones. *Anticancer Research* 2018; **38**: 4449-57. 491-492
38. Mao M-Y, Yang Y-M, Li K-Z et al. The *mnc* gene promotes exopolysaccharide synthesis and represses the *vicRKX* gene expressions via microRNA-size small RNAs in *Streptococcus mutans*. *Front Microbiol* 2016; **7**: 687. 493-494
39. Chen H, Tang Y, Weir MD et al. Effects of *S. mutans* gene-modification and antibacterial calcium phosphate nanocomposite on secondary caries and marginal enamel hardness. *RSC Adv* 2019; **9**: 41672-83. 495-496
40. Wang Y, Wang X, Jiang W et al. Antimicrobial peptide GH12 suppresses cariogenic virulence factors of *Streptococcus mutans*. *J Oral Microbiol* 2018; **10**: 1442089. 497-498
41. Escobar IE, Rossatto FCP, Kim SM et al. Repurposing Kinase Inhibitor Bay 11-7085 to Combat *Staphylococcus aureus* and *Candida albicans* Biofilms. *Front Pharmacol* 2021; **12**. 499-500
42. Li Z, Xiang Z, Zeng J et al. A GntR family transcription factor in *Streptococcus mutans* regulates biofilm formation and expression of multiple sugar transporter genes. *Front Microbiol* 2019; **9**: 3224. 501-502-503