

Why Do Bio-carbonates Exist?

Luis Pomar¹, Pamela Hallock^{2,5}, Guillem Mateu-Vicens³, and Juan I. Baceta⁴

¹ Catedra Guillem Colom, Universitat de les Illes Balears, E-07122 Palma de Mallorca, Spain

² College of Marine Science, University of South Florida, St Petersburg, FL 33701, USA

³ Department of Biology, Universitat de les Illes Balears, E-07122 Palma de Mallorca, Spain

⁴ Department of Geology, Universidad del Pais Vasco, E-48940, Leioa, Basque Country, Spain

* Correspondence: pmuller@usf.edu

Abstract: Calcium carbonate precipitation associated with biotic activity is first recorded in Archaean rocks. The oldest putative fossils related to hydrothermal vents have been dated at ~3.77 Ga (possibly 4.29 Ga). Stromatolites, the oldest dated at 3.70 Ga, have since occurred through Earth history, despite dramatic changes in physical and chemical conditions in aquatic environments. A key question is: What advantages do photosynthesizing aquatic prokaryotes and algae gain by precipitating carbonates? We propose the “Phosphate Extraction Mechanism” (PEM) to explain the benefits of biomineralization in warm, oligotrophic, alkaline, euphotic environments. Carbonate precipitation enhances access to otherwise limited carbon dioxide and phosphate in such environments. This mechanism also provides an explanation for prolific production of carbonates during times of elevated atmospheric carbon dioxide at intervals in the Phanerozoic.

Keywords: Phosphate; nutrient limitation; carbon dioxide; Archaean; Proterozoic; cyanobacteria; calcareous algae; coccolithophores

1. Introduction

Carbon, which is the major component of fossil fuels, is the key ingredient in “life”. Carbon is also a key ingredient in limestones and other carbonate rocks. The ultimate sources of carbon are the primordial components of the Earth. Even at present, volcanic sources emit ~1–4x10⁸ metric tons of carbon dioxide (CO₂) per year. Through geologic history, the dynamics of carbon chemistry have played major roles in oceanic, terrestrial and even subsurface crustal processes. Humans are now burning the equivalent of several hundred years of photosynthetic sequestration of carbon each year, releasing CO₂ into the Earth’s atmosphere, with climatic consequences that are becoming more apparent every year.

The systematic measure of CO₂ concentrations in the atmosphere was started by C. David Keeling of the Scripps Institution of Oceanography in March 1958 at a NOAA (National Oceanic and Atmospheric Administration) facility at Mauna Loa Observatory, Hawaii, USA [1]. NOAA initiated its own CO₂ measurement in May 1974 and, since then, NOAA and Scripps have run the measurements in parallel [2]. Prior to Keeling’s research, CO₂ measurements were inconsistent, but the Keeling’s methods revealed a pattern of seasonal oscillations of CO₂, with peaks in May and lows in November, with the averaged values of successive years progressively increasing (Fig. 1). Keeling envisaged the yearly cycles to reflect the vegetation cycles that prevail across the northern hemisphere while the increase over time was thought to be caused by human activities, especially the burning of fossil-fuels [3].

Citation: Lastname, F.; Lastname, F.; Lastname, F. Title. *J. Mar. Sci. Eng.* **2022**, *10*, x.
<https://doi.org/10.3390/xxxxx>

Academic Editor: Firstname Lastname

Received: date

Accepted: date

Published: date

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).




Figure 1. CO₂ concentrations in the atmosphere, started by C. David Keeling of the Scripps Institution of Oceanography in March 1958.

By the 1960's, greenhouse gas emissions and their link to global climate change became a serious concern, with both scientists and the public considering CO₂ to be an invisible pollutant. Attention and concern for greenhouse gases and their role in climate change intensified, and in 1988 the United Nations created the Intergovernmental Panel on Climate Change (IPCC) [4]. Concern for ocean acidification, *climate change's equally evil twin*, emerged somewhat later. Although scientists had been tracking ocean pH for more than 30 years, biological studies emerged in the 1990s [5] and have accelerated since the introduction of the term "ocean acidification" in 2003 [6].

Burning of fossil fuels, combined with widespread changes in land use, has resulted in rapidly increasing concentrations of atmospheric CO₂ that are causing the decline in the pH of surface seawater [6–9]. Will progressive warming and acidification cause mass extinctions and evolutionary turnover of marine biotas, as occurred at previous events that caused major perturbations of ocean chemistry [10]?

This question reveals a major carbonate paradox in the geologic record. Through the Phanerozoic, extended Greenhouse World times of higher CO₂ (Silurian–Devonian and Jurassic–Cretaceous) were characterized by prolific accumulations of biogenic carbonates, especially those associated with cyanobacterial and algal calcification (Fig. 2). In the Cenozoic, coral reefs emerged as major carbonate factories as Icehouse World conditions developed in the Oligocene and became predominant. In the Neogene, a conundrum noted more than 45 years ago [25].




Figure 2. Carbonate precipitation nodes, genetic macroevolution rates, and estimates of oxygen and CO₂ in the atmosphere. **A:** Supercontinents and crustal growth; precratonic, accretionary orogens and supercontinents assembly (redrawn from [11]). **B:** carbonate precipitation modes through Earth's history (multiple sources). [1]: Oldest fossils [11]. [2]: Isua oldest stromatolites [13]. **C:** pH evolution from [14]; **a:** nominal model, in which the median Archean surface temperature is slightly higher than modern surface temperatures. Archean land fraction was anywhere between 10% and 75% of modern land fraction; **b:** no Archean land endmember scenario; **c:** model with assumed 100 ppm Proterozoic methane and 1% Archean methane levels. **D:** Maximum and minimum estimates of atmospheric partial oxygen pressure (after [15]). **E:** estimates of atmospheric CO₂ concentration; [3] from [16], upper and lower boundaries reflect average surface temperature for an ice-free (20°C) and ice covered (5°) Earth; [4] Acritarch isotopic composition [17]; [5] C-isotope reservoir modeling

[18]; [6] Phanerozoic GEOCARB III models [19]; [7] Royer compilation [20]; [8] from [21]; [9] paleo-soil mass balances [22]; [10] from [23]; [11] Picoplanktonic whitening and partial sheath calcification commenced ~1400-1300 Ma ago (33 CO₂ PAL, and cyanobacteria CCMs were induced when pCO₂ = 10 PAL [24]. PAL: present atmospheric level; GOE: the Great Oxygenation Event; NOE: Neoproterozoic oxygenation event.

Another paradox associated with changes in ocean chemistry over the course of Earth history relates to another essential element for life, yet is commonly the most limiting nutrient required for essential processes including photosynthesis, growth and reproduction. Bioavailable phosphorus (P), typically occurring as phosphate (PO₄³⁻), provides the backbone of nucleotides (DNA and RNA) and phospholipid membranes of cells, and is used in many other essential functions including ATP-energy storage and energy transfer, and for the synthesis of proteins and enzymes. Kempe and Degens [26] postulated that abundant dissolved PO₄³⁻ in the Archaean seas helped to foster the evolution of life; see also [27, 28]. The goal of our paper is to explore the hypothesis that decline in both CO₂ and PO₄³⁻, and the history of changes in modes of calcification through Earth history, are tied to the close and complex relationships between biogenic carbonates and phosphate.

2. The Hypothesis

The “Phosphate Extraction Mechanism” (PEM) provides an explanation for how calcifying microbial and algal biotas can thrive in oligotrophic conditions. During daylight, calcification can be coupled to photosynthesis. With energy from active ion transport, protons can be split from bicarbonate, providing a carbonate ion (CO₃²⁻) for calcification and a CO₂ molecule for photosynthesis. Phosphate is adsorbed during calcium carbonate (CaCO₃) precipitation. At night, respiration releases CO₂ and can promote partial dissolution of diurnally precipitated carbonates. Phosphate that is adsorbed onto precipitating CaCO₃ during daylight can be desorbed at night, making it available for uptake by the calcifying cyanobacteria or algae.

3. The Hadean–Archaean

Conditions on the Hadean Earth were clearly very different from present, and even from those of the Proterozoic. One hypothesis is that the early Earth was hot, following the moon-forming impact, and cooled to the point where liquid water was present after about 10 million years [29,30]. Whether the Earth's surface continued to be thermophilic well into the Archaean is still debated, though ~4.3 Ga rocks near Hudson Bay are suspected to have formed under warm greenhouse conditions [31].

Over the past 4.5 Ga, heat-, gravity-, and tectonically-driven processes have concentrated Iron (Fe) and Nickel (Ni) in the Earth's core. The dominant elements in the Earth's crust are Oxygen (O), Silicon (Si), Aluminum (Al), Iron (Fe), Calcium (Ca), Magnesium (Mg), Sodium (Na), and Potassium (K). In the mantle, Mg is nearly an order of magnitude more prevalent than in the crust, where proportions especially of Si and Al are higher. Basaltic, komatiitic (ultramafic) and mantle ejecta are potent CO₂ sinks (all are rich in Mg²⁺ and Fe²⁺, with somewhat less Ca²⁺) and their subduction may have drawn down atmospheric CO₂ [30,32]. Alkaline rocks in India indicate carbonate subduction occurred by 4.26 Ga [30].

Through the Archaean, the combination of high atmospheric CO₂ concentrations and the prevalence of Mg²⁺, Fe²⁺, and Ca²⁺ resulted in formation of massive abiogenic dolomites and limestones, indicating extreme oversaturation of waters (Fig. 2B). The earliest hints of life date back ~3.8 billion years (early–mid Archaean, Fig. 2A, B), and by ~3.7 Ga, some metacarbonate rocks, found in the Isua supercrustal belt in southwest Greenland, contain 1–4-cm-high isolated and aggregated stromatolites [33]. Through the Proterozoic, massive abiogenic carbonates declined in prevalence relative to biogenic carbonates and are unknown in modern marine environments, even in those that are strongly oversaturated with respect to CaCO₃. In a water body containing abundant Ca²⁺ and dissolved inorganic

carbon, modest increases in saturation, whether by warming, reduced hydrostatic pressure, evaporation, or by biogenic processes such as photosynthetic uptake of CO_2 or microbial sulfate reduction, can trigger CaCO_3 or $\text{CaMg}(\text{CO}_3)_2$ precipitation.

So the questions we pose are these: In post Archaean oceans, did biogenic processes become so ubiquitous that they became the drivers and controllers of CaCO_3 precipitation, resulting in the overwhelming predominance of biogenic carbonate production? Was the sequence of carbonate factories through geologic history, from predominantly abiogenic precipitation in the Archaean, to predominance of biologically-induced geochemical precipitation in the Proterozoic, to the addition of biologically-controlled calcification during the Phanerozoic (Fig. 2B), consistent with and driven by evolutionary processes involving luminosity of the Sun, the stratification of the Earth, and the emergence of life, with their combined effects on ocean chemistry (Fig. 2C)?

The luminosity of the Sun has increased ~30% during the past 4.5 Ga [34]. The composition of the Earth's atmosphere has changed (Fig. 2D,E), with substantial decline in greenhouse gases, including CO_2 (Fig. 2E) and methane (CH_4). For example, CO_2 has been estimated to have been ~100 times higher in the Hadean than modern levels and ~10 times higher in the Cambrian than present [30]. As noted above, ultramafic ejecta are potent CO_2 sinks and their subduction may have drawn down atmospheric CO_2 in the Hadean [32].

4. Pertinent Chemical Processes

Before delving into biological processes, we briefly examine some chemical factors associated with CO_2 in aqueous environments (Fig. 3). Two measures associated with but not restricted to CO_2 are pH and alkalinity. Both are important in water chemistry, but their interactions can be confusing when considering calcification. The concentration of hydrogen ions (H^+), indicating how acidic or basic a substance is, is measured as pH. Alkalinity indicates the ability of a solution to neutralize acids (i.e., buffering). Alkalinity consists of ions that can incorporate H^+ (protons) into their structure, limiting the availability of those protons that would otherwise lower pH. Carbonate alkalinity measures the concentrations of CO_3^{2-} and HCO_3^- ions, which are typically present in the highest concentrations in natural waters. Total Alkalinity reflects primarily CO_3^{2-} and HCO_3^- ions, but also includes PO_4^{3-} , borate, orthosilicate, sulfides, and organic acids. Shallow water bodies can be strongly influenced by rising temperature and evaporation, concentrating the alkaline ions, and promoting the precipitation of carbonate minerals, assuming that appropriate cations are available (e.g., Ca^{2+} , Mg^{2+}). Moreover, any process that takes up protons can also promote such precipitation.




Figure 3. In an aqueous solution, carbonate, bicarbonate, carbon dioxide, and carbonic acid exist together in a dynamic equilibrium. When CO_2 is absorbed by seawater, a series of chemical reactions increase the concentration of H^+ and cause the seawater acidity to increase. When calcium carbonate (e.g., limestone) reacts with acidic free hydrogen (H^+) ions in seawater, the solid calcium carbonate dissolves, releasing free calcium (Ca^{2+}) ions and free bicarbonate (HCO_3^-) ions.

As elements that are relatively abundant in both crustal and mantle rocks, Mg^{2+} , Fe^{2+} , and Ca^{2+} ions are soluble in aquatic environments. Thus, the combination of high carbonate alkalinity and abundant reactive cations, especially in warm waters, provided prolific sources of ions to produce abiogenic carbonates, providing critical storage of atmospheric CO_2 on geological time scales [35].

5. Pertinent Biological Processes

All living organisms carry traces of the histories of their ancestors within their genetic makeup and, in the case of the Eukarya, literally within their cells. The diversity of metabolic pathways in the prokaryotic Archaea and Eubacteria far surpasses the comparatively limited pathways found in the Eukarya, which are metabolically limited to those of their Proteoarchaeota and α -proteobacterial symbiotic predecessors [36].

Early microbial forms evolved in aquatic environments lacking free oxygen (O_2). Autotrophic processes in aquatic systems require: a) an energy source, b) source of dissolved inorganic carbon (DIC), c) a source of protons, and d) nutrients (fixed N, P, Fe, and trace elements) necessary for cell growth and reproduction [37]. Energy sources include oxidation of inorganic compounds (e.g., H_2S , CH_4) by chemoautotrophs and sunlight by photoautotrophs [38]. The primary sources of DIC are CO_2 or CH_4 . In photoautotrophy, proton donors can be H_2 , Fe^{2+} or H_2S , which do not release O_2 (anoxygenic photosynthesis), or H_2O that releases O_2 as a byproduct (oxygenic photosynthesis).

RuBisCO is the enzyme that catalyzes the first major step of carbon fixation in the Calvin cycle, the process of photosynthesis. As the most abundant protein on Earth, RuBisCO is found in all three domains of life (Eubacteria, Archaea and Eukarya), and fixes more than 90% of the inorganic carbon that is converted into biomass [39]. However, this enzyme evolved in an O_2 -free atmosphere, prior to the Great Oxygenation Event, and does not efficiently discriminate between CO_2 and O_2 . Enzyme activity and specificity are reciprocally linked: faster RuBisCO has a higher error rate and a more specific RuBisCO has a lower catalytic rate.

6. CCMs: CO_2 Concentrating Mechanisms

It may seem paradoxical that RuBisCO, this ubiquitous and essential enzyme, has not become more efficient or been replaced by more efficient enzymes. However, CCMs (CO_2 concentrating mechanisms) evolved instead by adaptation of active transport processes and compartmentalization of accumulated HCO_3^- [40]. The CCMs in cyanobacteria and microalgae enhance photosynthetic carbon fixation by energy-driven uptake of HCO_3^- and its conversion to CO_2 . The minimum requirements for a cyanobacterial CCM are one energy-driven active transporter that accumulates HCO_3^- in the cytosol, and an anion-permeable carboxysome containing intracellular carbonic anhydrase [41]. Central to the functioning of the cyanobacterial CCM is the carboxisome (Fig. 4), a protein micro-compartment within the cell that contains RuBisCO and a carbonic anhydrase. The latter converts HCO_3^- into CO_2 within the carboxysome, concentrating CO_2 up to 1000-fold around the active site of RuBisCO [40], thus overcoming the inefficiency of RuBisCO [24] and its inability to distinguish O_2 from CO_2 .




Figure 4. Central to the cyanobacterial CCM is the carboxysome, a cellular micro-compartment containing the RuBisCo and a carbonic anhydrase (CA) that by converting HCO_3^- into CO_2 rises CO_2 concentration up to 1000-fold around the active site of RuBisCO [40].

For eukaryotic algae with CCMs, the presence of pyrenoids is often associated with the occurrence of a CCM, although there are exceptions [41]. Eukaryotic algal CCMs likely originated independently in different clades of algae, as indicated by the diversity of inorganic C transporters and of carbonic anhydrases, as well as the lack of a strict correlation between occurrence of a CCM and the presence of a pyrenoid [41].

7. Phosphorous

Another essential element for life forms that has fluctuated downward in bioavailability through Earth history is phosphorous (P). Ranking 11th among the elements in the Earth's crust, P occurs primarily as phosphate (PO_4^{3-}) in apatite. Although found in most igneous and metamorphic rocks, mafic rocks commonly contain at least an order of magnitude more apatite than most felsic rocks (granites and rhyolites) [28]. Thus, as Fe- and Mg-rich materials have been concentrated in the Earth's mantle and core, so too has P. By the late Archean, continental-scale granitic cratons had developed on the Earth's crust, contributing to the decline in PO_4^{3-} in aquatic environments.

Phosphorus is one of the eleven macro-biogenic elements that make up living organisms [C, O, H, N, S, P, Na, K, Ca, Mg, Cl, Fe]. Bioavailable PO_4^{3-} is essential for all forms of life, although it has received less attention than C, N, and Fe in the evolution of autotrophy in relation to environmental changes during the last 4 Ga [42,43,44]. As noted in the Introduction, PO_4^{3-} is essential to cellular structure and processes, including nucleotides (DNA and RNA), [phospholipid](#) membranes, energy transfer and storage, and for synthesis of proteins and enzymes [26]. However, PO_4^{3-} became one of the most limiting nutrients for primary productivity in Phanerozoic marine ecosystems due to its low solubility in oxygenated waters and because, unlike bioavailable dissolved N (DIN), PO_4^{3-} cannot be fixed from the atmosphere [28,37,44].

Microbial uptake of PO_4^{3-} takes place through at least two kinetically distinct processes. A "low affinity component" operates continuously, apparently driven by proton-motive force. A "high affinity component" requires energy from ATP and operates when internal PO_4^{3-} pools are depleted. Was the decline in available PO_4^{3-} in Proterozoic oceans related to changes in major precipitation modes of CaCO_3 ?

A key factor in the bioavailability of PO_4^{3-} in the Archean and early Proterozoic oceans was the lack of free oxygen [36,46]. By the early Proterozoic, cyanobacteria were producing free oxygen, triggering the Great Oxidation Event (~2.4–2.1 Ga) (Fig. 2). With the emergence of O_2 into the atmosphere, terrestrial weathering dramatically changed,

resulting in oxidation of sulfides ubiquitous in continental rocks. A major result was the delivery of substantial amounts of sulfates to the oceans, and those dissolved sulfates provided the oxidation potential for sulphate-reducing microbes, which are obligate anaerobes. The result was an oxygenated mixed layer/photoc zone to perhaps 200 m depth, but the vast ocean depths remained anoxic and sulphidic (i.e., euxinic) through most of the Proterozoic [36,46].

The bioavailability of PO_4^{3-} declined in surface waters, based on the relative insolubility of PO_4^{3-} in oxygenated waters compared to its much greater solubility in euxinic waters. Shallow waters exposed to sunlight became depleted in PO_4^{3-} as a consequence of reduced solubility in terrestrial-runoff waters and in oxygenated surface waters, and further diminished by biological uptake by photosynthetic cyanobacteria and microalgae, and by geochemical uptake into ferric oxyhydroxides. At the same time, the much greater volumes of subsurface, aphotic waters, which were euxinic and therefore in which PO_4^{3-} was highly soluble, became major repositories. In that respect, like today, hydrodynamic processes such as mixing and upwelling must have been critical mechanisms for supplying PO_4^{3-} to photosynthesizing microorganisms in the surface waters.

By the late Neoproterozoic, sufficient photosynthetic oxygen production had occurred to oxidize most oceanic waters. Two additional contributors to the increase in atmospheric O_2 and dissolved O_2 throughout the oceans were primary production by eukaryotic algae and increased solubility of O_2 in cold oceanic waters associated with Cryogenian glaciation. The latter also would have increased the rates of ocean turnover and delivery to surface waters of dissolved PO_4^{3-} from subsurface waters (Fig. 5). Shen et al. [47] postulated that the oxygenation of the oceans allowed diverse animal lineages to increase in size.




Figure 5. Evolution of Phosphorous (P) concentration through the late Neoproterozoic and Cambrian and its relationship with the emergence of major groups of organisms. Notice that the first weakly calcified metazoans (red bar) appeared during a minimum of P at the end of the Ediacaran.

Modern lineages of metazoans with hard skeleton appeared by mid-Cambrian during another minimum in P concentration.

On scales of major plate-tectonics processes, massive basalt flows and tectonic uplift increase delivery of weathering-mobilized PO_4^{3-} to the oceans. On intermediate time scales, changes in rates of deep-ocean circulation influence rates of delivery of DIN and PO_4^{3-} to surface waters, where they promote primary productivity. In some cases, regional changes in ocean circulation have brought PO_4^{3-} rich subsurface waters into oxygenated conditions, creating major phosphatization events that have resulted in economically valuable phosphate ore deposits such as the prolific Lutetian-Messinian phosphorites of Morocco [48]. Most PO_4^{3-} in the ocean is cycling within the biotic system, and is returned to the ocean by microbial activity. Only a small fraction of the phosphorus is buried in sediments, as reduced conditions within the sediments promotes dissolution of PO_4^{3-} into sediment pore waters and its return to ocean cycling [49].

8. Calcification as a Sink and a Source of Both CO_2 and PO_4^{3-}

Precipitation of carbonate minerals is a long-term sink for CO_2 , but in the short term, can serve as a source (e.g., [35] and references therein). Calcification typically involves two bicarbonate ions, incorporating one CO_3^{2-} into the CaCO_3 mineral structure, and releasing the other into the environment as CO_2 [50]. In aquatic environments, photosynthetic uptake of CO_2 or HCO_3^- reduces availability of CO_2 , increases pH and can lead to oversaturation with respect to calcite. Dissolved PO_4^{3-} is adsorbed onto and co-precipitates with calcite, with incorporation of some of the surface PO_4^{3-} into the bulk structure as crystal growth proceeds (Fig. 6). However, higher concentrations of dissolved phosphate inhibit the growth of calcite crystals and can stop growth completely [51].




Figure 6. Simplified diagram of the adsorption process. Atoms, molecules or ions adhere to a surface by weak residual forces (e.g., van der Waals, electrostatic). It can be multilayer or monolayer.

Numerous studies have documented removal of PO_4^{3-} associated with CaCO_3 precipitation, for example during dense blooms of some cyanobacteria. Guldbrandsen and Cremer [52] experimentally demonstrated co-precipitation of CaCO_3 and PO_4^{3-} , and Kitano et al. [53] found that stirring increases the amount of PO_4^{3-} that co-precipitates. Millero et al. [54] demonstrated that PO_4^{3-} uptake on CaCO_3 minerals is a multistep process. Depending upon water chemistry and temperature, adsorption or desorption can occur rapidly, in minutes to a few hours, with much slower processes lasting more than one week. Up to 80% of the adsorbed PO_4^{3-} can be released from CaCO_3 over one day. The amount of PO_4^{3-} left on the CaCO_3 is close to equilibrium adsorption [54]. Thus, the

adsorption of PO_4^{3-} during CaCO_3 precipitation can be both a sink and a source of phosphates (Fig. 7).




Figure 7. Co-precipitation of phosphate and calcium carbonate.

9. Prokaryotic Organo-Sedimentary Systems

Stromatolites, the fossil evidence of calcified microbial mats, can be viewed as the first biotic process to produce a hard body [55]. By the Late Archaean, ooids—the second biotic process to produce a hard body—appeared when photosynthetic O_2 production by cyanobacteria was occurring. What might have been the advantage for these benthic microbial systems to produce coated-mineral structures? Such coatings would have restricted light penetration, as well as diffusion of DIC and nutrients required for photosynthesis, growth and reproduction. And what could be the advantage of inducing carbonate-mud precipitates (whittings) by neritic photosynthetic prokaryotes (picoplankton) and eukaryotes (phytoplankton)? This third microbially-mediated organo-sedimentary system to produce carbonate precipitates emerged by the Mesoproterozoic, following the Great Oxygenation Event (GOE) (Fig. 2).

Cyanobacteria have remarkably few nutritional requirements. They use light as source of energy, H_2O or H_2S as electron donors and CO_2 (or HCO_3^-) as the source of inorganic carbon [56]. Additionally, many species can fix atmospheric N_2 to NH_4^+ , which makes them independent of DIN sources such as nitrate, ammonium or organic nitrogen. Hence, the most critical nutrients for cyanobacteria are PO_4^{3-} , which can be stored intracellularly as polyphosphate, and, to a lesser extent, iron. Moreover, cyanobacteria can minimize their PO_4^{3-} requirements through the synthesis of sulfolipids [57], allowing them to colonize low-nutrient environments, as demonstrated by their ubiquity in both terrestrial and aquatic environments.

Thus, precipitation of CaCO_3 during the day augments CO_2 availability for photosynthesis [58], while PO_4^{3-} adsorption sequesters that scarce but essential nutrient. At night, desorption of PO_4^{3-} makes it available for uptake by the cells. In microbial mats (Fig.

8), in darkness, anoxic conditions resulting from continued sulfate reduction, aerobic respiration and sulfide oxidation likely contributes to the dissolution of CaCO_3 and increases the potential for PO_4^{3-} uptake by the microbial assemblage.




Figure 8. Oxygen and sulfide concentration (ppm) (in A) and pH (in B) variation during a diel cycle in the upper 12 mm of modern marine stromatolites in the Exuma Cays, Bahamas. (C): Light intensity and temperature at the surface of the stromatolite. (Adapted from [59]).

Giant ooids, associated with cyanobacterial mats and stromatolites, are common in Late Archaean rocks. They have been described in the 2.64 Ga old Ghaap Group in South Africa [60] and in the 2.72 Ga old Pilbara group in Australia [61], becoming abundant and characteristic of many Neoproterozoic carbonates [62]. Microbially-mediated organo-mineralization processes are likely involved in the origin of ooids. They can either be biologically induced (e.g., by-products of metabolic activities that increase environmental alkalinity and trigger carbonate precipitation), or biologically influenced (e.g., microbial extracellular polymeric substances serve as templates for carbonate mineralization [63]). Again, the advantage for the microbial consortium to precipitate CaCO_3 coatings around particles was likely in PO_4^{3-} adsorption, associated with the precipitation of metastable amorphous carbonate [64] (Fig. 9).




Figure 9. Conceptual multiphase model for the oolite cortex accretion, based on [64,65].

A significant source of lime-mud production in modern oligotrophic shallow marine and lacustrine environments also is linked to microbial calcification [66]. Whitings are the drifting milky clouds that are commonly observed in warm, shallow seas. Robbins et al. [67] estimated that whitings production might account for more than 40% of the bank-top- and peri-platform Holocene muds on the west side of Great Bahama Bank. The production of whitings is linked to blooms of cyanobacteria and planktic green algae in oligotrophic aquatic environments via CO_2 uptake [68]. Photosynthetic carbon fixation during blooms induces an increase in extracellular pH, favoring crystal nucleation [66]. What would it be the advantage for photosynthetic pico- and microplankton to induce carbonate-mud precipitation in the water? Blooming means that CO_2 and nutrient acquisition must keep pace with the increase of organic matter required for cell division. With the diurnal cycle of rapid adsorption and desorption of PO_4^{3-} , the milky cloud can efficiently capture this scarce nutrient, which would be of particular advantage in extremely oligotrophic conditions.

10. Bio-calcification in Photosynthetic Eukaryotes

The basic eukaryotic cell is descended from an anaerobic Proteoarchaeota that engulfed purple, non-sulfur α -proteobacteria that could utilize oxygen but were not obligate aerobes [36]. All algae have that basic eukaryotic cell, and all algal plastids are believed to have originated from cyanobacterial endosymbionts [69]. Molecular phylogenetic studies

have demonstrated that the morphological diversity of the algae results from their polyphyletic origins within the Eukarya [69,70]. An interesting question then, is: Why do some marine algae within different phylogenetic groups invest energy into producing CaCO_3 skeletal structures, whereas most do not?

Tropical macroalgae cope with three strong selective pressures: limited access to $\text{CO}_2/\text{HCO}_3^-$ due to warm water and competition among photosynthetic organisms, limited access to PO_4^{3-} , and herbivory. Calcification is clearly an anti-herbivory defense. And there are striking differences in percent tissue PO_4^{3-} between calcified and fleshy algae, indicating different nutrient acquisition strategies. The effects of nutrient enrichment with nitrogen (DIN) and PO_4^{3-} on productivity and calcification of fleshy and calcareous algae differ. DIN and PO_4^{3-} frequently enhance productivity of fleshy algae, but do not increase calcification rates of calcareous species [71].

Halimeda, a codiacean macroalga common in warm, shallow-marine environments, can add up to one segment per day, per branch (Fig. 10). However, the algae require nutrient uptake for growth and reproduction in warm waters where PO_4^{3-} levels are often below detection limits. Calcification occurs during daytime within an organic matrix in the outer utricle walls of 1-day-old segments. Photosynthetic removal of CO_2 by diffusion across a membrane and enzymatic anhydrase activity induces aragonite precipitation in the utricle space [72,73]. At night, respiration elevates CO_2 and decreases CO_3^{2-} saturation, which decreases seawater pH in the utricle space. The aragonite needles partially dissolve, break and recrystallize into micro-anhedral crystals ($<1\ \mu\text{m}$). Such partial dissolution at night would facilitate desorption of PO_4^{3-} from the skeleton




Figure 10. Calcification stages of *Halimeda* [72,73]. **A:** Growth of short skeletal needles on the utricle wall; **B:** Recrystallization of short needles to micron-sized anhedral crystals. **C:** secondary calcification by long and dense euhedral skeletal aragonite needles, **D:** the primary inter-utricular space In the rim of the segment is filled with micron-sized anhedral crystals.

Coralline red algae occur as thin crusts on hard substrata that can produce free-living rhodoliths, or as thalli with articulated branches. Most precipitate high-Mg calcite rather than aragonite. Calcification dynamics (Fig. 11) depend on nutrient concentrations; PO_4^{3-} inhibits the crystalline lattice and hampers calcite precipitation [74]. Enrichment of PO_4^{3-} , along with other nutrients, favors the growth of macroalgae and phytoplankton, which reduces water transparency and impedes coralline algal growth. Calcification associated with photosynthesis exhibits day-night cyclicity that is also recorded in diel pH fluctuations. Abiotic re-precipitation of CaCO_3 can occur in crustose coralline red algae during

nighttime [75] and pH drop at the boundary layer in darkness has been documented [76]. Thus, partial dissolution occurring at night would provide favorable conditions to remobilize adsorbed PO_4^{3-} from the calcitic skeleton and facilitate uptake by the algal cells.




Figure 11. Skeleton formation in red algae (based on Nash et al. [74]). (A) Primary cell wall is forming. Hemicellulose microfibrils act as nucleating substrate for micro-granules. (B) Cellulose microfibrils extrude from plasma membrane and leaks through broken primary cell wall. (C) Inter-filament grains initiate attached to the external part of the primary cell wall and growth in the leaked cellulose microfibril. (D) Secondary cell wall thickens, extruded cellulose microfibrils stops leaking and mineralizes as Mg-calcite. (E) Plasmalemma pushed inwards as cellulose microfibril continues to extrude and mineralize. Secondary cell wall fully formed.

Coccolithophores are photosynthetic protists that produce small calcitic disks (coccoliths). These oval-shaped plates of CaCO_3 consist of double discs, composed of radial arrays of minute, elaborately-shaped crystal units. They form delicate crystalline lace or open web-like patterns in which the rate of the CaCO_3 crystal surface/volume is very high. Originating by the Late Triassic (Fig. 2), they proliferated during the Cretaceous, and, together with planktic foraminifera, allowed widespread carbonate production in the open ocean [77]. Coccolithophores can produce as much as two coccoliths per hour [78]. The variability of coccolith shapes indicates they serve diverse functions. Young [79] noted features that defy simple physical functions, particularly complex mesh structures and the variety of elaborate forms, and suggested that coccolith might be adaptations for nutrient uptake.

Coccolithophores thrive in waters with minimal PO_4^{3-} , insufficient to promote growth of other phytoplankton. Coccolith formation is less adversely affected by nutrient deficiency than is cell division and growth [80]. The nutrient requirements for the organic cellular components of a coccolithophorid cell are similar to those for non-calcifying phytoplankton. However, forming coccoliths requires minimal nutrient cost, as coccolith production continues even when cell division ceases because of nutrient limitation [81].

In modern *E. huxleyi*, coccoliths nucleate and grow in a Golgi-derived coccolith vesicle, from where they migrate to the cell surface (Fig. 12). Within the coccolith vesicle, polysaccharides are thought to regulate nucleation and subsequent growth of coccoliths [78]. Sviben et al. [83] identified a reservoir compartment, distinct from the coccolith vesicle, with high concentrations of Ca^{2+} and PO_4^{3-} . Only Ca^{2+} is delivered to the coccolith vesicle [85], whereas PO_4^{3-} is diverted to the cell for growth and reproduction.




Figure 12. Calcification processes in coccolithophorids (based on [82, 83, 84, 85]. [A] Coccolith formation occurs within the Golgi-derived coccolith vacuole (CV). Intracellular CaCO_3 precipitation from bicarbonate (HCO_3^-) releases equimolar H^+ that must be rapidly removed from the CV to maintain pH for CaCO_3 precipitation [B]. Some H^+ may be utilized by photosynthesis [C]. Not yet fully understood, it is hypothesized Ca^{2+} to be recruited through Ca-channels and V-type ATPase pumps [D]. A reservoir compartment (RC) concentrates Ca^{2+} and P_i [E]. The coccolith-associated polysaccharide [F] may drive the Ca^{2+} from the coccosphere into the RC, from where only Ca^{2+} is derived to the coccolith vesicle (CV) and phosphorous is diverted to the cell for algal growth and reproduction. Transfer of Ca^{2+} from the RC into the CV uses Ca^{2+} channels and transporters.

11. CO_2 and Carbonate Depositional History

In aqueous solutions, CO_3^{2-} , HCO_3^- , CO_2 , and H_2CO_3 co-exist in dynamic equilibria. When CO_2 reacts with water (H_2O), it complexes to H_2CO_3 , which dissociates to H^+ and HCO_3^- , thereby increasing the concentration of H^+ and cause the seawater acidity to increase (i.e., the pH to decline). When CaCO_3 (e.g., limestone) reacts with free hydrogen (H^+) ions in seawater, the solid CaCO_3 dissolves, forming free calcium (Ca^{2+}) ions and free bicarbonate (HCO_3^-) ions. As noted previously, the concentrations of ions of carbonate, bicarbonate, phosphate, borate, orthosilicate, sulfides, and organic acids constitute alkalinity in aquatic environments, especially HCO_3^- . The consumption of protons results in decreased H^+ activity. Thus, CaCO_3 acts to neutralize or buffer the solution by consuming H^+ .

In the modern world, burning of fossil fuels, combined with widespread changes in land use, have resulted in rapidly increasing concentrations of atmospheric CO_2 that are causing the decline in the pH of surface seawater (e.g., [6–9]). Will the progressive warming and acidification cause mass extinctions and evolutionary turnover of marine biotas, as has been documented by paleoclimate evidence [10]?

From this question emerges another major “carbonate paradox”. Contrary to predictions of increased carbonate dissolution in response to increasing CO_2 , extended high CO_2 Greenhouse World periods during the Phanerozoic (Silurian–Devonian and Jurassic–Cretaceous) were characterized by prolific accumulations of biogenic carbonates, especially those associated with cyanobacterial and algal calcification (Fig. 2). In the Cenozoic, coral reefs emerged as major carbonate factories as Icehouse World conditions developed and

became predominant in the Oligocene through the Holocene, a conundrum noted many years ago [25].

The solution to this paradox lies in two parts. First, there is the “rate” factor. Mass extinction events in the Phanerozoic record are characterized by carbonate depositional hiatuses that indicate times of global carbonate dissolution [86]. Very rapid increases in atmospheric CO₂ (or other atmospheric compounds that unite with water to form acids, e.g., SO₂ + H₂O → H₂SO₄ + 2H⁺), whether caused by massive extrusion of flood basalts, bolide impact, methane-hydrate release, extensive burning of fossil fuels, or some combination of events, can indeed trigger carbonate dissolution resulting in hiatuses of a hundred thousand years (Paleocene–Eocene) or hundreds of thousand years (Cretaceous–Paleocene) or millions of years (Permian–Triassic).

But eventually dissolution, combined with high atmospheric CO₂, over time will increase alkalinity and restore the carbonate factory. The massive accumulations of Silurian–Devonian and Jurassic–Cretaceous carbonates occurred during times of relatively rapid sea-floor spreading, which elevated both CO₂ and Ca²⁺ concentrations, triggered warming temperatures, sea-level rise, and expansive areas of relatively shallow basins and shelves. Reduced land areas limited freshwater input of dissolved nutrients, and warm waters reduced rates of deep-ocean circulation. During Greenhouse-world conditions, the trophic resource continuum expanded (compared to Icehouse conditions), with limited extremely rich regions of upwelling and expansive regions of extreme oligotrophy, fostering extensive production of carbonates [87]. Elevated evaporation rates, relatively shallow seas, and high alkalinity favored calcifying cyanobacterial and microalgal taxa that could sequester scarce PO₄³⁻. The result, massive carbonates dominated by carbonate muds or chalks.

During daylight hours, in warm, shallow waters, abundant microalgae take up available DIN and PO₄³⁻, while high rates of photosynthesis deplete immediate access to CO₂/HCO₃⁻. Calcification associated with photosynthesis alleviates both deficiencies. Two HCO₃⁻ provides one CO₂ for photosynthesis and a CO₃²⁻ for calcification, somewhat alleviating local CO₂ depletion. At the same time, PO₄³⁻ adsorption occurs during daytime calcification, and is partially desorbed at night, becoming available for uptake by the cyanobacteria or algae. In addition, some dissolution of the CaCO₃ releases HCO₃⁻, and along with community respiration, results in early morning maximum concentrations of HCO₃⁻. With the onset of light, photosynthesis makes energy available to CCMs, providing the primary producers with access to and storage of CO₂, which becomes increasingly unavailable over the course of the day.

And finally, why did extensive coral reefs only become prevalent with the onset of the Icehouse World climates [25]. Again, the answer is likely climatic and geochemical influences on biological processes [88,89]. High-latitude cooling, compression of tropical habitats, and increasing temperature gradients between high and low latitudes [90] and between surface and deeper waters, were major factors. Coral communities flourished circumtropically beginning in the late Oligocene. By the early Miocene, reefs and associated biotas expanded latitudinal distributions by more than 10° north and south [91]. The expansion in reef-building capacity corresponded to increasing Mg/Ca ratios in seawater [92] and falling atmospheric CO₂ concentrations [93], both of which promoted aragonite precipitation in warm, tropical waters. The majority of extant Symbiodiniaceae lineages diversified since Middle Miocene [94]. Some global cooling, including much cooler subsurface waters, likely benefited the coral-Symbiodiniaceae symbioses, which are sensitive to photo-oxidative stress under elevated temperatures [95]. The substantial increase in rates of circulation of deep, cold waters resulted in more “intermediate” conditions of oceanic oligotrophy [85]. Pomar and Hallock [87] further postulated that Neogene cooling supported the co-evolution and expansion of zooxanthellate corals and coralline algae into shallow, high-energy waters, where their carbonate production potential was highest. In such environments, both calcification and PO₄³⁻ extraction are also potentially optimized.

12. Conclusions

Photosynthesis typically involves daytime CO₂ uptake and O₂ release predominating over respiration; at night, respiration consumes O₂ and releases CO₂. The result is strong diel variation in both pH and dissolved O₂ concentrations in aquatic environments. In warm-water environments in which alkalinity is relatively high and PO₄³⁻ concentrations are minimal, these variations have the potential to promote precipitation of CaCO₃ and associated adsorption of PO₄³⁻ in daylight when photosynthesis is active. At night, lower pH and oxygen availability can promote some CaCO₃ dissolution and PO₄³⁻ desorption. This combination of calcification and PO₄³⁻ extraction occurs in photosynthesizing cyanobacteria, as well as in a diverse array of calcifying nanophytoplankton and calcareous macroalgae, indicating that this process allows both prokaryotic and eukaryotic photosynthetic organisms to thrive in warm, alkaline, oligotrophic waters. Concentrations of both atmospheric CO₂ and surface-ocean PO₄³⁻, which are essential for biological productivity, have declined by at least 1–2 orders of magnitude over Earth history. The “Phosphate Extraction Mechanism”, associated with photosynthetically-induced calcification, has played a major role in the production and accumulation of carbonates throughout much of the Proterozoic and the Phanerozoic. Recognition of this relationship helps resolve the apparent paradox that periods in the Phanerozoic when atmospheric CO₂ levels were considerably higher than present and predicted Anthropocene concentrations, were times of massive accumulation of carbonates, predominantly produced by photosynthesizing cyanobacteria and calcifying algae. This synthesis also stresses the importance of rates of change, as times of rapid increase in atmospheric CO₂ concentrations were associated with mass extinction events characterized by global carbonate-depositional hiatuses, requiring hundreds of thousands or even millions of years for surface waters of the oceans to regain sufficient alkalinity to sustain accumulation of massive carbonates.

Acknowledgments: Prof Gabriel Moyà (UIB) for the seminal discussions on this topic, and to the volume editor, H. Westphal, for her patience and suggestions.

Author Contributions: Conceptualization: LP, PH, GMV and JIB; Methodology: LP, PH, GMV and JIB; Investigation: LP, PH, GMV and JIB; Writing: Original Draft Preparation, LP and PH; Writing: Review & Editing, PH and GMV.

References

1. Keeling, C. D. The concentration and isotopic abundances of carbon dioxide in the atmosphere. *Tellus* 1960, v. 12, p. 200–203.
2. Scripps CO₂ Program. Global Stations CO₂ Concentration Trends. 2022. https://scrippsco2.ucsd.edu/graphics_gallery/other_stations/global_stations_CO2_concentration_trends.html (accessed on 19/09/2022).
3. Harris, D. C. Charles David Keeling and the story of atmospheric CO₂ measurements. *Analytical Chemistry* 2010, v. 82, p. 7865–7870.
4. IPCC (Intergovernment Panel on climate Change), <https://www.ipcc.ch/about/>; (accessed on 22/08/2022).
5. Kleypas, J. A., Buddemeier, R. W., Archer, D., Gattuso, J-P., Langdon, C., Opdyke, B. N. Geochemical consequences of increased atmospheric CO₂ on coral reefs. *Science* 1999, v. 284, p. 118–120.
6. Caldeira, K., Wickett, M. E. Anthropogenic carbon and ocean pH. *Nature* 2003, v. 425, p. 365–365.
7. Wigley, T.M. A combined mitigation/geoengineering approach to climate stabilization. *Science* 2006, v. 314(5798), p. 452–454.
8. Wolf-Gladrow, D., Riebesell, U., Burkhardt, S., Bijma, J. Direct effects of CO₂ concentration on growth and isotopic composition of marine plankton. *Tellus* 1999, v. 51B, p. 461–476.
9. Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R. M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., Najjar, R. G., Plattner, G-K., Rodgers, K. B., Sabine, C. L., Sarmiento, J. L., Schlitzer, R., Slater, R. D., Totterdell, I. J., Weirig, M-F., Yamanaka, Y., Yool, A. Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature* 2005, v. 437(29), p. 681–686.
10. Clarkson, M. O., Kasemann, S. A., Wood, R. A., Lenton, T. M., Dains, S. J., Richoz, S., Ohnemüller, F., Meixner, A., Poulton, S. W., Tipper, E. Ocean acidification and the Permo-Triassic Mass Extinction. *Science* 2015, v. 348, p. 229–232.
11. Young, J. R. Possible functional interpretations of coccolith morphology. *Abh. Geol. Bundesanstalt-A* 1987, v. 39, p. 305–313.
12. Dodd, M., Papineau, D., Grenne, T., Slack, J. F., Rittner, M., Pirajno, F., O’Neil, J., Little, C. T. S. Evidence for early life in Earth’s oldest hydrothermal vent precipitates. *Nature* 2017, v. 543, 60–64.

13. Nutman, A. P., Bennett, V. C., Friend, C. R. L., Van Kranendonk, M. J., Chivas, A. R., Rapid emergence of life shown by discovery of 3,700-million-year-old microbial structures. *Nature* 2016, v. 537, p. 535–538. 586
14. Krissansen-Totton, J., Arney, G. N., Catling, D. C. Constraining the climate and ocean pH of the early Earth with a geological carbon cycle model. *Proc. Natl. Acad. Sci. USA* 2018, v. 115, p. 4105–4110. 587
15. Holland, H. D. The oxygenation of the atmosphere and oceans. *Phil. Trans. Royal Soc. B-Biol. Sci.* 2006, v. 362, p. 903–915. 588
16. Kasting, J. F. Earth's early atmosphere. *Science* 1993, 259(5097), p. 920–926. 589
17. Kaufman, A. J., Xiao, S. High CO₂ levels in the Proterozoic atmosphere estimated from analyses of individual microfossils. *Nature* 2003, v. 425, p. 279–282. 590
18. Kah, L.C., Bartley, J.K. Effect of marine carbon reservoir size on the duration of carbon isotope excursions. Interpreting the Mesoproterozoic carbon isotope record. *Geol. Soc. of Amer. Abstr.* 2004, v. 36, no. 5, p. 78. 591
19. Berner, R. A., Kothavala, Z. GEOCARB III: A revised model of atmospheric CO₂ over Phanerozoic time. *Amer. J. Sci.* 2001, v. 301, p. 182–204. 592
20. Royer, D. L., Berner, R. A., Montañez, I. P., Tabor, N. J., Beerling, D. J. CO₂ as a primary driver of Phanerozoic climate. *GSA Today* 2014, v. 14; no. 3, doi: 10.1130/1052-5173(2004)0142.0.CO;2. 593
21. Kah, L. C., Riding, R. Mesoproterozoic carbon dioxide levels inferred from calcified cyanobacteria. *Geology* 2007, v. 35, p. 799–802. 594
22. Sheldon, N. D. Precambrian paleosols and atmospheric CO₂ levels. *Precam. Res.* 2006, v. 147, p. 148–155. 595
23. Sleep, N. H., Zahnle, K. Carbon dioxide cycling and implications for climate on ancient Earth. *J. Geophys. Res.* 2001, v. 106, p. 1373–1399. 596
24. Riding, R. Cyanobacterial calcification, carbon dioxide concentrating mechanisms, and Proterozoic–Cambrian changes in atmospheric composition. *Geobiology* 2006, v. 4, p. 299–316. 597
25. Frost, S. H. Cenozoic reef systems of Caribbean. Prospects for paleoecologic synthesis. *Am. Assoc. Petrol. Geol. Stud. Geol.* 1977, v. 4, p. 93–110. 598
26. Kempe, S., Degens, E. T. An early soda ocean. *Chem. Geol.* 1985, v. 53, p. 95–108. 599
27. Crockford, P., Halevy, I. Questioning the paradigm of a phosphate-limited Archaean biosphere. *Geophys. Res. Lett.* 2022, 49, e2022GL099818. 600
28. Ingalls, M., Grotzinger, J. P., Present, T., Rasmussen, B., Fischer, W. W. Carbonate-associated phosphate (CAP) indicates elevated phosphate availability in Neoproterozoic shallow marine environments. *Geophys. Res. Lett.* 2022, 49, e2022GL098100. 601
29. Zahnle, K., Arndt, N., Cockell, C., Halliday, A., Nisbet, E., Selsis, F., Sleep, N. H. Emergence of a habitable planet. *Space Sci. Rev.* 2007, v. 129, p. 35–78. 602
30. Sleep, N.H. The Hadean-Archaeon environment. *Cold Spring Harb Perspect Biol.* 2010, v. 2(6), a002527. 603
31. O'Neil, J., Carlson, R. W., Francis, D., Stevenson, R. K. Neodymium-142 evidence for Hadean mafic crust. *Science* 2009, v. 321, p. 1828–1831. 604
32. Sleep, N. H., Zahnle, K., 2001, Carbon dioxide cycling and implications for climate on ancient Earth. *J. Geophys. Res.* 2001, v. 106, p. 1373–1399. 605
33. Nutman, A. P., Bennett, V. C., Friend, C. R. L., Van Kranendonk, M. J., Chivas, A. R. Rapid emergence of life shown by discovery of 3,700-million-year-old microbial structures. *Nature* 2016, v. 537, p. 535–538. 606
34. Goldblatt, C., Zahnle, K. J. . Clouds and the faint young Sun paradox. *Climate of the Past* 2011, v. 7(1), p. 203–220. 607
35. Hallock, P. Changing influences between life and limestones in Earth history. In *Coral Reefs in the Anthropocene*, Birkeland, C. Ed.; Springer, Dordrecht, Netherlands, 2015, pp. 17–42. 608
36. Mentel, M., Martin, W. Energy metabolism among eukaryotic anaerobes in light of Proterozoic ocean chemistry. *Phil. Trans. Royal Soc. B-Biol. Sci.* 2008, v. 363, p. 2717–2729. 609
37. Ward, L. M., Rasmussen, B., Fischer, W. W. Primary productivity was limited by electron donors prior to the advent of oxygenic photosynthesis. *J. Geophys. Res.: Biogeosci.* 2019, 124(2), 211–226. 610
38. Fischer, W. W., Hemp, J., Johnson, J. E. Evolution of oxygenic photosynthesis. *Ann. Rev. Earth Planet. Sci.* 2016, 44(1), 647–683. 611
39. Erb, T. J., Zarzycki, J. A short history of RubisCO: the rise and fall (?) of Nature's predominant CO₂ fixing enzyme. *Curr. Opin. Biotech.* 2018, v. 49, p. 100–107. 612
40. Badger, M. R., Price, G. D. CO₂ concentrating mechanisms in cyanobacteria: molecular components, their diversity and evolution. *J. Exper. Bot.* 2003, v. 54, p. 609–622. 613
41. Raven, J.A., Beardall, J., Sánchez-Baracaldo, P. The possible evolution and future of CO₂-concentrating mechanisms. *J. Exper. Bot.* 2017, v. 68, p. 3701–3716. 614
42. Papineau, D. Global biogeochemical changes at both ends of the Proterozoic: Insights from phosphorites. *Astrobiology* 2010, v. 10, p. 165–181. 615
43. Large, R. R., Halpin, J. A., Lounejeva, E., Danyushevsky, L. V., Maslennikov, V. V., Gregory, D., Sack, P. J., Haines, P. W., Long, J. A., Makoundi, C., Stepanov, A. S. Cycles of nutrient trace elements in the Phanerozoic ocean. *Gondwana Res.* 2015, v. 28, p. 1282–1293. 616
44. Pufahl, P. K., Groat, L.A. Sedimentary and igneous phosphate deposits: Formation and exploration: An invited paper. *Econ. Geol.* 2017, v. 112, p. 483–516. 617
45. Paytan, A., McLaughlin, K. The oceanic phosphorus cycle. *Chem. Rev.* 2007, v. 107, p. 563–576. 618

46. Poulton, S. W., Canfield, D. E. Ferruginous conditions: A dominant feature of the ocean through Earth's history. *Elements* 2011, v. 7, p.107–112 646
47. Shen, J., Webb, G.A., Jell, J. S. Platform margins, reef facies and microbial carbonates; a comparison of Devonian reef complexes in the Canning Basin, Western Australia and the Guilin region, South China. *Earth-Sci. Rev.* 2008, v. 88, p. 33–59. 647
48. Chernoff, C. B., Orris, G. J. Data set of world phosphate mines, deposits, and occurrences: Part A. geologic data; Part B. location and mineral economic data. *U.S. Geol. Surv. Open-File Report*, C6, 2002, p. 680. 648
49. Baturin, G. N. Phosphorus cycle in the ocean. *Lithol. Min. Res.* 2003, v. 38, p. 101–119. 649
50. Vescei A., Berger W. H. Increase of atmospheric CO₂ during deglaciation: constraints on the coral reef hypothesis from patterns of deposition. *Global Biogeochem Cycles* 2004, 18(1):GB1035 650
51. Danen-Louwerse, H. J., Lijklema, L., Coenraats, M., Coprecipitation of phosphate with calcium carbonate in Lake Veluwe. *Water Resources* 1995, v. 29, p. 1781–1785. 651
52. Gulbrandsen, R. A., Cremer, M. Coprecipitation of carbonate and phosphate from seawater. *U.S. Geol. Surv. Prof. Pap.* 1970, v. 700-C, p. C125–C126. 652
53. Kitano, Y., Okumura, M., Idogaki, M. Uptake of phosphate ions by calcium carbonate. *Geochem. Jour.* 1978, v. 12, p. 29–37. 653
54. Millero, F., Huang, F., Zhu, X., Liu, X., Zhang, J.-Z. Adsorption and desorption of phosphate on calcite and aragonite in seawater. *Aquatic Geochem.* 2001, v. 7, p. 33–56. 654
55. Dupraz, C., Reid, R. P., Braissant, O., Decho, A. W., Norman, R. S., Visscher, P. T. Processes of carbonate precipitation in modern microbial mats. *Earth-Sci. Rev.* 2009, v. 96, p. 141–162. 655
56. Stal, L. J. Coastal microbial mats: the physiology of a small-scale ecosystem. *South African J. Bot.* 2001, v. 67, p. 399–410.. 656
57. Van Mooy, B. A. S., Rocap, G., Fredricks, H. F., Evans, C. T., Devol, A. H. Sulfolipids dramatically decrease phosphorus demand by picocyanobacteria in oligotrophic marine environments. *Proc. Natl. Acad. Sci. USA* 2006, v. 103, p. 8607–8612. 657
58. McConnaughey, T. A., Whelan, J. F. Calcification generates protons for nutrient and bicarbonate uptake. *Earth-Sci. Rev.* 2006, v. 42, p. 95–117. 658
59. Visscher, P. T., Hoefft, S. E., Surgeon, T.-M. L., Rogers, D. R., Bebout, B. M., Thompson, J. S., Reid, R. P. Microelectrode measurements in stromatolites. Unraveling the Earth's past? In *Environmental Electrochemistry: Analyses of Trace Element Biogeochemistry*, Taillefert, M, Rozan, T, Eds. *Am. Chem. Soc. Symp. Ser. 811*. Oxford Univ Press: New York, NY, 2002, p. 265–282. 659
60. Wright, D. T., Altermann, W. Microfacies development in Late Archaean stromatolites and oolites of the Ghaap Group of South Africa. *Geol. Soc. London Spec. Pub.* 2000, v. 178, p. 51–70. 660
61. Batchelor, M. T., Burne, R. V., Henry, B. I., Li, F., Paul, P. 2018, A biofilm and organomineralisation model for the growth and limiting size of ooids. *Sci. Rep.* 2018, v. 8, p. 559. 661
62. Grotzinger, J. P., James, N. P. Precambrian carbonates: Evolution of understanding; In *Carbonate Sedimentation and Diagenesis in the Evolving Precambrian World*; Grotzinger, J. P., James, N. P. Eds., SEPM Society for Sedimentary Geology Spec. Publ., Tulsa, OK; 2000, v. 67, p. 3–20. 662
63. Brehm, U., Krumbein, W. E., Palinska, K. A. Biomicrospheres generate ooids in the laboratory. *Geomicrobiol. J.* 2006, v. 23, p. 545–550. 663
64. Diaz, M. R., Eberli, G. P., Blackwelder, P., Phillips, B., and Swart, P.K., 2017, Microbially mediated organomineralization in the formation of ooids. *Geology*, v. 45, p. 771–774. 664
65. Diaz, M. R., Eberli, G. P. Decoding the mechanism of formation in marine ooids: A review. *Earth-Sci. Rev.* 2019, v. 190, p. 536–556. 665
66. Yates, K. K., Robbins, L. L. Microbial lime-mud production and its relation to climate change. In *Geological perspectives of global climate change*, Gerhard, L.C., Harrison, W.H., Hanson, B. M., Eds., AAPG Stud. in Geol., Tulsa, OK, USA, 2001, v. 47, p. 267–283. 666
67. Robbins, L. L., Tao, Y., Evans, C. A., Temporal and spatial distribution of whittings on Great Bahama Bank and a new lime mud budget. *Geology* 1997, v. 25, p. 947–950. 667
68. Robbins, L. L., Blackwelder, P. L. Biochemical and ultrastructural evidence for the origin of whittings: A biologically induced calcium carbonate precipitation mechanism. *Geology* 1992, v. 20, p. 464–468. 668
69. Bhattacharya, D., Medlin, L. The phylogeny of plastids: A review based on comparisons of small-subunit ribosomal RNA coding regions. *J. Phycol.* 1995, v. 31, p. 489–498. 669
70. Stiller, J.W. and Hall, B.D., 1997, The origin of the red algae: implications for plastid evolution. *P. Natl. Acad. Sci-Biol* 1997, v. 94, p. 4520–4525. 670
71. Demes, K. W., Littler, M. M., Littler, D. S. Comparative phosphate acquisition in giant-celled rhizophytic algae (Bryopsidales, Chlorophyta): Fleshy vs. calcified forms. *Aquat. Bot.* 2010, v. 92, p. 157–160. 671
72. Wize mann, A., Meyer, F. W., Westphal, H. A new model for the calcification of the green macro-alga *Halimeda opuntia* (Lamouroux). *Coral Reefs* 2015, v. 33, p. 951–964. 672
73. Wize mann, A., Meyer, F. W., Hofmann, L. C., Wild, C., Westphal, H. Ocean acidification alters the calcareous microstructure of the green macro-alga *Halimeda opuntia*. *Coral Reefs* 2015, v. 34, p. 941–954. 673
74. Nash, M. C., Diaz-Pulido, G., Harvey, A. S., Adey, W. Coralline algal calcification: A morphological and process-based understanding. *PLOS ONE* 2019, v. 14, p. e0221396. 674

75. Comeau, S., Edmunds, P. J., Spindel, N. B., Carpenter, R. C. The responses of eight coral reef calcifiers to increasing partial pressure of CO₂ do not exhibit a tipping point. *Limnol. Oceanogr.* 2003, v. 58, p. 388–398. 705
76. Hurd, C. L., Cornwall, C. E., Currie, K., Hepburn, C. D., McGraw, C. M., Hunter, K. A., Boyd, P. W. Metabolically induced pH fluctuations by some coastal calcifiers exceed projected 22nd century ocean acidification: a mechanism for differential susceptibility? *Global Change Biol.* 2011, v. 17, p. 3254–3262. 706
77. Hay, W. W. Carbonate fluxes and calcareous nannoplankton. In *Coccolithophores—From Molecular Processes to Global Impact*; Thierstein, H., Young, J. R., Eds.; Springer, New York, USA, 2004, p. 509–527. 707
78. Monteiro, F. M., Bach, L. T., Brownlee, C., Bown, P., Rickaby, R. E. M., Poulton, A. J., Tyrrell, T., Beaufort, L., Dutkiewicz, S., Gibbs, S., Gutowska, M. A., Lee, R., Riebesell, U., Young, J., Ridgwell, A. Why marine phytoplankton calcify. *Sci. Adv.* 2016, v. 2 (7), e1501822. 708
79. Young, J. R. Possible functional interpretations of coccolith morphology. *Abh. Geolog. Bundesanstalt-A* 1987, v. 39, p. 305–313. 709
80. Paasche, E. Roles of nitrogen and phosphorus in coccolith formation in *Emiliania huxleyi* (Prymnesiophyceae). *Eur. J. Phycol.* 1998, v. 33, p. 33–42. 710
81. Sheward, R. M., Poulton, A. J., Gibbs, S. J., Daniels, C. J., Bown, P. R. Physiology regulates the relationship between coccosphere geometry and growth phase in coccolithophores. *Biogeosciences* 2017, v. 14, p. 1493–1509. 711
82. Brownlee, C., Wheeler, G. L., and Taylor, A. R. Coccolithophore biomineralization: new questions, new answers. *Sem. Cell Devel. Biol.* 2015, v. 46, p. 11–16. 712
83. Sviben, S., Gal, A., Hood, M. A., Bertinetti, L., Politi, Y., Bennet, M., Krishnamoorthy, P., Schertel, A., Wirth, R., Sorrentino, A., Pereiro, E., Faivre, D., and Scheffel A. A vacuole-like compartment concentrates a disordered calcium phase in a key coccolithophorid alga. *Nature Com.* 2016, v. 7, p. 11228. 713
84. Walker, C. E., Heath, S., Salmon, D. L., Smirnoff, N., Langer, G., Taylor, A. R., Brownlee, C., Wheeler, G.L. An extracellular polysaccharide-rich organic layer contributes to organization of the coccosphere in coccolithophores. *Front. Mar. Sci.* 2018, v. 5, Art. 306. 714
85. Gal, A., Sviben, S., Wirth, R., Schreiber, A., Lassalle-Kaiser, B., Faivre, D., Scheffel, A. Trace-element incorporation into intracellular pools uncovers calcium-pathways in a coccolithophore. *Adv. Sci.* 2017, v. 4, p. 1700088. 715
86. Newell, N.D. Mass extinctions at the end of the Cretaceous Period. *Science* 1965, v. 149 (3687), p. 922–924. 716
87. Hallock, P. Fluctuations in the trophic resource continuum: A factor in global diversity cycles? *Paleoceanography* 1987, v. 2, p. 457–471. 717
88. Hallock, P. Why are larger foraminifera large? *Paleobiology* 1985, v. 14, p. 250–261. 718
89. Pomar, L. Hallock, P. Carbonate factories: A conundrum in sedimentary geology. *Earth-Sci. Rev.* 2008, v. 87, p. 134–169. 719
90. Lear, C. H., Rosenthal, Y., Wright, J. D. The closing of a seaway: Ocean water masses and global climate change. *Earth Planet. Sci. Lett.* 2003, v. 210, p. 425–436. 720
91. Adams, C. G., Lee, D. E., Rosen, B. R. Conflicting isotopic and biotic evidence for tropical sea-surface temperatures during the Tertiary. *Palaeogeog. Palaeoecol.* 1990. v. 77, p. 289–313. 721
92. Stanley, S.M., Hardie, L.A. Secular oscillations in the carbonate mineralogy of reef-building and sediment-producing organisms driven by tectonically forced shifts in seawater chemistry. *Palaeogeog. Palaeoecol.* 1998, v. 144, p. 3–19. 722
93. Pearson, P. N., Palmer, M. R. Atmospheric carbon dioxide concentrations over the past 60 million years. *Nature* 2000, v. 406, p. 695–699. 723
94. Pochon, X., Montoya-Burgos, J. I., Stadelmann, B., Pawlowski, J. Molecular phylogeny, evolutionary rates, and divergence timing of the symbiotic dinoflagellate genus *Symbiodinium*. *Mol. Phylogenet. Evol.* 2006, v. 38, p. 20–30. 724
95. Wooldridge, S. A. Breakdown of the coral-algae symbiosis: Towards formalising a linkage between warm-water bleaching thresholds and the growth rate of the intracellular zooxanthellae. *Biogeosciences* 2013, v. 10, p. 1647–1658 725