

# Article

## Different responses to adventitious rhizogenesis under indole-3-butyric acid and seaweed extracts in ornamentals: first results in *Photinia x Fraseri* 'Red Robin'.

Danilo Loconsole<sup>1\*</sup>, Anna Elisa Sdao<sup>1</sup>, Giuseppe Cristiano<sup>1</sup> and Barbara De Lucia<sup>1</sup>

<sup>1</sup> Department of Soil, Plant and Food Sciences (Di.S.S.P.A.), University of Bari, "Aldo Moro", Italy. Via Amendola 165/A, 70126 Bari, Italy;

\* Correspondence: danilo.loconsole@uniba.it (D.L.); phone: +39-080-544-30-39 giuseppe.cristiano@uniba.it (G.C.); anna.sdao@uniba.it (A.E.S.); barbara.delucia@uniba.it (B.D.L.).

**Abstract:** Fraser's photinia 'Red Robin' (*Photinia x fraseri* Dress, *Rosaceae* family) is an important primary ornamental landscaping species with optimal hedge or screen effects and low maintenance, but it is difficult to root when propagated by cuttings, although high concentrations of phytohormones are used to optimize rhizogenesis. To our knowledge, there is currently no feasible enhanced method for photinia vegetative propagation through stem cuttings, using seaweed extract-based biostimulants as root promoters. Given the economic importance of the species, this research aims to assess the effects of indole-3-butyric acid (IBA) and seaweed extract-based stimulators on the quality of photinia 'Red Robin' cuttings, in terms of rooting indicators and ground and aboveground agronomic features. The treatments applied were different concentrations of commercial rooting stimulators compared to an untreated control: C0: distilled water; Rhizopon AA: 1% IBA (R1); Kelpak®: 2 mL L<sup>-1</sup> (K2); Kelpak®: 3 mL L<sup>-1</sup> (K3); Goteo®: 2 mL L<sup>-1</sup> (G2); Goteo®: 3 mL L<sup>-1</sup> (G3). The first results showed different responses to adventitious rhizogenesis under IBA and both seaweed extract treatments. At 70 DAC (days after cutting), the seaweed extract stimulated the production of over 80% of cuttings with callus; at 240 DAC, the percentage of rooted cuttings treated under R1 was the highest = 34.3%; the worst results were obtained by both biostimulant treatments at the highest doses: K3 = 21.3% and G3 = 20.7%. Furthermore, R1 produced 3.07 roots per cutting, which was 50% higher than the average of all other treatments. The applications of Kelpak® and Goteo® biostimulants, at both concentrations, resulted in an inhibition of root length with values below the untreated control. Rooted cuttings under R1 showed the highest ground (0.35g) and aboveground (0.47) dry value. Neither seaweed extract, Kelpak® or Goteo®, at different concentrations, improved both ground and aboveground weights of rooted cutting, compared to untreated control, indicating that these natural products are not suitable for Fraser's photinia 'Red Robin' propagation using this methodology. The overall quality of cuttings in IBA treatment was the strongest, with 1%, being the optimum concentration. Further research must be conducted to propose effective agronomic protocols by investigating: application methods, doses and number of applications and to clarify the biochemical and molecular mechanisms of action of these seaweed extracts.

**Citation:** To be added by editorial staff during production.

Academic Editor: Firstname  
Lastname

Received: date  
Revised: date  
Accepted: date  
Published: date



**Copyright:** © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Keywords:** *Ascophyllum nodosum*, auxin, biostimulants, cutting, *Ecklonia maxima*, ornamental plants, quality rooted cutting, root architecture, vegetative propagation.

## 1. Introduction

Nowadays, one of the critical challenges to address is increasing global agricultural production and crop quality, through reducing the impact of agriculture on ecosystems [1, 2]. Increasing the quality and yield of crops, maintaining agroecosystems, reducing chemical substances, increasing nutrient uptake and use efficiencies, and stimulating the plant's natural defense, must be the major goals [3, 4, 5]. The quality of propagation material is one of the most important factors for increasing the productivity of any ornamental crop. Several authors have dealt with defining the quality of the propagation material from a genetic, sanitary and agronomic point of view. From a genetic point of view, quality is related to the endogenous content of auxins [6]. Sanitary quality is characterized by the absence of parasites and pathogens [7]; the agronomic quality of a rooted cutting is based on the morphological features of the aerial parts and adventitious roots, which influence health, vigor and uniformity [8, 9]. A high number of very fine roots, which are essential for continuous access to water and nutrients, can help the plant withstand transplant shock, increasing survival and plant growth [10]. The process of adventitious root formation is influenced by a number of internal and external factors. Among the internal factors, the most important role is ascribed to phytohormones, especially auxins. Natural (indole-3-butyric acid, IBA) and synthetic auxins (naphthalene acetic acid, NAA), applied exogenously to cuttings [11], play a key role in generating adventitious roots [12, 13] and balanced shoots in ornamental species that are recalcitrant to clonal propagation, due to a low concentration of endogenous auxins [10, 14 – 22]. Commercial products based on indole-3-butyric acid (IBA), such as Rhizopon, are one of the most widely used exogenous sources of auxin and they can be delivered to cuttings in talc or dissolved in alcohol to be used as a quick dip [23]. The most recent literature has expanded the research on other natural substances with rooting activity such as biostimulants [24, 25, 26]. Biostimulants are defined as “product stimulating plant nutrition processes, independently of the product's nutrient content, with the sole aim of improving one or more of the following characteristics of the plant or the plant rhizosphere: (a) nutrient use efficiency; (b) tolerance to abiotic stress; (c) quality traits; (d) availability of confined nutrients in soil or rhizosphere” [27].

Seaweed-extract based biostimulants [28], particularly from the brown alga *Ascophyllum nodosum* (L.) Le Jolis [29, 30], commonly used in the agriculture industry, have been observed to be beneficial for plants because of the cell signaling activity of some molecules, such as polysaccharides [31, 32, 33], polyphenols, peptides, carotenoids [34], betaines [35], and macro- and micronutrients. Moreover, phytohormones (e.g., auxins, gibberellins and cytokinins), found in seaweed extracts, accelerate metabolism and development [36, 37], as well as other hormone-like substances [38, 39, 40]. Among the numerous commercial biostimulants, Goteo® (Goteo—Goactiv, UPL, Cesena, Italy), a biologically active filtrate, called GA142, from the seaweed *A. nodosum*, is a source of polysaccharides, vitamins, auxins and cytokinins [41, 42]. A previous study carried out by Loconsole et al, proved that Goteo® improved the aerial and root quality traits of ornamental cuttings at a dose of 3 mL L<sup>-1</sup>, in wild sage, with a greater number of roots, better growth traits, root morphologies and carbohydrate content compared to IBA. Moreover, in glossy Abelia, the same authors [43], suggested the application of a 1 mL L<sup>-1</sup> concentration of Goteo® to obtain high-quality rooted stem cuttings.

The biostimulant Kelpak® (Kelp Products Ltd., Cape Town, South Africa), an extract of brown seaweed, in particular *Ecklonia maxima* (Osbeck) Papenfuss, stimulates the growth and enlargement of the root system, due to the very high auxin-to-cytokinin ratio. This leads to a better absorption and translocation of macro- and microelements [44]. Moreover, the research carried out by Szabó et al. [45] on *Prunus* ‘Marianna’, showed that application of the Kelpak® biostimulant resulted in the highest rooting rate and increased fresh weight of cuttings during rooting compared to the control. Finally, Traversari et al. [46], observed that the application of Kelpak® improved both rooting percentage and root biometric parameters on two rose cultivar cuttings proving that a

sustainable replacement of synthetic products used for rooting promotion is possible and desirable.

Red-tip photinia (*Photinia x fraseri* Dress.), a hybrid from *P. serrulata* Lindl. x *P. glabra* Thunb. Maxim., belonging to the *Rosaceae* family and discovered at the Fraser Nursery in Birmingham (Alabama, USA) around 1940, is an evergreen shrub. It is probably the most popular hedging plant of the last 20 years with great ornamental value; it has young red foliage and heads of long-lasting creamy flowers in late spring [47], and is widely used in the design of green areas [48]. Used by landscape designers as a low maintenance hedge or screen that provides spring interest, it attracts pollinating insects with its nectar and the larvae of some species of *Lepidoptera* feed on its leaves. However, despite its important role in floriculture, this species is considered difficult to propagate [49]; although high concentrations of phytohormones are used to optimize rhizogenesis [50], limiting its commercial use [48]. *P. x Fraseri* 'Red Robin' differs from the species in that the young shoots are a particularly strong red color and it is particularly robust.

Despite many studies having illustrated the value of seaweed extracts in promoting growth, quality and yield when applied to the plant or rhizosphere in the production of cereals, fruit, vegetables and ornamental plants [51, 52, 53], their use in vegetative propagation by cutting has been little investigated in terms of the current bibliography [54 – 57]. To our knowledge, there is currently no feasible enhanced method for vegetative propagation of *P. x fraseri* 'Red Robin' through stem cuttings, by applying algae extracts as root promoters.

Given the economic importance of the species, this research aims to assess the effects of IBA and seaweed extract-based stimulators on the quality of photinia cuttings, in terms of rooting indicators and ground and aboveground agronomic features.

## 2. Materials and Methods

### 2.1. Rooting environment

The experiment was carried out from 4 March to 30 October 2021 (240 days after cutting, DAC) in a commercial propagation greenhouse located in Apulia (southern Italy) (40°54'19.1" N, 17°18'21.4" E; 66 m a.s.l.), covered with an ethylene-vinyl acetate film with a net providing 50% shading. During the experiment, the greenhouse environmental parameters were: air temperature ranging from a minimum (night time) of 12°C to a maximum (daytime) of 27°C; seedbeds (bottom heating during the winter period) at the temperature of 18 ± 1 °C; and misting for 60 s at 20 min intervals (with droplet average size of 100 µm), daily from 8 a.m. to 6 p.m.

### 2.2. Mother plants and cuttings

Twenty stock mother plants of *P. x fraseri* 'Red Robin' were randomly chosen, grown in open field conditions and regularly pruned to prevent flowering. Ninety median and semi-hardwood stem cuttings were harvested from each mother plant. Each cutting was chosen in order to ensure uniformity, vigour, absence of disease, trueness to type, and a length of 4 cm with three nodes, removing the basal leaves and maintaining two leaves per cutting.

### 2.3. Rooting promoters and cutting propagation

Cuttings were rooted using three different commercial rooting promoters: auxins were applied in the form of commercially available rooting powder: Rhizopon AA (identified as R), at 1% IBA (Sigma, St. Louis, MO, USA), directly to the bases of cuttings; the concentration was chosen according to previous studies that observed how high doses of IBA improve the quality of rooted cuttings. For example, [58] and [59], have achieved the best results with 0.8%, while Bonamino and Blazich [50] have applied IBA at 1%. Kelpak® (identified as K), is composed of 34% (w/w) *E. maxima* extract 11.16 mg L<sup>-1</sup> auxin and 0.031 mg L<sup>-1</sup> cytokinin (with a auxin:cytokinin ratio of 360:1), alginates (1.5 L<sup>-1</sup>), amino acids (total 441.3 mg 100 g<sup>-1</sup>), mannitol (2261 mg L<sup>-1</sup>), neutral sugars (1.08 g L<sup>-1</sup>),

and small amounts of macroelements (N 0.09%, P 90.7 mg kg<sup>-1</sup>, K 7163.3 mg kg<sup>-1</sup>, Ca 190.4 mg kg<sup>-1</sup>, Mg 337.2 mg kg<sup>-1</sup>, Na 1623.7 mg kg<sup>-1</sup>) and microelements (mean composition: Mn 17.3 mg kg<sup>-1</sup>, Fe 40.7 mg kg<sup>-1</sup>, Cu 13.5 mg kg<sup>-1</sup>, Zn 17.0 mg kg<sup>-1</sup>, B 33.0 mg kg<sup>-1</sup>) [60, 44]. The (producer's) recommended dose is 10mLL<sup>-1</sup> before transplanting and the same dose three times after transplanting (7 days between each treatment); the dose chosen in our experiment was 2 and 3 mLL<sup>-1</sup> respectively because they were applied by foliar application and not by drenching. Goteo® (Goteo – Goactiv, UPL, Italy) (identified as G), a liquid formulation used as a source of auxins, cytokinins, polysaccharides and vitamins. GA142 is supplemented by the company with organic mineral fertilizers (w/v): 13% P<sub>2</sub>O<sub>5</sub>, 5% K<sub>2</sub>O and 1.3-2.4% organic matter. In terms of concentration, the company recommends a 0.1% solution (1mLL<sup>-1</sup>) for vegetable crops, while ornamental plants are not specified. Previous experiments, from Gajc-Wolska [61] and Matysiak [62], suggested 3 to 4 treatments with 0.2% solution (2 mLL<sup>-1</sup>) every 2 weeks to accelerate root regeneration.

On 4 March 2021, plastic trays (100 holes, 3.5 cm in diameter) were sterilized with a fresh chlorine solution: one part bleach (5.25% sodium hypochlorite) to nine parts water for a final strength of 0.5%. They were filled with paper tubes containing the substrate, a mixture of brown and blond peat and perlite (v:v = 80:20; pH 5.0-6.0; organic carbon, 35%; organic nitrogen, 0.8%; organic matter, 85%) (Jiffy® Products International BV, Toul, France) and watered to saturation. Cuttings were inserted to the depth of 2 cm into the substrate.

#### 2.4. Experimental design

The experiment, started on 4 March, consisted of six treatments, each in three replications, each replication containing 100 cuttings.

The treatments applied were different concentrations of rooting stimulators compared to an untreated control:

- C0: distilled water;
- R1: Rhizopon AA, 1%IBA;
- K2: 2 mL L<sup>-1</sup>;
- K3: 3 mL L<sup>-1</sup>;
- G2: 2 mL L<sup>-1</sup>;
- G3: 3 mL L<sup>-1</sup>.

IBA was applied directly to the bases of cuttings (10 mm) in the form of Rhizopon AA, the commercially available rooting powder.

Starting from 4 March, the Kelpak® and Goteo® solutions were sprayed with a hand sprayer every two weeks until they ran off the leaves of the cuttings, a total of 4 times.

The treatment design was a randomized complete block design of 18 experimental units (6 concentrations x 3 replicates).

Experimental analysis were conducted in the Floriculture and ornamental plants laboratory of Department of Soil, Plant and Food Sciences, at the University of Bari.

#### 2.5. Callus initiation and rooting formation

Beginning on day 5, three cuttings were randomly taken each time to observe the changes in cutting morphological traits. At 70 DAC (13 May), callus initiation and rooting formation was investigated. The counting method was used to quantify the callus production rate and rooting percentage; gross morphological changes in the base of the stem cutting were captured using a Nikon SMZ800N microscope with a Nikon DS-Fi1 camera (Nikon Corporation, Tokyo, Japan) and Nis Elements 4.0 digital software at a resolution of 96 dpi. A cutting was considered rooted if it had at least one primary root ≥1 mm long; unrooted cutting percentage was also evaluated.

On 13 May, 12 June (at 100 DAC) and 30 October (at 240 DAC), the callused (number of cuttings that produced calli/total number of cuttings x 100%) cuttings and

rooted (number of rooted cuttings/total number of cuttings tested  $\times$  100%) cuttings percentages were recorded for twelve randomly chosen cuttings per treatment. At 240 DAC, the cutting survival percentage was also monitored. Percentage data were subjected to arcsine square root transformation before ANOVA analysis. At 240 DAC, the number of cuttings with 1, 2, 3, 4, 5 and 6 roots and the average root number per cutting for each treatment were recorded.

#### 2.6. Roots architecture

At 240 DAC, twelve rooted cuttings per treatment were sampled and analysed for the morphology of the adventitious rooting system. Each rooted cutting was washed and separated into roots and leaves plus stems. The rooting substrate was gently washed out using water and a soft brush. The roots were scanned at 400 dpi using an Epson v700 Perfection (Japan) scanner. The images were then processed using image analysis software (WinRHIZO v. 2005b©, Regent Instruments Inc., Québec, QC, Canada, [www.regentinstruments.com](http://www.regentinstruments.com) accessed on 29 December 2022) for the evaluation of total root length, root surface area, root diameter, number of root tips, forks and crossings.

#### 2.7. Ground biomass

At 240 DAC, root fresh and dry weights (g) were measured on twelve rooted cuttings per treatment: fresh samples were dried in a thermo-ventilated oven at 70 °C until it reached a constant mass.

#### 2.8. Aboveground quality features

At 240 DAC, the same samples analyzed for root parameters were recorded for aerial growth traits. Three new and fully opened leaves were sampled for analyzing the chlorophyll index (SPAD) (Konica Minolta Chlorophyll Meter SPAD-502 Plus, Solna, Sweden). The number of leaves per cutting was counted and the total leaf area per rooted cutting was measured with a leaf area meter (Delta-T; Decagon Devices, Pullman, WA, USA). Aboveground (leaves + stems) fresh and dry weights (g) were measured: samples were dried in a thermo-ventilated oven at 70 °C until it reached a constant mass.

#### 2.9. Statistical analysis

A one-way analysis of variance (ANOVA) was performed to test the effects of different concentrations of the rooting promoters on rooting performance and morphological traits. All the above data analyses were performed using SAS version 9.3 statistical software (SAS, 1999); treatment means were separated by the SNK (Student-Newman-Keuls) test ( $p \leq 0.05$ ).


### 3. Results

Table 1 shows the percentage values of callus, rooted and unrooted cuttings at 70 DAC. A higher percentage of callused cuttings was observed with both seaweed extracts (K and G) at different concentrations compared to the IBA - Rhizopon treatment (R1) and the control. The percentage of callused cutting was 84.3% on average in both the seaweed extracts, greater than 1.3 and 1.4 times compared to the control and the R treatment. The rooted cutting percentage was significantly higher in the R1 treatment (29.2%) followed by the G3 treatment (16.7%); the C0, K2, K3 and G2 treatments recorded the lowest rooted cutting percentage values, with no differences between them. The same table shows that the highest value of unrooted cuttings was found in the control (average 33.3%), while the cuttings treated with G3 all had calli or roots.

**Table 1.** Callused, rooted and unrooted cutting (%) at 70 DAC in *P. x fraseri* 'Red Robin' influenced by rooting promoters concentration (RPC).

| RPC | Cuttings (%) |              |              |
|-----|--------------|--------------|--------------|
|     | Callused     | Rooted       | Unrooted     |
| C0  | 66.7 ± 4.2 b | 0.0 ± 0.0 c  | 33.3 ± 4.2 a |
| R1  | 58.3 ± 4.0 c | 29.2 ± 3.0 a | 12.5 ± 0.0 b |
| K2  | 87.5 ± 0.0 a | 0.0 ± 0.0 c  | 12.5 ± 0.0 b |
| K3  | 83.3 ± 4.2 a | 0.0 ± 0.0 c  | 16.7 ± 4.2 b |
| G2  | 83.3 ± 4.2 a | 0.0 ± 0.0 c  | 16.7 ± 4.2 b |
| G3  | 83.3 ± 4.2 a | 16.7 ± 4.2 b | 0.0 ± 0.0 c  |

In columns, different letters indicate significant differences within parameters (S.N.K. test,  $p \leq 0.05$ ; mean  $\pm$  SD,  $n = 3$ ) C0: untreated control; R1: Rhizopon AA (1% IBA); K: Kelpak®; G: Goteo®; K2: 2 mL L<sup>-1</sup>; K3: 3 mL L<sup>-1</sup>; G2: 2 mL L<sup>-1</sup>; G3: 3 mL L<sup>-1</sup>.



**Figure 1.** On the left (a) the production of callous tissue in G2 treated cutting; on the right (b) the adventitious root in R1 treated cutting.

Figure 1 is a microscope image: we can see, on the left (a), the production of callous tissue in the G2 treated cutting; on the right (b), the adventitious root in the R1 treated cutting. Table 2 shows the callused and rooted cutting percentage at 100 and 240 DAC. At 100 DAC, the greatest callused cutting percentage values were obtained in the untreated control and Goteo® at the highest dose. The R1 application resulted in 51% fewer callused cuttings compared to the control; the same trend was found at 240 DAC. Furthermore, K3 and G3 showed a lower production of callused cuttings, compared to the corresponding lower dose of the same biostimulant (K3: 11.3%; G3: 10.3%; K2: 18.0%; G2: 15.3%). The same Table 2 shows that, at 100 and 240 DAC, the percentage of rooted cuttings treated under R1 was statistically the highest (25% at 100 DAC and 34.3% at 240 DAC), with an increase of 15.5% compared to the control at 240 DAC. At 100 DAC, treatments with biostimulants showed the lowest values of rooting compared to IBA. At 240 DAC, the worst results were obtained by both biostimulant treatments at the highest doses, with no statistically significant differences between them (K3 = 21.3% and G3 = 20.7%). At 240 DAC, cuttings treated under R1 resulted in the best survival percentage (data not shown).

**Table 2.** Callused and rooted cutting (%) at 100 and 240 DAC in *P. x fraseri* ‘Red Robin’ influenced by rooting promoters concentration (RPC).

| RPC | Cuttings (%)  |               |               |               |
|-----|---------------|---------------|---------------|---------------|
|     | Callused      |               | Rooted        |               |
|     | 100 DAC       | 240 DAC       | 100 DAC       | 240 DAC       |
| C0  | 33.0 ± 1.1 a  | 13.7 ± 0.7 bc | 7.7 ± 0.7 d   | 29.7 ± 1.4 b  |
| R1  | 16.0 ± 1.0 d  | 3.3 ± 0.3 e   | 25.0 ± 1.0 a  | 34.3 ± 0.9 a  |
| K2  | 30.0 ± 1.0 b  | 18.0 ± 1.1 a  | 12.0 ± 1.0 bc | 25.0 ± 0.6 c  |
| K3  | 29.0 ± 1.0 bc | 11.3 ± 0.7 cd | 13.3 ± 1.3 b  | 21.3 ± 0.7 d  |
| G2  | 27.3 ± 0.9 c  | 15.3 ± 0.9 b  | 14.3 ± 1.3 b  | 27.0 ± 1.0 bc |
| G3  | 34.0 ± 0.6 a  | 10.3 ± 0.9 d  | 9.0 ± 1.6 cd  | 20.7 ± 0.9 d  |

In columns, different letters indicate significant differences within parameters (S.N.K. test,  $p \leq 0.05$ ; mean ± SD, n=3) C0: untreated control; R1: Rhizopon AA (1% IBA); K: Kelpak®; G: Goteo®; K2: 2 mL L<sup>-1</sup>; K3: 3 mL L<sup>-1</sup>; G2: 2 mL L<sup>-1</sup>; G3: 3 mL L<sup>-1</sup>.

Table 3 shows, at 240 DAC, the number of rooted cuttings per treatment with a different number of roots, between 1 and 6. Treatment with IBA positively influenced the number of cuttings with 3, 4, 5 and 6 roots.

**Table 3.** Cuttings (no.) with 1, 2, 3, 4, 5 and 6 roots at 240 DAC in *P. x fraseri* ‘Red Robin’ influenced by rooting promoters concentration (RPC).


| RPC | Rooted cuttings (no.) |             |             |             |             |              |
|-----|-----------------------|-------------|-------------|-------------|-------------|--------------|
|     | 1 root                | 2 roots     | 3 roots     | 4 roots     | 5 roots     | 6 roots      |
| C0  | 13.7 ± 0.7 a          | 7.0 ± 0.6 a | 4.3 ± 0.3 b | 3.7 ± 0.3 a | 1.0 ± 0.0 b | 0.0 ± 0.0 c  |
| R1  | 7.0 ± 0.6 c           | 7.0 ± 1.0 a | 8.7 ± 0.3 a | 4.3 ± 0.3 a | 3.0 ± 0.6 a | 4.3 ± 0.3 a  |
| K2  | 10.3 ± 0.7 b          | 7.3 ± 0.3 a | 4.3 ± 0.3 b | 1.7 ± 0.3 b | 1.0 ± 0.0 b | 0.4 ± 0.3 bc |
| K3  | 9.7 ± 0.3 b           | 5.3 ± 0.3 a | 2.7 ± 0.3 c | 2.0 ± 0.0 b | 1.0 ± 0.0 b | 0.7 ± 0.3 bc |
| G2  | 14.0 ± 0.6 a          | 6.3 ± 0.3 a | 2.0 ± 0.6 c | 4.0 ± 0.6 a | 0.7 ± 0.3 b | 0.0 ± 0.0 c  |
| G3  | 10.4 ± 0.7 b          | 5.7 ± 0.7 a | 2.3 ± 0.3 c | 1.3 ± 0.3 b | 0.0 ± 0.0 b | 1.3 ± 0.3 b  |

In columns, different letters indicate significant differences within parameters (S.N.K. test,  $p \leq 0.05$ ; mean ± SD, n=3) C0: untreated control; R1: Rhizopon AA (1% IBA); K: Kelpak®; G: Goteo®; K2: 2 mL L<sup>-1</sup>; K3: 3 mL L<sup>-1</sup>; G2: 2 mL L<sup>-1</sup>; G3: 3 mL L<sup>-1</sup>.

Data on root morphological features are provided in Tables 4-5. In Table 4, at 240 DAC the statistically highest value of number of roots per cutting was obtained in the

treatment carried out with the R1 application (3.07 roots per cutting). This was 50% higher than the average of all other treatments, which do not have statistically significant differences between them. The same table shows that the greatest root length development was obtained in cuttings treated with Rhizopon, compared to the other treatments; the applications of Kelpak® and Goteo® biostimulants at both concentrations resulted in an inhibition of root length with values below the untreated control. The same trend was recorded for the surface area.

Table 5 shows the average number of tips, forks and crossings at 240 DAC. Cuttings treated with R1 showed a 105% increase in tips compared to the untreated control and G3. The K2, K3 and G2 treatments were equal to each other and below the control, by an average of 32%. Regarding the number of forks and crossings (Table 5), the highest value was obtained with cuttings treated with Rhizopon, +61% forks and +15% crossings compared to the control; the cuttings treated with the biostimulants at the different concentrations obtained lower values than the control.



**Figure 2.** Root architecture in *P. x fraseri* ‘Red Robin’ influenced by application of rooting promoters concentration.

**Table 4.** Root morphological traits: roots number (no.), length (mm), surface area (mm<sup>2</sup>) and average diameter (mm), at 240 DAC in *P. x fraseri* influenced by rooting promoters concentration (RPC).

| RPC | Roots (no.)  | Length (mm)   | Surface area (mm <sup>2</sup> ) | Diameter (mm) |
|-----|--------------|---------------|---------------------------------|---------------|
| C0  | 2.0 ± 0.04 b | 133.0 ± 1.1 b | 23.8 ± 0.9 b                    | 0.60 ± 0.01 a |

|    |              |               |              |               |
|----|--------------|---------------|--------------|---------------|
| R1 | 3.1 ± 0.07 a | 162.1 ± 7.1 a | 33.9 ± 0.9 a | 0.73 ± 0.02 a |
| K2 | 2.1 ± 0.10 b | 54.9 ± 3.0 cd | 12.5 ± 0.4 c | 0.67 ± 0.02 a |
| K3 | 2.1 ± 0.05 b | 51.1 ± 2.3 d  | 11.8 ± 0.6 c | 0.66 ± 0.01 a |
| G2 | 1.9 ± 0.08 b | 67.6 ± 4.0 cd | 14.5 ± 0.1 c | 0.70 ± 0.02 a |
| G3 | 2.0 ± 0.07 b | 72.2 ± 7.0 c  | 16.3 ± 0.2 c | 0.73 ± 0.01 a |

In columns, different letters indicate significant differences within parameters (S.N.K. test,  $p \leq 0.05$ ; mean ± SD, n=3) C0: untreated control; R1: Rhizopon AA (1% IBA); K: Kelpak®; G: Goteo®; K2: 2 mL L<sup>-1</sup>; K3: 3 mL L<sup>-1</sup>; G2: 2 mL L<sup>-1</sup>; G3: 3 mL L<sup>-1</sup>.

**Table 5.** Root morphological traits: root tips (no.), forks (no.) and crossing (no.), at 240 DAC in *P. x fraseri* influenced by rooting promoters concentration (RPC).

| RPC | Root        |              |                 |
|-----|-------------|--------------|-----------------|
|     | Tips (no.)  | Forks (no.)  | Crossings (no.) |
| C0  | 330 ± 7.0 b | 755 ± 12.5 b | 99 ± 1.8 b      |
| R1  | 677 ± 5.6 a | 1215 ± 7.7 a | 114 ± 3.8 a     |
| K2  | 217 ± 1.1 c | 326 ± 6.3 e  | 25 ± 1.1 e      |
| K3  | 225 ± 6.7 c | 298 ± 5.5 e  | 30 ± 1.5 e      |
| G2  | 226 ± 3.8 c | 459 ± 7.9 d  | 38 ± 0.6 d      |
| G3  | 340 ± 8.7 b | 621 ± 13.3 c | 49 ± 1.7 c      |

In columns, different letters indicate significant differences within parameters (S.N.K. test,  $p \leq 0.05$ ; mean ± SD, n=3) C0: untreated control; R1: Rhizopon AA (1% IBA); K: Kelpak®; G: Goteo®; K2: 2 mL L<sup>-1</sup>; K3: 3 mL L<sup>-1</sup>; G2: 2 mL L<sup>-1</sup>; G3: 3 mL L<sup>-1</sup>.

Data on fresh and dry ground weights are provided in Table 6: rooted cuttings under R1 show the highest, and statistically different, dry value (0.35g) compared to the other treatments, resulting in +40% in comparison to the control.

**Table 6.** Ground fresh and dry weights (g) per cutting, at 240 DAC in *P. x fraseri* influenced by rooting promoters concentration (RPC).

| RPC | Ground weights (g) |               |
|-----|--------------------|---------------|
|     | Fresh              | Dry           |
| C0  | 0.81 ± 0.07 ab     | 0.25 ± 0.09 b |
| R1  | 0.97 ± 0.02 a      | 0.35 ± 0.01 a |
| K2  | 0.47 ± 0.02 c      | 0.15 ± 0.01 c |

|    |                |               |
|----|----------------|---------------|
| K3 | 0.72 ± 0.02 b  | 0.23 ± 0.01 b |
| G2 | 0.81 ± 0.07 ab | 0.24 ± 0.01 b |
| G3 | 0.65 ± 0.03 b  | 0.18 ± 0.01 c |

In columns, different letters indicate significant differences within parameters (S.N.K. test,  $p \leq 0.05$ ; mean  $\pm$  SD,  $n = 3$ ) C0: untreated control; R1: Rhizopon AA (1% IBA); K: Kelpak®; G: Goteo®; K2: 2 mL L<sup>-1</sup>; K3: 3 mL L<sup>-1</sup>; G2: 2 mL L<sup>-1</sup>; G3: 3 mL L<sup>-1</sup>.

Table 7 shows the aboveground morpho-biometric features: no statistically significant differences were among the treatments, in terms of number of leaves per cutting and chlorophyll index. At 240 DAC, the statistically highest value of leaf area per cutting was obtained in the treatment carried out with the R1 application (27.9 cm<sup>2</sup>); K3 and G3 were similar to the control, with lower values compared to R1. The worst performances were observed in the K2 and G2 treatments.

**Table 7.** Above-ground morpho-biometric traits: leaves per cutting (no.), leaf area (mm<sup>2</sup>) and chlorophyll index (SPAD), at 240 DAC in *P. x fraseri* influenced by rooting promoters concentration (RPC).

| RPC | Leaves per cutting (no.) | Chlorophyll index (SPAD) | Leaf area per cutting (cm <sup>2</sup> ) |
|-----|--------------------------|--------------------------|------------------------------------------|
| C0  | 3.7 ± 0.7 a              | 570 ± 13.7 a             | 22.0 ± 1.2 b                             |
| R1  | 4.7 ± 0.3 a              | 530 ± 11.9 a             | 27.9 ± 0.8 a                             |
| K2  | 3.3 ± 0.3 a              | 533 ± 15.0 a             | 18.1 ± 0.3 c                             |
| K3  | 4.0 ± 0.6 a              | 554 ± 2.91 a             | 21.9 ± 0.9 b                             |
| G2  | 3.3 ± 0.3 a              | 529 ± 20.4 a             | 18.4 ± 0.4 c                             |
| G3  | 4.00 ± 0.6 a             | 582 ± 11.6 a             | 22.3 ± 1.4 b                             |

In columns, different letters indicate significant differences within parameters (S.N.K. test,  $p \leq 0.05$ ; mean  $\pm$  SD,  $n = 3$ ) C0: untreated control; R1: Rhizopon AA (1% IBA); K: Kelpak®; G: Goteo®; K2: 2 mL L<sup>-1</sup>; K3: 3 mL L<sup>-1</sup>; G2: 2 mL L<sup>-1</sup>; G3: 3 mL L<sup>-1</sup>.

Table 8 showed that the aboveground dry weight was significantly influenced by treatments: the rooting effect under IBA (R1) was higher than that under the control and K3.

**Table 8.** Aboveground fresh and dry weights (g) per cutting, at 240 DAC in *P. x fraseri* influenced by rooting promoters concentration (RPC).

| RPC | Above-ground weights (g) |               |
|-----|--------------------------|---------------|
|     | Fresh                    | Dry           |
| C0  | 1.11 ± 0.14 a            | 0.40 ± 0.09b  |
| R1  | 1.29 ± 0.13 a            | 0.47 ± 0.11 a |

|    |               |                |
|----|---------------|----------------|
| K2 | 0.99 ± 0.11 a | 0.35 ± 0.04 bc |
| K3 | 1.11 ± 0.09 a | 0.39 ± 0.08 b  |
| G2 | 0.96 ± 0.08 a | 0.33 ± 0.08 c  |
| G3 | 1.04 ± 0.06 a | 0.35 ± 0.03 bc |

In columns, different letters indicate significant differences within parameters (S.N.K. test,  $p \leq 0.05$ ; mean  $\pm$  SD,  $n = 3$ ) C0: untreated control; R1: Rhizopon AA (1% IBA); K: Kelpak®; G: Goteo®; K2: 2 mL L<sup>-1</sup>; K3: 3 mL L<sup>-1</sup>; G2: 2 mL L<sup>-1</sup>; G3: 3 mL L<sup>-1</sup>.

#### 4. Discussion

Calkins [63] reported that Fraser's photinia is an important primary ornamental landscaping species with optimal hedge or screen effects and low maintenance, but it is difficult to root when propagated by cuttings. Given the economic importance of the species in the ornamental industry, this research aims to assess the effects of IBA and algae extract-based stimulators on the quality of photinia cuttings, in terms of rooting indicators and ground and aboveground agronomic features. Vegetative propagation by cuttings has numerous advantages, one of them is to provide more uniform and agronomically superior commercial plantlets than those obtained by heterozygous seeds [64]. First results show different responses to adventitious rhizogenesis under in-dole-3-butyric acid and seaweed extract treatments. The application of exogenous auxin and cytokinin stimulates callus differentiation in various species [65]. Several studies showed that an intermediate ratio of auxin and cytokinin promotes callus induction, while a high ratio of auxin-to-cytokinin or cytokinin-to-auxin induces root and shoot regeneration, respectively [66]. Plants of *Arabidopsis thaliana* treated with *A. nodosum* extract biostimulant, showed an increased cytokinin-like responses, suggesting a contribution in the cytokinin-like activity of the extracts's compounds [67]. From our results obtained at 70 DAC, it would seem that the seaweed extracts stimulated the production of over 80% of cuttings with callus (Figure 1, Table 1). Comparing this result to IBA (R1), it can be seen that this, in contrast, obtained the highest rooting percentage (Table 1). Furthermore, it would appear that the conspicuous production of callus tissue prevents the adventitious rooting of photinia cuttings, representing a structural obstacle to the emission of adventitious roots. Our preliminary results agree with Monder et al. [68]: in the rhizogenesis of 'Hurdal' rose, the authors showed that an increase in rooting percentage was only strictly connected to a decrease in the percentage of cuttings with calli only. Callus overgrowth in photinia could be an unfavorable phenomenon for fast rhizogenesis. Conversely, Costa et al. [69] in *Rosa 'Madelon'* and Fouda and Schmidt [70] in *Rosa rugosa* stem cuttings found that the new parenchyma tissue (callus) precedes root initiation. Martins et al. [71] showed an inverse relationship between rooting and callus formation for olive tree cuttings. In our study, the application of IBA (R1) improved the rooting percentage at 240 DAC (Table 2) and the number of cuttings with six roots (Table 3) as compared to the untreated control and seaweed extracts. In our experimental conditions, the application of IBA at the concentration of 1g L<sup>-1</sup> produced 33% of rooted cuttings, while Bonaminio and Blazich, [50] in 1983, found that 5000 and 10000 mg L<sup>-1</sup> of IBA solutions applied to the terminal, semi-hardwood cuttings of Fraser's photinia, promoted rooting more effectively than the control, and increased rooting percentage significantly (100 and 93%). Cutting success, entailing quality AR formation with high rooting percentage, depends on numerous factors, such as cutting type, environmental conditions, nutritional levels of the stock plant, rooting medium and phytohormone application [72, 73, 74]. IBA has been reported to increase *in vivo* adventitious root formation, overall quality and uniformity of roots in many ornamental species [75 - 84]. Untreated stem

cuttings (C0) were also able to root, but with a lower rooting percentage, in comparison to IBA-treated cuttings, possibly due to the presence of stored carbohydrates and endogenous auxin contents in the cuttings [85]. Until now, no research has compared the morphological quality of adventitious roots treated with different IBA and seaweed extract concentrations in Fraser's Photinia cuttings. To improve plant performance and provide protection against the deleterious effects of numerous abiotic stressors, several amendments such as biostimulators and bioelicitors have been used [86]. The favourable impacts of using seaweed extract as natural regulators have resulted in better crop growth and production [87, 88]. In our experimental conditions, neither K nor G biostimulants, at different concentrations, increased the rooting percentage (Tables 2) and the root architecture (Figure 2, Tables 4-5) compared to the IBA treatment. On the contrary, previous studies have shown the efficacy of biostimulants in promoting rhizogenesis. For example, the use of a 40% concentrated *A. nodosum* extract increased the rooting of *Passiflora actinia* by about 10% [89]. Even if unsuccessful, our results in the treatments with biostimulants agree with those of Traversari et al. [46] on rose rhizogenesis: cuttings treated with Phylgreen, a commercial biostimulant made from *A. nodosum*, through a low temperature mechanical extraction, had low values of both survival rate and root biometric parameters. Since biostimulating effects are clearly species-specific and product-specific, results regarding one biostimulant or one species only do not directly apply to another biostimulant or another plant species [90, 91, 92]. In our study, both Kelpak® and Goteo® negatively affected root length, surface area and diameter compared to IBA and the control (Table 4). On the contrary, positive effects of Goteo on rooting were reported for *Physocarpus opulifolius* [77], *Hydrangea paniculata* [93], *Ornithogalum arabicum* [94] and rose [59]. In Fraser's Photinia, both Kelpak® and Goteo® at different concentrations decreased the number of root tips, forks and crossings compared to the IBA treatment (Table 5). Based on these findings, the overall development and morphology of Photinia cuttings treated with Kelpak® and Goteo® were inhibited.

Seaweed extracts often stimulate and accelerate cell division, elongation, differentiation and protein synthesis [38, 95]; Makhaye et al. [96] have verified the potential stimulatory effect of biostimulants (especially Kelpak®) on the germination of *A. esculentus* seeds. The application of seaweed extracts based biostimulants in our experiment did not positively influence the number of leaves, chlorophyll index (SPAD) and leaf area index (Table 7). These results are in disagreement with that found in *Lantana camara*, *Abelia x grandiflora* [43] and *Cornus alba* 'Aurea' and 'Elegantissima' [97]. Our preliminary results (Tables 6, 8) disagree with Ratore et al. [98], Kocira et al. [99], Gajc-Wolska et al. [61] and Caccialupi et al. [100] who obtained a positive effect, by applying seaweed extracts, on plant growth, development and yield. Our results agree with Francke et al. [42] who showed a lower yield of shallots than that of the control (4%) by applying Goteo biostimulant. The limited information available in the literature does not allow further discussion of this biostimulant since contrasting results were observed in Fraser's photinia 'Red Robin' cuttings with respect to other investigated species. The overall quality of cuttings in the IBA treatment was the strongest, with 1000 mg L<sup>-1</sup>, being the optimum concentration, according to studies by Quan et al. [101].

## 5. Conclusion

In conclusion, the main outcomes of this study can be summarized as follows: we suggest that the use of 1% IBA (Rhizopon AA), compared to an untreated control, may be beneficial to ornamental nursery farmers wishing to produce Fraser's photinia 'Red Robin' quality cuttings with a well-developed root system and, therefore, capable of achieving rapid establishment at transplantation. On the contrary, neither seaweed extract, Kelpak® or Goteo®, at different concentrations, improved root percentage and architecture, and ground and above ground weights of rooted cutting, compared to the

untreated control. This result indicates that these natural products are not suitable for Fraser's photinia 'Red Robin' propagation using this methodology. Further research must be conducted to propose effective agronomic protocols by investigating: application methods, doses and number of applications and to clarify the biochemical and molecular mechanisms of action of these seaweed extracts.

**Author Contributions:** Conceptualization, B.D.L. and G.C.; methodology, B.D.L., G.C. and D.L.; software, G.C.; data curation, B.D.L., G.C. and D.L.; writing—original draft preparation, B.D.L., G.C., D.L. and A.E.S.; writing—review and editing, B.D.L., G.C., D.L. and A.E.S.; funding acquisition, B.D.L. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by Regione Puglia (Italy) MIS. 16 – Cooperazione – sottomisura 16.2 – Sostegno a progetti pilota e allo sviluppo di nuovi prodotti, pratiche, processi e tecnologie. Grant number B79J20000140009. Agreement "Transfer of protocols for quarantine and harmful organisms and for the selection of sanitary materials improved for the Apulian nursery- ProDi-QuaVi" project ("Trasferimento di protocolli per organismi da quarantena e nocivi e per la selezione di materiali sanitariamente migliorati per il vivaismo pugliese (B79J20000140009)" realizzato nell'ambito del PSR 2014-2020 – Regione Puglia MIS. 16 – Cooperazione – sottomisura 16.2 – Sostegno a progetti pilota e allo sviluppo di nuovi prodotti, pratiche, processi e tecnologie – Avviso pubblico approvato con DAG n. 194 del 12/09/2018, in Italian). Paper no. 3.

**Acknowledgments:** The authors thank Vivai Capitanio (Monopoli, BA, Italy) and Giovanni Fanizzi for technical support in the field and the materials used in the experiments. The authors thanks Claudia Ruta and Patrick Angelo Guarini for providing huge help in microscope image acquisition.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

- Del Buono, D. Can biostimulants be used to mitigate the effect of anthropogenic climate change on agriculture? It is time to respond. *Sci. Total Environ.* **2020**, *751*, 141763. <https://doi.org/10.1016/j.scitotenv.2020.141763>.
- Davydov, R.; Sokolov, M.; Hogland, W.; Glinushkin, A.; Markaryan, A. The application of pesticides and mineral fertilizers in agriculture. *MATEC Web of Conferences*. **245**, EDP Sciences, **2018** p. 11003. <https://doi.org/10.1051/mateconf/201824511003>.
- Costa, J.A.V.; Freitas, B.C.B.; Cruz, C.G.; Silveira, J.; Morais, M.G. Potential of micro algae as biopesticides to contribute to sustainable agriculture and environmental development. *J. Environ. Sci. Health B.* **2019**, *54*(5), 366–375. <https://doi.org/10.1080/03601234.2019.1571366>.
- Singh, N.; Joshi, E.; Sasode, D.S.; Dangi, R.S.; Chouhan, N. Soil fertility, macro and micro nutrient uptake and their use efficiencies under integrated nutrient management in groundnut (*Arachis hypogaea* L.). *Int. J. Chem. Stud.* **2020**, *8*(1), 1983–1987. <https://doi.org/10.22271/chemi.2020.v8.i1ac.8556>.
- Neely, C.; Bourne, M.; Chesterman, S.; Kouplevatskaya-Buttoud, I.; Bojic, D.; Vallée, D. *Implementing Agenda 2030 in Food and Agriculture: accelerating policy impact through cross-sectoral coordination at the country level*; Food and Agricultural Organization of The United Nations: Rome, Italy, 2017.
- Blakesley, D.; Chaldecott, M.A. The role of endogenous auxin in root initiation: part II. Sensitivity and evidence from studies on transgenic plant tissues. *Plant Growth Regulation* **1993**, *13*, 77–84. <https://doi.org/10.1007/BF00207595>.
- Gidoin, Y. Innovation to quality in the horticulture industry: feedback from a CEO. *Acta Hort.* **2019**, *1245*, 1–10. <https://doi.org/10.17660/ActaHortic.2019.1245.1>.
- Song, S.J.; Ko, C.H.; Shin, U.S.; Oh, H.J.; Kim, S.Y.; Lee, S.Y. Successful stem cutting propagation of *Patrinia rupestris* for horticulture. *Rhizosphere* **2019**, *9*, 90–92. <https://doi.org/10.1016/j.rhisph.2018.12.001>.
- Gallegos-Cedillo, V.M.; Diáñez, F.; Nájera, C.; Santos, M. Plant Agronomic Features Can Predict Quality and Field Performance: A Bibliometric Analysis. *Agronomy* **2021**, *11*, 2305. <https://doi.org/10.3390/agronomy11112305>.
- Loconsole, D.; Cristiano, G.; De Lucia, B. Image Analysis of Adventitious Root Quality in Wild Sage and Glossy Abelia Cuttings after Application of Different Indole-3-Butyric Acid Concentrations. *Plants* **2022**, *11*, 290. <https://doi.org/10.3390/plants11030290>.

11. Hąc-Wydro, K.; Flasiński, M. The studies on the toxicity mechanism of environmentally hazardous natural (IAA) and synthetic (NAA) auxin—the experiments on model *Arabidopsis thaliana* and rat liver plasma membranes. *Colloids and Surfaces B: Biointerfaces* **2015**, *130*, 53–60. <https://doi.org/10.1016/j.colsurfb.2015.03.064>.
12. Gonin, M.; Bergougnoux, V.; Nguyen, T.D.; Gantet, P.; Champion, A. What Makes Adventitious Roots? *Plants* **2019**, *8*, 240. <https://doi.org/10.3390/plants8070240>.
13. Abu-Zahra, T. R.; Al-Shadaideh, A. N.; Abubaker, S. M.; Qrunfleh, I. M. Influence of auxin concentrations on different ornamental plants rooting. *Int. J. Bot.* **2013**, *9*(2), 96–99. <https://doi.org/10.3923/ijb.2013.96.99>.
14. Zheng, L.; Xiao, Z.B.; Song, W.T. Effects of substrate and exogenous auxin on the adventitious rooting of *Dianthus caryophyllus* L. *Hortic. Sci.* **2020**, *55*, 170–173. <https://doi.org/10.21273/HORTSCI14334-19>.
15. Babaie, H.; Zarei, H.; Nikdel, K.; Firoozjani, M.N. Effect of different concentrations of IBA and time of taking cutting on rooting, growth and survival of *Ficus binnendijkii* ‘Amstel Queen’ cuttings. *Not. Sci. Biol.* **2014**, *6*, 163–166. <https://doi.org/10.15835/nsb629281>.
16. Bryant, P.H.; Trueman, S.J. Stem anatomy and adventitious root formation in cuttings of *Angophora*, *Corymbia* and *Eucalyptus*. *Forests* **2015**, *6*, 1227–1238. <https://doi.org/10.3390/f6041227>.
17. Lei, C.; Fan, S.; Li, K.; Meng, Y.; Mao, J.; Han, M.; Zhao, C.; Bao, L.; Zhang, D. iTRAQ-based proteomic analysis reveals potential regulation networks of IBA-induced adventitious root formation in apple. *Int. J. Mol. Sci.* **2018**, *19*, 667. <https://doi.org/10.3390/ijms19030667>.
18. Hartmann, H.T.; Kester, D.E.; Davies, F.T.; Geneve, R.L. *Plant Propagation: Principles and Practices*; Prentice Hall: Hoboken, NJ, USA, 2002.
19. Erçili, S.; Eşitken, A.; Anapali, O.; Şahin, U. Effects of substrate and iba-concentration on adventitious root formation on hardwood cuttings of *rosa dumalis*. *Acta Hortic.* **2005**, *690*, 149–152. [10.17660/ActaHortic.2005.690.22](https://doi.org/10.17660/ActaHortic.2005.690.22).
20. Ribeiro, M.M.; Collado, L.M.; Antunes, M.A. The influence of indole-3-butyric-acid in *Prunus laurocerasus* vegetative propagation. *Acta Hortic.* **2008**, *885*, 277–283. <https://doi.org/10.17660/ActaHortic.2010.885.38>.
21. Grigoriadou, K.; Sarropoulou, V.; Krigas, N.; Maloupa, E. Vegetative and in vitro propagation of the medicinal and ornamental plant *Astragalus suberosus* subsp. *Haarbachii* (Fabaceae). *Eur. J. Hortic. Sci.* **2022**, *87*, 1–9. <https://doi.org/10.17660/eJHS.2022/017>.
22. Cano, A.; Sánchez-García, A.B.; Albacete, A.; González-Bayón, R.; Justamante, M.S.; Ibáñez, S.; Pérez-Pérez, J.M. Enhanced conjugation of auxin by GH3 enzymes leads to poor adventitious rooting in carnation stem cuttings. *Front. Plant Sci.* **2018**, *9*, 566. <https://doi.org/10.3389/fpls.2018.00566>.
23. Shiri, M.; Mudywa, R.M.; Takawira, M.; Musara, C.; Gama, T. Effects of rooting media and indole-3-butyric acid (IBA) concentration on rooting and shoot development of *Duranta erecta* tip cuttings. *Afr. J. Plant Sci.* **2019**, *13*, 279–285. <https://doi.org/10.5897/AJPS2019.1851>.
24. Bulgari, R.; Franzoni, G.; Ferrante, A. Biostimulants application in horticultural crops under abiotic stress conditions. *Agronomy* **2019**, *9* (6), 306. <https://doi.org/10.3390/agronomy9060306>.
25. Cuadrado, C.J.L.; Pinillos, E.O.; Tito, R.; Mirones, C.S.; GamarraMendoza, N.N. Insecticidal properties of capsaicinoids and glucosinolates extracted from *Capsicum chinense* and *Tropaeolum tuberosum*. *Insects* **2019**, *10*(5), 132. <https://doi.org/10.3390/insects10050132>.
26. Campobenedetto, C.; Mannino, G.; Beekwilder, J.; Contartese, V.; Karlova, R.; Berteau, C. M. The application of a biostimulant based on tannins affects root architecture and improves tolerance to salinity in tomato plants. *Scientific reports* **2021**, *11*(1), 1–15. <https://doi.org/10.1038/s41598-020-79770-5>.
27. EU. Regulation of the European Parliament and of the Council Laying Down Rules on the Making Available on the Market of EU Fertilising Products and Amending Regulations (EC) No 1069/2009 and (EC) No 1107/2009 and Repealing Regulation (EC) No 2003/2003. *Off. J. Eur. Union* **2019**, *62*, 1–114. Available online: <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=OJ:L:2019:170:TOC> (accessed on 3 June 2022).
28. Ali, O.; Ramsubhag, A.; Jayaraman, J. Biostimulant properties of seaweed extracts in plants: Implications towards sustainable crop production. *Plants* **2021**, *10*, 531. <https://doi.org/10.3390/plants10030531>.
29. Ertani, A.; Francioso, O.; Tinti, A.; Schiavon, M.; Pizzeghello, D.; Nardi, S. Evaluation of seaweed extracts from *Laminaria* and *Ascomyllum nodosum* spp. As biostimulants in *Zea mays* L. using a combination of chemical, biochemical and morphological approaches. *Front. Plant Sci.* **2018**, *9*, 428. <https://doi.org/10.3389/fpls.2018.00428>.
30. Shukla, P.S.; Mantin, E.G.; Adil, M.; Bajpai, S.; Critchley, A.T.; Prithiviraj, B. *Ascomyllum nodosum*-based biostimulants: Sustainable applications in agriculture for the stimulation of plant growth, stress tolerance, and disease management. *Front. Plant Sci.* **2019**, *10*, 655. <https://doi.org/10.3389/fpls.2019.00655>.
31. Afonso, N.C.; Catarino, M.D.; Silva, A.; Cardoso, S.M. Brown macroalgae as valuable food ingredients. *Antioxidants* **2019**, *8*, 365. <https://doi.org/10.3390/antiox8090365>.
32. Bulgari, R.; Franzoni, G.; Ferrante, A. Biostimulants Application in Horticultural Crops under Abiotic Stress Conditions. *Agronomy* **2019**, *9*, 306. <https://doi.org/10.3390/agronomy9060306>.

33. Franzoni, G.; Cocetta, G.; Prinsi, B.; Ferrante, A.; Espen, L. Biostimulants on Crops: Their Impact under Abiotic Stress Conditions. *Horticulturae* **2022**, *8*, 189. <https://doi.org/10.3390/horticulturae8030189>.
34. Hrólfssdóttir, A.P.; Arason, S.; Sveinsdóttir, H.I.; Gudjónsdóttir, M. Added Value of *Ascophyllum nodosum* Side Stream Utilization during Seaweed Meal Processing. *Mar. Drugs* **2022**, *20*, 340. <https://doi.org/10.3390/md20060340>.
35. Battacharyya, D.; Babgohari, M. Z.; Rathor, P.; Prithiviraj, B. Seaweed extracts as biostimulants in horticulture. *Scientia Horticulturae* **2015**, *196*, 39–48 <https://doi.org/10.1016/j.scienta.2015.09.012>.
36. Zhang, X.; Ervin, E.H. Impact of seaweed extract-based cytokinins and zeatin riboside on creeping bent grass heat tolerance. *Crop Sci.* **2008**, *48*, 364–370. <https://doi.org/10.2135/cropsci2007.05.0262>.
37. Wang, Y.; Fu, F.; Li, J.; Wang, G.; Wu, M.; Zhan, J.; Chen, X.; Mao, Z. Effects of seaweed fertilizer on the growth of *Malus hupehensis* Rehd. Seedlings, soil enzyme activities and fungal communities under replant condition. *Eur. J. Soil Biol.* **2016**, *75*, 1–7. <https://doi.org/10.1016/j.ejsobi.2016.04.003>.
38. Stirk, W.A.; Rengasamy, K.R.R.; Kulkarni, M.G.; van Staden, J. *Plant Biostimulants from Seaweed*. In The Chemical Biology of Plant Biostimulants; John Wiley & Sons, Ltd.: Hoboken, NJ, USA, 2020; pp. 31–55. ISBN 978-1-119-35725-4.
39. Kulkarni, M.G.; Rengasamy, K.R.R.; Pendota, S.C.; Gruz, J.; Plačková, L.; Novák, O.; Doležal, K.; Van Staden, J. Bioactive molecules derived from smoke and seaweed *Ecklonia maxima* showing phytohormone-like activity in *Spinacia oleracea* L. *New Biotechnol.* **2019**, *48*, 83–89. <https://doi.org/10.1016/j.nbt.2018.08.004>.
40. Petropoulos, S.A. Practical Applications of Plant Biostimulants in Greenhouse Vegetable Crop Production. *Agronomy* **2020**, *10*, 1569. <https://doi.org/10.3390/agronomy10101569>.
41. Stepowska, A. Effects of GA 142 (Goëmar Goteo) and GA 14 (Goëmar BM86) extracts on sweet pepper yield in non-heated tunnels. In *Solanaceous Crops*; Zbigniew, T.D., Ed.; Editorial House Wie's Jutra: Warsaw, Poland, 2008; p. 45.
42. Francke, A.; Majkowska-Gadomska, J.; Kaliniewicz, Z.; Jadwisieńczyk, K. No Effect of Biostimulants on the Growth, Yield and Nutritional Value of Shallots Grown for Bunch Harvest. *Agronomy* **2022**, *12*, 1156. <https://doi.org/10.3390/agronomy12051156>.
43. Loconsole, D.; Cristiano, G.; De Lucia, B. Improving Aerial and Root Quality Traits of Two Landscaping Shrubs Stem Cuttings by Applying a Commercial Brown Seaweed Extract. *Horticulturae* **2022**, *8*, 806. <https://doi.org/10.3390/horticulturae8090806>.
44. Lötze, E.; Hoffman, E.W. Nutrient composition and content of various biological active compounds of three south African-based commercial seaweed biostimulants. *J. Appl. Phycol.* **2016**, *28*, 1379–1386. <https://doi.org/10.1007/s10811-015-0644-z>.
45. Szabó, V.; Sárvari, A.; Hrotkó, K. Treatment of stockplants with biostimulators and their effects on cutting propagation of *Prunus mariantha* 'GF 8-1'. *Acta horticulturae* **2011**, *923*, 277–281.
46. Traversari, S.; Cacini, S.; Nesi, B. Seaweed Extracts as Substitutes of Synthetic Hormones for Rooting Promotion in Rose Cuttings. *Horticulturae* **2022**, *8*, 561. <https://doi.org/10.3390/horticulturae8070561>.
47. Gilman, G.F. and Watson, D.G. 1994. Photinia × fraseri – Fraser Photinia. Fact Sheet ST-447, <http://hort.ufl.edu/trees/PHOFRAA.pdf>.
48. Larraburu, E.E.; Apóstolo, N.M.; Llorente, B.E. Anatomy and morphology of photinia (*Photinia × fraseri* Dress) in vitro plants inoculated with rhizobacteria. *Trees* **2010**, *24*, 635–642. <https://doi.org/10.1007/s00468-010-0433-x>.
49. Dirr, M. A. Tolerance of Leyland Cypress and Red-Tip Photinia to Soil and Foliar Applied Sodium Chloride. *J. Environ. Hortic.* **1990**, *8*(3), 154–155. <https://doi.org/10.24266/0738-2898-8.3.154>.
50. Bonaminio, V.P.; Blazich, F.A. Response of Fraser's photinia stem cuttings to selected rooting compounds. *J Environ Hortic* **1983**, *1*:9–11 <https://doi.org/10.1007/s00468-010-0433-x>.
51. Khan, W.; Rayirath, U.P.; Subramanian, S.; Jithesh, M.N.; Rayorath, P.; Hodges, D.M. Seaweed extracts as biostimulants of plant growth and development. *J. Plant Growth Regul.* **2009**, *28*, 386–399. <https://doi.org/10.1007/s00344-009-9103-x>.
52. Paradiković, N.; Teklić, T.; Zeljković, S.; Lisjak, M.; Špoljarević, M. Biostimulants research in some horticultural plant species—A review. *Food Energy Secur.* **2019**, *8*, e00162. <https://doi.org/10.1002/fes3.162>.
53. Kisvarga, S.; Farkas, D.; Boronkay, G.; Neményi, A.; Orlóci, L. Effects of Biostimulants in Horticulture, with Emphasis on Ornamental Plant. *Agronomy* **2022**, *12*, 1043. <https://doi.org/10.3390/agronomy12051043>.
54. Cardoso, J.C.; Vendrame, W.A. Innovation in Propagation and Cultivation of Ornamental Plants. *Horticulturae* **2022**, *8*, 229. <https://doi.org/10.3390/horticulturae8030229>.
55. Monder, M.J.; Kozakiewicz, P.; Jankowska, A. Anatomical structure changes in stem cuttings of rambler roses induced with plant origin preparations. *Sci. Hortic.* **2019**, *255*, 242–254. <https://doi.org/10.1016/j.scienta.2019.05.034>.
56. Pacholczak, A.; Nowakowska, K.; Mika, N.; Borkowska, M. The effect of the biostimulator Goteo on the rooting of nine-bark stem cuttings. *Folia Hortic.* **2016**, *28*, 109–116. <https://doi.org/10.1515/fhort-2016-0013>.
57. Pacholczak, A.; Nowakowska, K. The Effect of Biostimulators and Indole-3-Butyric Acid on Rooting of Stem Cuttings of Two Ground Cover Roses. *Acta Agrobot.* **2020**, *73*, 1. <https://doi.org/10.5586/aa.731>.
58. Hammo, Y. H.; Kareem, B. A.; Salih, M. I. Effect of planting date and IBA concentration on rooting ability of stem cutting of Fraser's Photinia (*Photinia x fraseri*). *IOSR J. of Agr. and Vet. Sci.* **2013**, *5*(4), 51–54.

59. Hasan, N. S.; Hammo, Y. H. Influence of IBA and media on rooting percentage and growth of semi-hardwood cuttings of red-tip Photinia plant (*Photinia x fraseri*). *J. of Duhok University* **2021**, *24*(1), 114–119.
60. Stirk, W. A.; Tarkowská, D.; Turečová, V.; Strnad, M.; Van Staden, J. Absciscic acid, gibberellins and brassinosteroids in Kelpak®, a commercial seaweed extract made from *Ecklonia maxima*. *J. Appl. Phycol.* **2014**, *26*(1), 561–567. <https://doi.org/10.1007/s10811-013-0062-z>.
61. Gajc-Wolska, J.; Kowalczyk, K.; Nowecka, M.; Mazur, K.; Metera, A. Effect of organic-mineral fertilizers on the yield and quality of endive (*Cichorium endivia* L.). *Acta Sci. Pol. Hortorum Cultus* **2012**, *11*, 189–200.
62. Matysiak, K.; Kaczmarek, S.; Kierzek, R.; Kardasz, P. Effect of seaweeds extracts and humic and fulvic acids on the germination and early growth of winter oilseed rape (*Brassica napus* L.). *J. Agric. Eng. Res.* **2010**, *55*, 28–32.
63. Calkins, M. *The sustainable sites handbook: A complete guide to the principles, strategies, and best practices for sustainable landscapes*. John Wiley & Sons: Hoboken, New Jersey, 2012, Volume 39.
64. Lustosa Sobrinho, R.; Zoz, T.; Finato, T.; Oliveira, C.E.d.S.; Neto, S.S.d.O.; Zoz, A.; Alaraidh, I.A.; Okla, M.K.; Alwasel, Y.A.; Beemster, G.; AbdElgawad, H. *Jatropha curcas* L. as a Plant Model for Studies on Vegetative Propagation of Native Forest Plants. *Plants* **2022**, *11*, 2457. <https://doi.org/10.3390/plants11192457>.
65. Ikeuchi, M.; Sugimoto, K.; Iwase, A.; Plant Callus: Mechanisms of Induction and Repression. *The Plant Cell* **2013**, *25*, 9, 3159–3173, <https://doi.org/10.1105/tpc.113.116053>
66. Skoog, F.; Miller, C.O. Chemical regulation of growth and organ formation in plant tissues cultured in vitro. *Symp. Soc. Exp. Biol.* **1957**, *11*, 118–130.
67. Khan, W.; Hiltz, D.; Critchley, A.T.; Prithiviraj, B. Bioassay to detect *Ascophyllum nodosum* extract-induced cytokinin-like activity in *Arabidopsis thaliana*. *J Appl Phycol* **2011**, *23*, 409–414. <https://doi.org/10.1007/s10811-010-9583-x>
68. Monder, M. J.; Kozakiewicz, P.; Jankowska, A. The Role of Plant Origin Preparations and Phenological Stage in Anatomy Structure Changes in the Rhizogenesis of *Rosa "Hurdal"*. *Front Plant Sci* **2021**, *7*, 12. <https://doi.org/10.3389/fpls.2021.696998>.
69. Costa, J. M.; Heuvelink, E.; Pol, P. A.; Put, H. M. C. Anatomy and morphology of rooting in leafy rose stem cuttings and starch dynamics following severance. *Acta Hort* **2007**, *751*, 495–502. <https://doi.org/10.17660/ActaHortic.2007.751.63>.
70. Fouda, R. A.; Schmidt, G. Histological changes in the stems of some *Rosa* species propagated by leafy cuttings as affected by IBA treatments. *Acta Agron. Hungar* **1994**, *43*, 265–275.
71. Martins, M.; Gomes, A. F. G.; da Silva, É. M.; da Silva, D. F.; Peche, P. M.; Magalhães, T. A.; Pio, R. Effects of anatomical structures and phenolic compound deposition on the rooting of olive cuttings. *Rhizosphere* **2022**, *23*, 100557. <https://doi.org/10.1016/j.rhisph.2022.100557>.
72. Lei, C.; Fan, S.; Li, K.; Meng, Y.; Mao, J.; Han, M.; Zhao, C.; Bao, L.; Zhang, D. iTRAQ-based proteomic analysis reveals potential regulation networks of IBA-induced adventitious root formation in apple. *Int. J. Mol. Sci.* **2018**, *19*, 667. <https://doi.org/10.3390/ijms19030667>.
73. Li, S.W.; Xue, L.; Xu, S.; Feng, H.; An, L. Mediators, genes and signaling in adventitious rooting. *Bot. Rev.* **2009**, *75*, 230–247. <https://doi.org/10.1007/s12229-009-9029-9>.
74. Della Rovere, F.; Fattorini, L.; D'Angeli, S.; Veloccia, A.; Falasca, G.; Altamura, M.M. Auxin and cytokinin control formation of the quiescent centre in the adventitious root apex of *Arabidopsis*. *Ann. Bot.* **2013**, *112*, 1395–1407.
75. Blazich, F.A. *Chemicals and formulations used to promote adventitious rooting*. In *Adventitious Root Formation in Cuttings*; Davis, T.D., Haissig, B.E., Sankhla, N., Eds.; Dioscorides Press: Portland, OR, USA, **1988**, 132–149.
76. Carpenter, W.J.; Cornell, J.A. Auxin application duration and concentration govern rooting of *Hibiscus* stem cuttings. *J. Am. Soc. Hort. Sci.* **1992**, *117*, 68–74. <https://doi.org/10.21273/JASHS.117.1.68>.
77. Iapichino, G.; Arnone, C.; Bertolino, M.; AmicoRoxas, U. Propagation of three *Thymus* species by stem cuttings. *Acta Hort.* **2006**, *723*, 411–413. <https://doi.org/10.17660/ActaHortic.2006.723.57>.
78. Sabatino, L.; D'anna, F.; Iapichino, G. Cutting type and IBA treatment duration affect *Teucrium fruticans* adventitious root quality. *Not. Bot. Horti Agrobot. Cluj Napoca* **2014**, *42*, 478–481. <https://doi.org/10.15835/nbha4229611>.
79. Kashefi, M.; Zarei, H.; Bahadori, F. The Effect of Indole Butyric Acid and the Time of Stem Cutting Preparation on Propagation of Damask Rose Ornamental Shrub. *JOP* **2014**, *4*, 237–243.
80. Pacholczak, A. The effect of the auxin application methods on rooting of *Physocarpus opulifolius* Maxim. cuttings. *Propag. Ornam. Plants* **2015**, *15*, 147–153.
81. Zhang, L.; Wang, S.; Guo, W.; Zhang, Y.; Shan, W.; Wang, K. Effect of Indole-3-Butyric Acid and rooting substrates on rooting response of hardwood cuttings of *Rhododendron fortune* Lindl. *Propag. Ornam. Plants* **2015**, *15*, 79–86.
82. Kaviani, B.; Gholami, S. Improvement of Rooting in *Forsythia x intermedia* Cuttings by Plant Growth Regulators. *JOP* **2016**, *6*, 125–131.
83. Sabatino, L.; D'Anna, F.; Iapichino, G. Improved Propagation and Growing Techniques for Oleander Nursery Production. *Horticulturae* **2019**, *5*, 55. <https://doi.org/10.3390/horticulturae5030055>.
84. Kashyap, U.; Chandel, A.; Sharma, D.; Bhardwaj, S.; Bhargava, B. Propagation of *Jasminum parkeri*: A Critically Endangered Wild Ornamental Woody Shrub from Western Himalaya. *Agronomy* **2021**, *11*, 331. <https://doi.org/10.3390/agronomy11020331>.

85. Dick, J.M.; Leakey, R.R.B. Differentiation of the dynamic variables affecting rooting ability in juvenile and mature cuttings of cherry (*Prunus avium*). *J. Hortic. Sci. Biotechnol.* **2006**, *81*, 296–302. <https://doi.org/10.1080/14620316.2006.11512065>.
86. Sikorska, A.; Gugala, M.; Zarzecka, K.; Domański, Ł.; Mystkowska, I. Morphological Features of Winter Rape Cultivars Depending on the Applied Growth Stimulators. *Agriculture* **2022**, *12*, 1747. <https://doi.org/10.3390/agriculture12101747>.
87. Chanthini, K.M.-P.; Senthil-Nathan, S.; Pavithra, G.-S.; Asahel, A.-S.; Malarvizhi, P.; Murugan, P.; Deva--Andrews, A.; Sivanesh, H.; Stanley-Raja, V.; Ramasubramanian, R.; Ghaith, A.; Abdel-Megeed, A.; Krutmuang, P. The Macroalgal Biostimulant Improves the Functional Quality of Tomato Fruits Produced from Plants Grown under Salt Stress. *Agriculture* **2023**, *13*, 6. <https://doi.org/10.3390/agriculture13010006>.
88. Sujeeth, N.; Petrov, V.; Guinan, K.J.; Rasul, F.; O'Sullivan, J.T.; Gechev, T.S. Current Insights into the Molecular Mode of Action of Seaweed-Based Biostimulants and the Sustainability of Seaweeds as Raw Material Resources. *Int. J. Mol. Sci.* **2022**, *23*, 7654. <https://doi.org/10.3390/ijms23147654>.
89. Gomes, E.N.; Vieira, L.M.; Tomasi, J.D.C.; Tomazzoli, M.M.; Grunennvaldt, R.L.; Fagundes, C.D.M.; Machado, R.C.B. Brown seaweed extract enhances rooting and roots growth on *Passiflora actinia* Hook stem cuttings. *Ornam. Hortic.* **2018**, *24*, 269–276.
90. Parađiković, N.; Teklić, T.; Zeljković, S.; Lisjak, M.; Špoljarević, M. Biostimulants research in some horticultural plant species—A review. *Food Energy Secur.* **2019**, *8*(2), e00162. <https://doi.org/10.1002/fes3.162>.
91. Toscano, S.; Ferrante, A.; Branca, F.; Romano, D. Enhancing the Quality of Two Species of Baby Leaves Sprayed with Moringa Leaf Extract as Biostimulant. *Agronomy* **2021**, *11*, 1399. <https://doi.org/10.3390/agronomy11071399>.
92. Kapczyńska, A.; Kowalska, I.; Prokopiuk, B.; Pawłowska, B. Rooting Media and Biostimulator Goteo Treatment Effect the Adventitious Root Formation of *Pennisetum 'Vertigo'* Cuttings and the Quality of the Final Product. *Agriculture* **2020**, *10*, 570. <https://doi.org/10.3390/agriculture10110570>.
93. Nowakowska, K.; Pacholczak, A. Effect of the biopreparation "Goteo" on rooting of *Hydrangea* stem cuttings (*Hydrangea paniculata* siebold Limelight and Vanille Freise®Renhy). *Propag. Ornam. Plants* **2017**, *17*, 126–133.
94. Salachna, P.; Zawadzinska, A.; Piechocki, R.; Wilas, J. Propagation of arabian star flower (*Ornithogalum arabicum* L.) by twin scales using seaweed extracts. *Folia Pomeranae Univ. Technol. Stetin.* **2014**, *310*, 105–112.
95. Aremu, A.O.; Makhaye, G.; Tesfay, S.Z.; Gerrano, A.S.; Du Plooy, C.P.; Amoo, S.O. Influence of Commercial Seaweed Extract and Microbial Biostimulant on Growth, Yield, Phytochemical Content, and Nutritional Quality of Five *Abelmoschus esculentus* Genotypes. *Agronomy* **2022**, *12*, 428. <https://doi.org/10.3390/agronomy12020428>.
96. Makhaye, G.; Aremu, A.O.; Gerrano, A.S.; Tesfay, S.; Du Plooy, C.P.; Amoo, S.O. Biopriming with seaweed extract and microbial-based commercial biostimulants influences seed germination of five *Abelmoschus esculentus* genotypes. *Plants* **2021**, *10*, 1327. <https://doi.org/10.3390/plants10071327>.
97. Pacholczak, A.; Szydło, W.; Jacygrad, E.; Federowicz, M. Effect of auxins and the biostimulator AlgaminoPlant on rhizogenesis in stem cuttings of two dogwood cultivars (*Cornus alba* 'Aurea' and 'Elegantissima'). *Acta Sci. Pol. Hortorum Cultus* **2012**, *11*, 93–103.
98. Rathore, S.; Chaudhary, D.; Boricha, G.; Ghosh, A.; Bhatt, B.; Zodape, S.; Patolia, J. Effect of seaweed extract on the growth, yield and nutrient uptake of soybean (*Glycine max*) under rainfed conditions. *S. Afr. J. Bot.* **2009**, *75*, 351–355. <https://doi.org/10.1016/j.sajb.2008.10.009>.
99. Kocira, A.; Lamorska, J.; Kornas, R.; Nowosad, N.; Tomaszewska, M.; Leszczyńska, D.; Kozłowicz, K.; Tabor, S. Changes in biochemistry and yield in response to biostimulants applied in bean (*Phaseolus vulgaris* L.). *Agronomy* **2020**, *10*, 189. <https://doi.org/10.3390/agronomy10020189>.
100. Caccialupi, G.; Caradonia, F.; Ronga, D.; Ben Hassine, M.; Truzzi, E.; Benvenuti, S.; Francia, E. Plant Biostimulants Increase the Agronomic Performance of Lavandin (*Lavandula x intermedia*) in Northern Apennine Range. *Agronomy* **2022**, *12*, 2189. <https://doi.org/10.3390/agronomy12092189>.
101. Quan, J.; Ni, R.; Wang, Y.; Sun, J.; Ma, M.; Bi, H. Effects of Different Growth Regulators on the Rooting of *Catalpa bignonioides* Softwood Cuttings. *Life* **2022**, *12*, 1231. <https://doi.org/10.3390/life12081231>