atmosphere

MDPI

Type of the Paper (Article)

Improving model performance of ecosystem carbon cycle by integrating soil erosion related processes

Jinliang Zhang[1,2] , Chao, Zhang[1,2], Wensi, Ma[1,2*], Wei Wang[1,2], Haofei, Li[1,2]

[1] Yellow River Engineering Consulting Co. Ltd, Zhengzhou, China.
[2] Key Laboratory of Water Management and Water Security for Yellow River Basin of Ministry of Water Resources , Zhengzhou, China
* Correspondence: mws@lzb.ac.cn; Tel.:13592610328

Abstract: Soil erosion is a key factor in soil quality degradation and carbon balance in arid ecosystems. However, many model ignores soil erosion process in arid regions, which may lead to limit the understanding of ecosystem processes in arid regions. In this study, we added the soil erosion process according to field observed data of soil hydrothermal regime and carbon flux. We validated this coupling version of IBIS (Integrated Biosphere Simulator) and RUSLE (RU-IBIS) by examining four different vegetation types and the carbon budget in the arid region on the Loess Plateau (LP). Our results indicated that the coupling model (RU-IBIS) produced more reliable simulations of the soil water content (the r from 0.23-0.90 to 0.71-0.97), evaporation (ET) (the average r is) and significantly improved the simulation of the leaf area index (LAI) (the average r is 0.95) , net primary production (NPP) (the average r is 0.95). We also conducted sensitivity experiments to determine how the soil texture and aerodynamic roughness (Z_{0m}) affect the soil water content. Moreover, it revealed that specific leaf area (SLA) plays a key role in the simulation of NPP and NEE. Our study suggested that coupled soil erosion process and parameterization can effectively improve the performance of IBIS in arid regions. These results need to be considered in future Earth system models.

Keywords: IBIS; RUSLE; soil water content; soil erosion process; carbon cycle; Loess Plateau

1. Introduction

As a component of terrestrial ecosystems, arid regions account for about 46% of global carbon pool [1]. Arid regions have low precipitation, high solar insolation, whose evaporation is much higher than precipitation, making the ecosystem relatively fragile and more vulnerable to external climate impacts [2]. Insufficient water and human activities such as overgrazing and excessive land reclamation have also led to serious land degradation in arid regions [3]. The land degradation releases carbon into the atmosphere, and may form a positive feedback loop within the carbon cycle [4]. Therefore, it is of great importance to study the processes of carbon cycling in arid region ecosystems in the context of climate change.

Simulating and predicting the arid ecosystem carbon cycle is inseparable from the dynamic global vegetation model (DGVM) [5]. Using DGVM, many studies have been conducted to simulate the change of carbon budget and its influencing factors on the arid regions [6-9]. Under the influence of long-term soil erosion, the nutrient-rich topsoil layer is gradually denuded and the soil nutrients are lost to runoff and sediment. Soil nutrients are lost with runoff and sediment, leading to a decrease in soil fertility and thus reducing the erosion resistance of the soil. This results in a vicious cycle of "soil erosion-nutrient loss-soil erosion" [10,11]. Soil erosion leads to thinning of the soil layer, decreasing in soil aeration, permeability and soil quality [12,13]. The soil erosion process strongly affects the energy and water exchange between soil and air, and thus definitely affects the simulation accuracy of carbon cycle in the arid regions. Soil erosion must be included in the study of

Citation: To be added by editorial staff during production.

Academic Editor: Firstname Lastname

Received: date
Revised: date
Accepted: date
Published: date

Copyright: © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/licenses/by/4.0/).

carbon cycle in arid regions [14-15]. However, in many DVGMs, this process was only expressed as simple soil erosion parameterization schemes [16-18].

IBIS has been developed to illuminate the interactions between different biosphere factors on time scales relevant to the carbon cycle process [19]. In recent years, some work have started to apply IBIS on a global scale [20]. Further refinements of the IBIS model include the creation of IBIS 2.0 and Agro-IBIS [21-23], which incorporate other models of atmospheric circulation, agricultural vegetation, and hydrological processes. In terms of model improvement, Liu et al. (2005) combined IBIS with the nitrogen cycle model to simulate the regional nitrogen cycle and to reveal the extent to which the nitrogen cycle affects the ecosystem [24]. While most ecosystems examined with IBIS are forests and farmland [25-27], recent updates to IBIS include adding more vegetation types and statistical models [28,29]. However, the IBIS model does not consider changes to the soil erosion regime. The absence of this process may produce model results that overestimate the moisture and carbon budget in arid regions.

We aim to using DVGMs to modelling the arid ecosystem carbon cycle, the objectives of the study were to (1) incorporate Revised Universal Soil Loss Equation (RUSLE) into the soil erosion and verified the accuracies of modelling soil water content, LAI and NPP simulations in Loess Plateau(LP); (2) determined sensitivity of the soil hydrothermal regime, ET(evapotranspiration) and the carbon flux to the changing model parameters. The modified IBIS model provides a useful tool for understanding the carbon cycle of arid regions in warming climate conditions.

2. Materials and Methods

2.1 Experimental sites

The Loess Plateau is the largest loess accumulation area in the world[30], which is located in the north-central of China, with soil erosion covering 71%, and the average thickness of loess in the region is 92.2 m. Due to its vegetation degradation, soil nutrient loss, land quality degradation, and ecological environment deterioration, the LP is very sensitive to climate change. We collected data from four regions in the LP (Table 1). These four stations are Shapotou (SPT: cropland), Changwu (CW: cropland) Ansai (AS: cropland) and Eerduosi (ERDS: desert). The detailed information of the four stations were summarized in Table 1.




Fig.1. Location of research stations in the LP.

Table 1. Site information for (SPT, CW, AS, ERDS) in LP.

Site		SPT	CW	AS	ERDS
Observation period		2001-2008	2004-2008	2004-2006	2005
Vegetation type		cropland	cropland	cropland	desert
Latitude	**N°**	37°27′	35°12′	36°51'	39°29′
Longitude	**E°**	104°57′	107°40′	109°19′	110°11′
Elevation	**m**	1250	1200	1068	1290
mean precipitation	**mm**	186	580	500	310
mean temperature	**℃**	9.6	9.1	8.8	6

2.2.Input and observed data

The meteorological data collected at four stations operated under the auspices of National Ecosystem Science Data Center, National Science & Technology Infrastructure of China (http://www.nesdc.org.cn). The period of single-point simulation is 1998-2008. The meteorological data include wind speed, relative humidity, pressure, radiation, temperature, precipitation, etc. The basic processing procedures include: anomaly rejection, blank interpolation, half-hourly synthesis of daily data, and unit conversion. The processed meteorological data are used as single-point simulated climate-driven data. Model validation data include soil water content, evapotranspiration, above-ground biomass, below-ground biomass, and leaf area index (LAI) [31-34]. In this study, the measured biomass values were converted to NPP values for model validation [35].

2.3. Model description and development

Foley et al. (1996) developed the biosphere-oriented, regional-scale Integrated Biosphere Simulator (IBIS) model [20]. In order to simulate the dynamic changes in the vegetation cover in Europe and Africa, the second iteration of the IBIS model was coupled with GENESIS (version 2.0), an atmospheric circulation model. The most recent version of IBIS, version 2.6 [35]. The IBIS model consists of five modules: land surface processes, vegetation dynamics, vegetation phenology, subterranean carbon and nitrogen cycling, and soil geochemistry. Our study focuses on improving and refining the land surface processes and vegetation dynamics modules.

(1) Land surface processes module

The land surface processes module is used to calculate the exchange of water, heat, and carbon dioxide. The thicknesses of the six soil layers, from top to bottom, are 0.10 m, 0.15 m, 0.30 m, 0.50 m, 1.00 m, and 2.00 m. In the soil module, the rate of change of the liquid soil water content is calculated using the Richard equation and the vertical flux of water is determined using Darcy's law. At the impermeable bottom boundary of the soil module, there is no diffusion of liquids or heat.

To simulate the soil hydrothermal regime, we calculate the soil temperature, the water content, and the ice content of a given soil layer:

$$\frac{\partial \theta}{\partial t} = \frac{\partial}{\partial z}[D(\theta)\frac{\partial \theta}{\partial z}] + \frac{\partial K(\theta)}{\partial z} - ST(z,t) \quad (1)$$

where $\theta(m^3m^{-3})$ is the actual soil water content content of a soil layer at a certain time, and $\theta i(m^3m^{-3})$ is the ice rate of the soil layer. $D(\theta)$ (mms^{-1})represents the soil diffusivity when the soil volumetric water content. $K(\theta)(mms^{-1})$ is the hydraulic conductivity, when the soil volumetric moisture content is $K(\theta)$ needs to be calculated by the saturated hydraulic conductivity Ks, b is the index of the campbell formula. $ST(z,t)(s^{-1})$ is a sink term resenting water uptake by plant roots as a function of ET.

$$K(\theta) = K_S\left(\frac{\theta}{\theta_S}\right)^{2b+3} \quad (2)$$

Using the heat diffusion equation, we can quantify the heat transfer in the soil:

$$C\frac{\partial T(z,t)}{\partial t} = \frac{\partial}{\partial z}\left[\frac{\lambda \partial T(z,t)}{\partial z}\right] + H_{fm} \quad (3)$$

$$H_{\text{fm}} = \rho_{ice} L_f \frac{\partial \theta_{ice}}{\partial t} \quad (4)$$

where C(Jkg^{-1}k^{-1}) and λ(Wm^{-1}k^{-1}) are the volumetric heat capacity and the thermal conductivity, respectively, t(s) is the time, z (mm) is the vertical depth. T(k) is the temperature. Hfm (Jkg^{-1}) is the latent heat released or consumed during phase change of soil water. ϱice (917 kgm^{-3}) is the density of ice, and θice(m^{3}m^{-3}) is the partial volume of ice content, Lf (0.3336×106 Jkg^{-1}) is the latent heat of fusion.

(2) Vegetation dynamic module

After registering a change in the biomass for each plant type, this module calculates the gross primary productivity (GPP), the net primary productivity (NPP), the net ecosystem exchange (NEE) in the vegetation coverage and type. The vegetation productivity calculation includes both the total photosynthesis and the breathing per hour. Vegetation biomass is the sum of the root system biomass and the plant leaves. The GPP calculation is:

$$GPP = \int A_g dt \quad (5)$$

and the NPP calculation is:

$$NPP = (1-\eta)\int (A_g - R_{\text{leaf}} - R_{\text{stem}} - R_{root})dt \quad (6)$$

where Ag (CO_2/m^2·s) is the net photosynthetic rate of the canopy. Rleaf (CO_2/m^2·s) is the leaf respiration rate. Rstem CO_2/m^2·s) is the respiration rate; Rroot (CO_2/m^2·s) is the root respiration rate. The rate of photosynmesis is simulated according to the Farquhar equations [36,37]. η is 0.33 or 0.3 [38], which is the proportion of carbon loss in tissue construction.

(3) Vegetation Phenology Module

In the phenological module, it is assumed that deciduous plants begin to germinate in winter when the growth of a particular plant functional type exceeds a certain threshold. This threshold is defined by the day when two conditions are met: 1) the average temperature (Avt10) is greater than 5°C for 10 consecutive days and 2) at least 100 days have experienced temperature increases greater than 0°C (GDD_0). When Avt10 is less than 5°C, the vegetation enters the withered yellow period. This threshold day forms the basis of the leaf area index (LAI) calculations:

$$LAI = LAI_{\max} * \mathrm{f}(T) \quad (7)$$

$$\mathrm{f}(T) = \begin{cases} (GDD_0 - T_b)/b & GDD_0 > T_b \text{且} avt10 > 5 \\ 1 & \\ \mathrm{f}(T) - 1/dfac & avt10 < 5 \end{cases} \quad (8)$$

where Tb is the threshold of GDD_0 in vegetation green-back period, b is the coefficient of controlling leaf spreading speed, *dfac* is the coefficient of controlling defoliation speed.

(4) Modified soil erosion parameterization scheme

Numerous studies have pointed out that soil erosion and erosion of poorly vegetated land are accompanied by changes in soil nutrients, structure, moisture, texture, etc. The soil microbial community will also be transformed during the restoration of soil erosion and erosion-damaged land [39,40]. As the main driver of soil organic carbon transfor-

mation, microbial community changes in erosion and revegetation systems largely determine the rate of organic carbon mineralization and fixation and the microbial contribution to soil organic carbon. The microbial communities of erosion and revegetation systems largely determine the rate of organic carbon mineralization and fixation and the apparent functional redundancy of microorganisms in the soil. Therefore, we used RUSLE to improves the soil erosion process module. The expression of general soil loss equation is :

$$A = R \cdot K \cdot LS \cdot C \cdot P \quad (9)$$

where A is the soil loss (kg km^{-2}), R is rainfall erosivity factor (J m^{-2}), K is soil erodibility factor (dimensionless), LS is slope length and gradient factor (dimensionless), C: vegetation cover and management factor (dimensionless) P: Soil and water conservation measures factor (dimensionless)




Fig. 2. Modeling framework and improvement modules (refer to [35])

(5)Model parameterization

Previous modifications to the IBIS model [41] suggested that accurate model results were dependent upon the accurate calibration of stomatal conductance coefficients (m , b), leaf respiration parameter (γ) and the maximum rubisco capacity (Vmax). For the photosynthesis of vegetation in arid regions, these parameters were set to $\gamma = 0.025$, m = 4.0, b = 0.04, and Vmax = 25 molCO_2 m^{-2} s^{-1} [42]. We set these four parameters in the Vegetation dynamic module.

Aerodynamic roughness is a parameter that is important for describing the aerodynamic characteristics in surface heat flux parameterization scheme, which results in large uncertainties in the soil hydrothermal condition calculations at smaller spatial scales [43]. Therefore, we changed the initial Z_{0m} (0.005 m) to be equal to the two annual average Z_{0m} for the two vegetation types along LP based on the measured data, the cropland (0.29) and the desert (0.0003) (Table 2) .

The specific leaf area (SLA) can be used not only to infer the processes of photosynthesis, evaporation, and ET, but also to estimate the net productivity of terrestrial ecosystems. In an individual plant or group of plants, the amount of light received by the vegetation is inversely proportional to the SLA value. The SLA value can also reflect the ability of plants to obtain light; plants with low SLA values can better adapt to poor treatment and drought environments, while plants with high SLA values can more easily preserve vital nutrients [44,45]. This study used SLA that correspond to the vegetation conditions in arid region to improved quality of our model results (Table 2)[46,47].

Table 2. Parameter differences between the O-IBIS and RU-IBIS model (M-IBIS).

Site		SPT		CW		AS		ERDS	
		O-IBIS	RU-IBIS	O-IBIS	RU-IBIS	O-IBIS	RU-IBIS	O-IBIS	RU-IBIS
Z_{0m}	**(m)**	0.005	0.29	0.005	0.29	0.005	0.29	0.005	0.0003
SLA	**(m^2/kg)**	20	16.04	20	11.69	20	10.9	20	11.69

*Note: Z_{0m}: Thermodynamic roughness. SLA: Specific leaf area. O-IBIS: original IBIS. RU-IBIS: modified IBIS References: [46,47]

2.5.Model validation

(1) Correlation analysis

We used many statistical methods to quantitatively evaluate the applicability of our models, including the relative error (RE), the correlation coefficient r, the root mean square error (RMSE) and the coefficient of determination R^2. Detailed descriptions of these statistical tests and formulas are found in Willmott et al. (1982) [48].

(2) Parameter sensitivity analysis

To investigate how parameters within IBIS could influence the simulation results, we ran 16 experiments. Individual model parameters were individually increased or decreased by 20% in each run. Overall model sensitivity to NPP, ET, SW, and ST (multilayer soil depth averages) was evaluated by comparing simulated annual averages from the modified parameter runs to the control runs.

(3) Redundancy analyses

We conducted redundancy analyses (RDA), both with and without variance partitioning, to estimate both the climate effects and the nature of the relationship between the site and the climate effects [49]. Here, RDA was used to analyse the relationships of carbon flux to climate variables. These analyses were performed using the vegan package in R.

3. Results

3.1.Soil water content conditions

Comparison of the simulated and measured soil water content values at the four sites is shown in Figure 3, where the simulated daily temperature trends are in good agreement with the observations. RU-IBIS generated improved soil water content estimates for the four stations, with a higher range of r values (from 0.23-0.90 to 0.71-0.97) and RMSE, ,RE values of 0.15-5.69 m^3/m^3, -0.12-0.27 m^3/m^3, respectively (Table 3). The underestimation of soil water content has been greatly improved. In terms of different depths: the correlation between the simulated results and the measured values of soil water content at 10 cm is the weakest. With the increase of soil depth, the deviation value decreases sequentially with the increase of soil depth, which is related to the fact that the deeper the soil depth, the smaller the actual amplitude. The RU-IBIS model has the highest simulation accuracy in the cropland (CW station), and the simulation effect is relatively poor in the desert region (ERDS). Our statistical analyses for these parameters are shown in Table 3.




Fig 3. The observed, O-IBIS, RU-IBIS values of monthly soil water content at soil depths of 10 cm, 30 cm, 50 cm, and 100 cm in different regions: (a-d) SPT, (e-h) AS, (i-l) CW, and (m-p) ERDS. The red and gray lines represent the linear fit of improved and original simulations, respectively.

Table 3. Statistical measures of modeled soil water content at monthly time scale.

Stations		10cm		30cm		50cm		100cm	
		RU-IBIS	O-IBIS	RU-IBIS	O-IBIS	RU-IBIS	O-IBIS	RU-IBIS	O-IBIS
SPT	r	0.71	0.45	0.88	0.78	0.91	0.82	0.92	0.90
	R^2	0.51	0.20	0.78	0.62	0.84	0.67	0.84	0.82
	RMSE	3.23	4.42	0.71	1.42	0.60	1.43	1.05	3.58
	RE	-0.08	0.09	-0.12	0.17	-0.12	-3.02	0.17	-0.04
AS	r	0.87	0.71	0.88	0.58	0.86	0.60	0.82	0.59
	R^2	0.75	0.51	0.77	0.33	0.73	0.36	0.68	0.34
	RMSE	2.67	3.62	1.77	5.60	1.81	5.25	1.89	4.81
	RE	0.07	-0.01	-0.03	-0.46	0.08	-0.31	-0.07	-0.43
CW	r	0.94	0.88	0.97	0.46	0.90	0.65	0.91	0.63
	R^2	0.88	0.77	0.94	0.21	0.82	0.42	0.82	0.40
	RMSE	5.69	7.26	1.44	5.29	0.64	3.78	1.06	3.69
	RE	0.27	0.37	-0.05	0.09	-0.11	-0.42	-0.09	-0.13
ERDS	r	0.84	0.34	0.87	0.74	0.91	0.60	0.76	0.23
	R^2	0.70	0.11	0.76	0.55	0.83	0.35	0.58	0.06
	RMSE	1.12	1.67	0.29	0.32	0.15	0.26	0.34	0.73
	RE	-0.08	-0.19	-0.10	-0.39	0.07	-2.91	0.19	-0.35

3.2 ET simulation results

Due to the limitation of monitoring conditions, only monthly average daily ET data of SPT (2003-2004), AS (2004-2006) and CW (2004) stations were observed. In Figure 4, it can be seen that the simulation result of evapotranspiration is higher than the measured

data. The r of ET were 0.61 (SPT), 0.83(AS), 0.86(CW). RMSE and RE were 2.06 mm, -0.05 mm(SPT), 2.07mm and -0.20mm (AS), 0.75 mm and 0.17mm (CW).




Fig.4 The simulated average daily ET was compared with the measurements in SPT (a), AS (b) and CW (c). The red and gray lines represent the linear fit of improved and original simulations, respectively.

Table 4. Statistical measures of modeled ET.

	SPT				AS				CW			
	r	R^2	RMSE	RE	r	R^2	RMSE	RE	r	R^2	RMSE	RE
RU-IBIS	0.61	0.38	2.06	-0.05	0.83	0.69	2.07	-0.20	0.86	0.75	0.75	0.17
O-IBIS	0.23	0.05	2.97	-23.93	0.41	0.17	3.49	0.76	0.59	0.35	1.18	0.95

3.3 LAI and NPP simulation

The original IBIS underestimated the LAI values, the RU-IBIS values, which had smaller uncertainties than the O-IBIS values, generally agreed with the observed LAI values (Fig. 5). The RU-IBIS model generated LAI correlation coefficient (0.95) and R^2 (0.90) results that were higher than those of the O-IBIS model (0.70 and 0.50, respectively). Furthermore, the RMSE and RE values simulated by the RU-IBIS results revealed improving O-IBIS results on LAI and NPP basis (Table 5). Clearly, the RU-IBIS dynamic vegetation results are more accurate than those of the O-IBIS model.

Figure (5c, d) showed the results between observed,O-IBIS and RU-IBIS NPP at SPT and AS stations, which demonstrated that the RU-IBIS simulations are maintained consistency the observed NPP. The NPP of O-IBIS was much lower than the observed value. After modifying IBIS, the underestimation phenomenon improved significantly. The correlation coefficients were 0.52 (O-IBIS) and 0.95 (RU-IBIS), respectively. The statistical metrics for accuracy and error improved when we used the RU-IBIS results instead of the O-IBIS results.




Fig 5. Comparison scatter plot of O-IBIS, RU-IBIS, and observed LAI, NPP value in the (a, c) SPT and (b, d) AS stations. The red and gray lines represent the linear fit of improved and original simulations, respectively.

Table 5. Statistical analyses of modeled monthly LAI and NPP.

		SPT				AS			
		r	R^2	RMSE	RE	r	R^2	RMSE	RE
LAI	RU-IBIS	0.94	0.88	0.60	0.02	0.96	0.92	0.47	-0.06
	O-IBIS	0.77	0.60	1.39	0.56	0.63	0.40	1.46	-9.67
NPP	RU-IBIS	0.96	0.92	474.11	0.29	0.93	0.87	318.18	-0.38
	O-IBIS	0.23	0.05	1188.11	-0.43	0.81	0.65	467.82	0.38

3.4 RDA analysis

Figures 6 and 7 show that the arid region ecosystem is highly sensitive to changes in the temperature. While warming temperatures are minimally impacted by precipitation, NPP tends to rise along with the temperature. The two consecutive RDA axes (Fig. 6)suggested the pattern of carbon fluxes with climate change and illustrate that the first RDA axis (66.84%) is absolutely dominant.The second RDA axis accounts for 17% of the carbon flux variation. Carbon flux variation on the second axis is indicative of warmer, higher radiation ecosystems with higher NPP, NEE, and Re values. Cloud amount, temperature, and soil temperature have the greatest influence on the carbon flux. In the LP, vegetation growth and soil respiration are very sensitive to temperature and radiation changes.

The pie chart in Figure 7 summarizes the contributions of climate and site to the different ecosystem carbon fluxes. Climate is shown nested within research sites because differences in climate are determined by the site location. The three temperature parameters jointly account for 31% of the NEE variation (Fig. 7), the most important controlling factor of NPP is the cloud amount (29.2%), and the soil respiration is most sensitive to changes in the site (63.2%). Different sites mean different soil textures, vegetation types, and ecosystems. Different ecosystems and ecosystem locations lead to different growth season onsets, growth season durations, and carbon emission intensities.




Fig 6. Climate-related carbon flux from redundancy analysis (RDA): coloured dots are different vegetation sites. The principal variables are: net primary production (NPP), net ecosystem exchange (NEE), soil respiration (Re). The climate variables are: the highest temperature(tmax), average temperature(tm), minimum temperature(tmin), precipitation(prec), wind speed(windm), cloud amount(cloud), relative moisture(em), pressure(pm), soil tempreture (ST), soil water content (SW).




Fig 7. Pie chart of total explained variation of environmental factors in simulated carbon flux

4. Discussion

4.1 Improvement of IBIS model

Our results revealed that soil erosion in O-IBIS is not realistic, as it largely underestimates soil water content (Fig. 3). Also, the simulated ET, LAI and NPP in O-IBIS is significantly lower than observations (Figs.4-5). We tentatively attribute this deficiency to the original IBIS, as most of the soil hydrothermal regime improvements and soil erosion process of arid region in IBIS require more validation and improvements. Previous works have suggested that the reduction of soil fertility and the deterioration of soil structure under the influence of long-term hydraulic erosion structural deterioration will further affect the growth and development of plants. In our study, we not only added the soil erosion process to the IBIS model, but we also recalibrated the physiological parameters according to the actual climate conditions in the study area. These modifications resulted in better fits with the observed data and reduced model uncertainties. We suggest to further collect such detailed ecological factors data at multiple locations to improve the arid vegetation modeling and reduce errors due to multi-site heterogeneity.

Further development always requires field-measured data for calibration. The LP is dominated by C3 plant. Physiological parameters of different vegetation types differ

greatly between C3 plant. RU-IBIS model, with its more complete soil erosion process, may provide better opportunities for parameter calibration in arid regions.

4.2 Parameter sensitivity

As shown in Table 6, the simulation results are most sensitive to changes in the Z_{0m}. For the cropland, a 20% increase in the Z_{0m} decreased ST and NEE by 26.68% and 18.25%, respectively, and increased SW, ET, and Re by 86.43%, 42.71%, and 38.07%, respectively. Therefore, the accuracy of the input parameters plays an important role in the simulation results. Soil water content and ET are greatly affected by the dynamic roughness, which reflects the weakening effect of different ground surface characteristics on the wind speed [43]. The vegetation characteristics and the wind speed cannot be ignored in sparsely vegetated areas. Z_{0m} is negatively correlated with wind speed, and wind speed is positively correlated with ET [50]. An increase in the Z_{0m} parameter represents a decrease in the wind speed, which in turn leads to a reduction in the ET value.

While the +20% change in the SLA value resulted in a moderate NPP increase of 5%-8%, the sensitivity of NEE to SLA is much higher than that of NPP. SLA values can directly affect the efficiency of vegetation in capturing light and CO_2[49]. Higher SLA values means that plants usually have higher productivity values, and that plants with lower specific leaf area values experience more efficient photosynthesis in limited resource conditions [51]. SLA is involved in the area-based biochemical parameterization of photosynthetic enzyme concentrations linking LAI to NPP and provides structural parameters linking leaf carbon distribution to LAI [44,52].

Table 6. The results of model parameters sensitivity experiments

Sites	Parameters	Value	Sensitivity (+20%)						Sensitivity (-20%)					
			ST	SW	ET	NPP	NEE	Re	ST	SW	ET	NPP	NEE	Re
			(%)						(%)					
SPT	Z_{0m}（m）	85	**-26.68**※	**86.43**	**42.71**	1.02	**-18.25**	**38.07**	**-9.62**	**38.02**	**16.84**	-0.31	**-23.09**	**46.37**
	SLA（m^2/kg）	6.85	0.17	0.00	0.01	-0.01	-0.15	0.01	-1.20	-0.01	-0.02	0.03	0.28	-0.02
AS	Z_{0m}（m）	66.4	-0.42	**-28.13**	**-15.64**	0.16	**-10.70**	**-32.72**	8.35	0.39	0.13	-0.17	0.56	0.07
	SLA（m^2/kg）	3.55	0.00	0.00	0.00	0.00	0.00	0.00	-0.31	0.01	0.00	0.01	-0.01	0.00
CW	Z_{0m}（m）	80	-1.30	-15.81	-5.27	0.49	**-62.27**	**-23.44**	**5.69**	**44.15**	**22.12**	-0.27	**10.93**	**41.47**
	SLA（m^2/kg）	4.34	0.96	0.00	0.01	-0.02	0.02	0.00	-0.37	0.00	-0.02	0.01	-0.05	-0.01
ERDS	Z_{0m}（m）	40.2	**13.15**	**-22.28**	9.20	0.60	**-53.31**	**30.55**	3.17	0.10	0.10	-0.11	-4.06	0.12
	SLA（m^2/kg）	54.05	**22.19**	**-22.16**	9.46	0.34	**-54.24**	**30.81**	-4.83	-0.17	-0.09	0.15	7.20	-0.25

*Note: +20%: model parameters increase by 20%; -20%: model parameters reduced by 20%; -26.68 ※: when the Z_{0m} increases by 20%, the soil temperature decreases by 26.68%.

4.3 Potential effects of climate change on the carbon cycle

As the concentrated rainfall, heavy storms and human interference in the Loess Plateau, soil erosion of LP is inevitable. Improving the activity and quantity of soil microorganisms results in accelerated soil respiration and CO_2 emission. Additionally, the organic carbon in the soil, which was originally sealed in the arid region, was released into the ground surface and eventually makes its way into the carbon cycle. However, the temperature increase can improve the photosynthetic rate of vegetation and prolong the growth period of vegetation, resulting in a longer period over which vegetation can draw CO_2 out

of the atmosphere. Similarly, arid region erosion results in decreased soil water content, which inhibited vegetation growth and CO_2 absorption. Increased soil erosion can reduce the soil water content volumetric under the same suction, which is unfavorable to soil water storage and reduces the drought-resistant performance of the soil. The infiltration rate of soil in the 0-10 cm layer gradually decreased with the increase of soil erosion, and the difference of infiltration rate of each soil layer in different vertical depths gradually decreased with the increase of soil erosion [49]. Therefore, the potential impact of arid region erosion on ecosystem carbon emission mainly depends on the arid region balance between the increase in the soil respiration potential caused by warming and the diminution of vegetation photosynthesis in conjunction with the increase in the soil water content [52,53], which is supported by additional field data from the LP [54]. NEE is numerically equal to the difference between NPP and soil heterotrophic respiration. Therefore, NPP could be largely canceled out by a higher ER value, which tends to respond positively to the ecosystem annual carbon emissions [52].

5. Conclusions

The IBIS model has been validated in a variety of different ecosystems, regions, and temperature zones worldwide. We used RU-IBIS to determine appropriateness of applying this model to arid regions in the LP. The results showed that：

1. The O-IBIS model generally underestimates the soil water content in all stations. After adding the soil erosion process scheme and adjusting the dynamic roughness, the ability of the IBIS model to simulate soil water content was greatly improved though still with underestimate bias (RE=-0.013 m^3/m^3). These results indicate that soil erosion plays an important role in controlling soil hydrothermal regime in arid regions. We should pay attention to the reliability of Z_{0m} value in the future simulation of the barren region with IBIS.

2. The underestimation of NPP and LAI was reduced by 68.7% and 88.5%, respectively, through the adjustment of the SLA. The modified model (RU-IBIS) accurately reproduces the vegetation growth conditions in the cropland region, and does a poor job of constraining the conditions in the desert. The results show that the complexities of applying to heterogeneous sites remain, and it need further improvement in a wider range area.

3. Sensitivity tests showed that the soil hydrothermal conditions and ET were sensitive to changes in the Z_{0m} values. SLA largely affects the simulated NPP, NEE at all stations. An increase in the SLA produces corresponding increases in the NPP and NEE values.

4. We infer that the temperature, radiation, and soil water content dictate the interannual variability of the carbon emission (absorption) strength in this simulation. Moreover, any soil water content lost during soil erosion may exacerbate water deficits and thus reduce NPP. However, it is unknown whether the arid region erosion will result in a net increase or decrease in the total carbon emissions in this region.

The RU-IBIS model improves the accuracy of the soil and vegetation modules. By incorporating the soil erosion process into IBIS, we hope to extend our single-site simulation to the whole arid region of the LP and explore the feedback of arid region on global climate change in the future.

Author Contributions: Conceptualization, Jinliang. Zhang.; and Wensi, Ma.; methodology, Wensi, Ma.; software, Chao.Zhang.; validation, Jinliang. Zhang.; and Wensi, Ma.; formal analysis, Wei.Wang.; investigation, Wei.Wang.; resources, Haofei.Li.; data curation, Jinliang. Zhang.; writing—original draft preparation, Wensi, Ma.; writing—review and editing, Jinliang. Zhang; visualization, Jinliang. Zhang.; supervision, Wensi, Ma.; project administration, Wensi, Ma. All authors have read and agreed to the published version of the manuscript." Please turn to the CRediT taxonomy for the term explanation. Authorship must be limited to those who have contributed substantially to the work reported.

Funding: This work is supported by the National Natural Science Foundation of China (41690142).

Data Availability Statement: The data that support the findings of this study are available from the corresponding author upon reasonable request.

Acknowledgments: Dataset is provided by National Ecosystem Science Data Center, National Science & Technology Infrastructure of China. (http://www.nesdc.org.cn).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Huang, J., H. Yu, X. Guan, G. Wang, and R. Guo. Accelerated dryland expansion under climate change, Nat. Clim. Chang, 2016,6, 166-171.
2. Huang, J., H. Yu, A. Dai, Y. Wei, and L. Kang. Drylands face potential threat under CO_2 global warming target, Nat. Clim. Change, 2017,7, 417-422.
3. Sun Dianchao, Li Yulin, Zhao Xueyong, Luo Yayong, Bi Jingdong. Effects of enclosure and grazing on net ecosystem carbon exchange in sandy grasslands of Horqin The effect of enclosure and grazing on the net ecosystem carbon exchange of kerchin sandy grassland. 2016, China Desert, 36(1):93-102.
4. Reynolds, J. F., et al. Global desertification: building a science for dryland development. Science, 2007,316,847-851.
5. Schuur, E. A., Mack, M. C. Ecological response to arid region thaw and consequences for local and global ecosystem services. Annual Review of Ecology, Evolution, and Systematics, 2018,49, 279-301.
6. Piao, S., J. Fang, P. Ciais, P. Peylin, Y. Huang, S. Sitch, and T. Wang. The carbon balance of terrestrial ecosystems in China. Nature, 2009,458(7241):1009-1013.
7. Yi, S., Wang, X., Qin, Y., Xiang, B., Ding, Y. Responses of alpine grassland on Qinghai–Tibetan plateau to climate warming and arid region degradation: a modeling perspective.Environmental Research Letters, 2014,9(7), 074014.
8. Zheng, G., Yang, Y., Yang, D., Dafflon, B., Lei, H., Yang, H. . Satellite-based simulation of soil freezing/thawing processes in the northeast Tibetan Plateau. Remote Sensing of Environment, 2019,231, 111269.
9. Chen, W., Zhu, D., Ciais, P., Huang, C., Viovy, N., Kageyama, M. Response of vegetation cover to CO_2 and climate changes between Last Glacial Maximum and pre-industrial period in a dynamic global vegetation model. Quaternary Science Reviews, 2019,218, 293-305.
10. Lal R. Soil erosion and the global carbon budget. Environ Int, 2003, 29:437-450.
11. Borrelli P, Robinson DA, Fleischer LR, Lugato E, Ballabio C, Alewell C, Meusburger K, Modugno S, Schutt B, Ferro V, Bagarello V, Van Oost K, Montanarella L, Panagos P. An assessment of the global impact of 21st century land use change on soil erosion. Nat Commun, 2017, 8.
12. Montgomery DR. Soil erosion and agricultural sustainability. P Natl Acad Sci USA, 2007, 104:13268-13272.
13. Verheijen FG A, Jones RJ A, Rickson RJ, Smith CJ. Tolerable versus actual soil erosion rates in Europe. Earth-Sci Rev, 2009, 94:23-38.
14. Van Oost K, Quine TA, Govers G, De Gryze S, Six J, Harden JW, Ritchie JC, Mccarty GW, Heckrath G, Kosmas C, Giraldez JV, Da Silva JR, Merckx R. The impact of agricultural soil erosion on the global carbon cycle. Science, 2007, 318:626-629
15. Maavara T, Lauerwald R, Regnier P, Van Cappellen P. Global perturbation of organic carbon cycling by river damming. Nat Commun, 2017, 8:15347.
16. Wang ZG, Hoffmann T, Six J, Kaplan JO, Govers G, Doetterl S, Van Oost K. Human-induced erosion has offset one-third of carbon emissions from land cover change. Nat Clim Change, 2017b, 7:345-349.
17. Beer, C., Lucht, W., Gerten, D., Thonicke, K., Schmullius, C. Effects of soil freezing and thawing on vegetation carbon density in Siberia: A modeling analysis with the Lund-Potsdam-Jena Dynamic Global Vegetation Model (LPJ-DGVM). Global Biogeochemical Cycles, 2007,21(1).
18. Wania, R., Ross, I., Prentice, I. C. Integrating peatlands and arid region into a dynamic global vegetation model: 2. Evaluation and sensitivity of vegetation and carbon cycle processes.Global Biogeochemical Cycles, 2009,23(3).
19. Wang, H., Ma, M., Wang, X., Yuan, W., Song, Y., Tan, J., Huang, G. Seasonal variation of vegetation productivity over an alpine meadow in the Qinghai–Tibet Plateau in China: modeling the interactions of vegetation productivity, phenology, and the soil soil erosion–thaw process.Ecological research, 2013,28(2), 271-282.
20. Foley, J. A., Prentice, I. C., Ramankutty, N., Levis, S., Pollard, D., Sitch, S., Haxeltine, A. An integrated biosphere model of land surface processes, terrestrial carbon balance, and vegetation dynamics. Global biogeochemical cycles, 1996,10(4), 603-628.
21. Delire, C., Foley, J. A. Evaluating the performance of a land surface/ecosystem model with biophysical measurements from contrasting environments. Journal of Geophysical Research: Atmospheres, 1999,104(D14), 16895-16909.
22. Twine, T. E., Kucharik, C. J. Evaluating a terrestrial ecosystem model with satellite information of greenness.Journal of Geophysical Research: Biogeosciences, 2008,113(G3).
23. Zipper, S. C., Soylu, M. E., Kucharik, C. J., Loheide II, S. P. Quantifying indirect groundwater-mediated effects of urbanization on agroecosystem productivity using MODFLOW-AgroIBIS (MAGI), a complete critical zone model. Ecological modelling, 2017,359, 201-219.
24. Liu, J., Price, D. T., Chen, J. M. Nitrogen controls on ecosystem carbon sequestration: a model implementation and application to Saskatchewan, Canada. Ecological Modelling, 2005,186(2), 178-195.

25. Yuan, Q., Zhao, D., Wu, S., Dai, E. Validation of the Integrated Biosphere Simulator in simulating the potential natural vegetation map of China. Ecological research, 2011,26(5), 917-929.
26. Liu, X., Guo, Q., Wang, C. Simulating net primary production and soil-surface CO2 flux of temperate forests in Northeastern China. Scandinavian journal of forest research, 2011,26(1), 30-39.
27. Luo, G., Yin, G. Net Primary Productivity of Farmland Ecosystem in Xinjiang: The Spatio-temporal Dynamics and Its Response to Climate Change Based on Agro-IBIS Model. Chinese Agricultural Science Bulletin, 2018,(34), 91-98.
28. Cao, X., Zhou, Z., Chen, X., Shao, W., Wang, Z. Improving leaf area index simulation of IBIS model and its effect on water carbon and energy-A case study in Changbai Mountain broadleaved forest of China. Ecological modelling, 2015,303, 97-104.
29. Yang, Y., Zhao, P., Zhao, J., Wang, H., Wang, B., Su, S., Peng, C. Trait-based climate change predictions of vegetation sensitivity and distribution in China. Frontiers in plant science, 2019,10, 908.
30. Cheng NN, He HM, Lu YJ, Yang SY. Coupling Analysis of Hydrometeorology and Erosive Landforms Evolution in Loess Plateau, China. Adv Meteorol, 2016, 2016.
31. Dong, M, Ye, X. H., Chu, Y. et al. Ecosystem Positioning Observation and Research Dataset of China. Grassland and desert ecosystems volume. Inner Mongolia Erdos Station:2004~2006 [M]. Beijing: China Agricultural Publishing House, 2011.
32. Liu, W. Z., Dang T. H. Chinese Ecosystem Positioning Observation and Research Dataset-Farmland Ecosystem Volume-Changwu Station, Shaanxi (1998-2008) [M]. Beijing: China Agricultural Publishing House, 2012a.
33. Liu, G. B., Li Q. X., Chen Y.M. China Ecosystem Positioning Observation and Research Dataset/Farmland Ecosystem-Shaanxi Ansai Station [M]. Beijing: China Agricultural Press, 2012b.
34. Li, X. R., Zhou H. Y., Wang X. P. China Ecosystem Positioning Observation and Research Dataset/Grassland and Desert Ecosystems-Ningxia Shapotou Station [M]. Beijing: China Agricultural Press, 2010.
35. Kucharik, C. J., Foley, J. A., Delire, C., Fisher, V. A., Coe, M. T., Lenters, J. D., Gower, S. T. Testing the performance of a dynamic global ecosystem model: water balance, carbon balance, and vegetation structure. Global Biogeochemical Cycles, 2000,14(3), 795-825.
36. Farquhar, G. D., Caemmerer, S. V., Berry, J. A. A biochemical model of photosynthetic CO_2 assimilation in leaves of C 3 species. Planta, 1980,149(1), 78-90.
37. Farquhar, G. D., Sharkey, T. D. Stomatal conductance and photosynthesis. Annual review of plant physiology, 1982,33(1), 317-345.
38. Amthor, J. The role of maintenance respiration in plant growth.Plant, Cell & Environment, 1984,7(8), 561-569.
39. Xiao, Y., Zhao, L., Dai, Y., Li, R., Pang, Q., & Yao, J. Representing permafrost properties in CoLM for the Qinghai–Xizang (Tibetan) Plateau. Cold Regions Science and Technology, 2013,87, 68-77.
40. Lamparter A, Bachmann J, Goebel MO, Woche SK. Carbon mineralization in soil: Impact of wetting-drying, aggregation and water repellency. Geoderma, 2009, 150:324-333.
41. Fischer, G. R., Costa, M. H., Murta, F. Z., Malhado, A. C., Aguiar, L. J., Ladle, R. J. Multi-site land surface model optimization: An exploration of objective functions. Agricultural and forest meteorology, 2013,182, 168-176.
42. Chang, K., Price, D., Chen, J. M., Kurz, W., Boisvenue, C., Hogg, E., Hember, R. Simulating impacts of water stress on woody biomass in the southern boreal region of western Canada using a dynamic vegetation model. Agricultural and forest meteorology, 2014,198, 142-154.
43. Zhou, Y., Sun, X., Zhu, Z., Zhang, R., Tian, J., Liu, Y., Yuan, G. Surface roughness length dynamic over several different surfaces and its effects on modeling fluxes. Science in China Series D: Earth Sciences, 2006,49(2), 262-272.
44. Reich, P. B., Walters, M. B., Ellsworth, D. S. From tropics to tundra: global convergence in plant functioning.Proceedings of the National Academy of Sciences, 94(25), 1997,13730-13734.
45. Weiher, E., Van Der Werf, A., Thompson, K., Roderick, M., Garnier, E., & Eriksson, O. Challenging Theophrastus: a common core list of plant traits for functional ecology. Journal of vegetation science, 1999, 10(5), 609-620.
46. Tian, G., Zhu, Z., Li.,Y. Effects of clipping, fertilizing,and watering on compensatory growth of Elymusnutans. Chinese Journal of Ecology, 2010,29(05), 869-875.
47. Qi, W., Guo, S., Cui, X., Yang, M., Zhang, Y., Du, G., Bu, H. Variation of seed mass and SLA of 63 plant species in different habitats at various altitudes on the eastern Loess Plateau.ACTA PRATACULTURAE SINICA, 2012,21(6), 42-50.
48. Willmott, C. J. Some comments on the evaluation of model performance. Bulletin of the American Meteorological Society,1982,63(11), 1309-1313.
49. Yang, Y., Zhu, Q., Peng, C., Wang, H., Chen, H. From plant functional types to plant functional traits: A new paradigm in modelling global vegetation dynamics. Progress in Physical Geography, 2015,39(4), 514-535.
50. Yu, M., Wu, B., Zeng, H., Xing, Q., Zhu, W. The Impacts of Vegetation and Meteorological Factors on Aerodynamic Roughness Length at Different Time Scales. Atmosphere, 2018,9(4), 149.
51. Vendramini, F., Díaz, S., Gurvich, D. E., Wilson, P. J., Thompson, K., Hodgson, J. G. Leaf traits as indicators of resource-use strategy in floras with succulent species. New Phytologist, 2002,154(1), 147-157.
52. Park, H., Iijima, Y., Yabuki, H., Ohta, T., Walsh, J., Kodama, Y., Ohata, T. The application of a coupled hydrological and biogeochemical model (CHANGE) for modeling of energy, water, and CO_2 exchanges over a larch forest in eastern Siberia. Journal of Geophysical Research: Atmospheres, 2011,116(D15).
53. Zhao, L., Wu, X., Wang, Z., Sheng, Y., Fang, H., Zhao, Y., Mo, B. Soil organic carbon and total nitrogen pools in arid region zones of the Loess Plateau. Scientific reports, 2018, 8(1), 3656.

54. Wang, J., Wang, G., Hu, H., & Wu, Q. The influence of degradation of the swamp and alpine meadows on CH_4 and CO_2 fluxes on the Loess Plateau. Environmental Earth Sciences,2010,60(3), 537-548.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.